



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:33 AM UTC

PDB ID : 5GL5 / pdb_00005gl5
Title : Sterol 3-beta-glucosyltransferase (ugt51) from *Saccharomyces cerevisiae*
(strain ATCC 204508 / S288c): UDPG complex
Authors : Feng, Y.; Chen, L.-Q.
Deposited on : 2016-07-08
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	498	8% 82% 5% 13%
1	B	498	5% 82% 5% 13%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7204 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sterol 3-beta-glucosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	431	Total	C	N	O	S	0	1	0
			3427	2211	585	620	11			
1	B	431	Total	C	N	O	S	0	1	0
			3419	2206	583	619	11			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	701	MET	-	expression tag	UNP Q06321
A	702	GLY	-	expression tag	UNP Q06321
A	703	SER	-	expression tag	UNP Q06321
A	704	SER	-	expression tag	UNP Q06321
A	705	HIS	-	expression tag	UNP Q06321
A	706	HIS	-	expression tag	UNP Q06321
A	707	HIS	-	expression tag	UNP Q06321
A	708	HIS	-	expression tag	UNP Q06321
A	709	HIS	-	expression tag	UNP Q06321
A	710	HIS	-	expression tag	UNP Q06321
A	711	SER	-	expression tag	UNP Q06321
A	712	SER	-	expression tag	UNP Q06321
A	713	GLY	-	expression tag	UNP Q06321
A	714	LEU	-	expression tag	UNP Q06321
A	715	VAL	-	expression tag	UNP Q06321
A	716	PRO	-	expression tag	UNP Q06321
A	717	ARG	-	expression tag	UNP Q06321
A	718	GLY	-	expression tag	UNP Q06321
A	719	SER	-	expression tag	UNP Q06321
A	720	HIS	-	expression tag	UNP Q06321
A	721	MET	-	expression tag	UNP Q06321
B	701	MET	-	expression tag	UNP Q06321
B	702	GLY	-	expression tag	UNP Q06321
B	703	SER	-	expression tag	UNP Q06321
B	704	SER	-	expression tag	UNP Q06321

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	705	HIS	-	expression tag	UNP Q06321
B	706	HIS	-	expression tag	UNP Q06321
B	707	HIS	-	expression tag	UNP Q06321
B	708	HIS	-	expression tag	UNP Q06321
B	709	HIS	-	expression tag	UNP Q06321
B	710	HIS	-	expression tag	UNP Q06321
B	711	SER	-	expression tag	UNP Q06321
B	712	SER	-	expression tag	UNP Q06321
B	713	GLY	-	expression tag	UNP Q06321
B	714	LEU	-	expression tag	UNP Q06321
B	715	VAL	-	expression tag	UNP Q06321
B	716	PRO	-	expression tag	UNP Q06321
B	717	ARG	-	expression tag	UNP Q06321
B	718	GLY	-	expression tag	UNP Q06321
B	719	SER	-	expression tag	UNP Q06321
B	720	HIS	-	expression tag	UNP Q06321
B	721	MET	-	expression tag	UNP Q06321

- Molecule 2 is URIDINE-5'-DIPHOSPHATE-GLUCOSE (CCD ID: UPG) (formula: $C_{15}H_{24}N_2O_{17}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			36	15	2	17	2		
2	B	1	Total	C	N	O	P	0	0
			36	15	2	17	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	133	Total 133	O 133	0	0
3	B	153	Total 153	O 153	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Sterol 3-beta-glucosyltransferase



• Molecule 1: Sterol 3-beta-glucosyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.66Å 80.56Å 154.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	77.29 – 1.90 77.29 – 1.90	Depositor EDS
% Data completeness (in resolution range)	89.5 (77.29-1.90) 89.5 (77.29-1.90)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.16 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.8.0157	Depositor
R, R_{free}	0.185 , 0.219 (Not available) , 0.226	Depositor DCC
R_{free} test set	3095 reflections (4.48%)	wwPDB-VP
Wilson B-factor (Å ²)	19.4	Xtriage
Anisotropy	0.196	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 41.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7204	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UPG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	1/3507 (0.0%)	0.97	3/4744 (0.1%)
1	B	1.00	1/3500 (0.0%)	1.01	5/4737 (0.1%)
All	All	0.99	2/7007 (0.0%)	0.99	8/9481 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1039	LYS	CA-C	6.22	1.56	1.52
1	B	804	VAL	CA-CB	5.53	1.60	1.54

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1034	ASP	N-CA-C	8.76	123.41	107.99
1	B	935	ASN	CB-CA-C	-7.30	95.06	110.31
1	A	947	PHE	CA-C-N	-6.47	115.00	119.66
1	A	947	PHE	C-N-CA	-6.47	115.00	119.66
1	A	739	SER	N-CA-C	5.64	117.27	109.15
1	B	947	PHE	CA-C-N	-5.62	114.60	120.38
1	B	947	PHE	C-N-CA	-5.62	114.60	120.38
1	B	811	VAL	CB-CA-C	-5.07	105.48	111.97

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	1033	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3427	0	3455	11	0
1	B	3419	0	3442	11	0
2	A	36	0	22	0	0
2	B	36	0	22	0	0
3	A	133	0	0	0	0
3	B	153	0	0	0	0
All	All	7204	0	6941	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:802:LEU:HD21	1:A:806:ASN:HD22	1.60	0.66
1:A:1083:LEU:O	1:A:1085:THR:HG23	1.97	0.64
1:A:941:ASN:HB3	1:A:963:TRP:CE2	2.43	0.54
1:B:826:ASP:OD1	1:B:912:GLN:NE2	2.41	0.53
1:A:1015:VAL:O	1:A:1018:ALA:O	2.26	0.53
1:B:946:ILE:HD12	1:B:1104:ILE:HD13	1.93	0.51
1:B:941:ASN:HB3	1:B:963:TRP:CE2	2.47	0.49
1:A:738:LYS:O	1:A:768:HIS:HE1	1.96	0.49
1:B:1016:MET:CE	1:B:1047:ASN:HD21	2.27	0.47
1:A:950:SER:HB3	1:A:953:PHE:CD1	2.50	0.47
1:B:813:MET:N	1:B:813:MET:HE2	2.31	0.46
1:B:776:SER:OG	1:B:791:GLU:CD	2.59	0.46
1:B:806:ASN:HB3	1:B:807:GLU:HG2	1.99	0.44
1:B:946:ILE:HD13	1:B:1079:LEU:HD11	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:738:LYS:O	1:A:768:HIS:CE1	2.72	0.41
1:A:895:TYR:CD1	1:B:809:MET:HG2	2.55	0.41
1:B:809:MET:CE	1:B:814:LEU:N	2.83	0.41
1:A:814:LEU:HD13	1:B:899:VAL:HG21	2.02	0.41
1:A:946:ILE:HD13	1:A:1079:LEU:HD11	2.03	0.41
1:A:990:LYS:HB2	1:A:1020:VAL:HG22	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	428/498 (86%)	415 (97%)	13 (3%)	0	100	100
1	B	428/498 (86%)	418 (98%)	10 (2%)	0	100	100
All	All	856/996 (86%)	833 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	UPG	A	1201	-	37,38,38	0.98	3 (8%)	55,58,58	1.42	6 (10%)
2	UPG	B	1201	-	37,38,38	0.97	3 (8%)	55,58,58	1.36	6 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UPG	A	1201	-	-	2/23/59/59	0/3/3/3
2	UPG	B	1201	-	-	3/23/59/59	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1201	UPG	C4-N3	-2.59	1.34	1.38
2	B	1201	UPG	O2-C2	2.57	1.27	1.23
2	A	1201	UPG	C2-N1	2.39	1.42	1.38
2	B	1201	UPG	PA-O3A	-2.23	1.57	1.59
2	B	1201	UPG	PB-O3A	-2.13	1.57	1.59
2	A	1201	UPG	C6-C5	2.07	1.39	1.35

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1201	UPG	O2B-PB-O3A	-4.45	95.25	107.27
2	A	1201	UPG	C5-C4-N3	4.10	120.55	114.80
2	A	1201	UPG	C4-N3-C2	-3.76	121.95	126.61
2	A	1201	UPG	O3A-PB-O1B	-3.62	99.82	110.70
2	A	1201	UPG	N3-C2-N1	3.36	119.27	114.89
2	B	1201	UPG	O4-C4-C5	-3.21	119.62	125.16
2	B	1201	UPG	C5-C4-N3	3.13	119.18	114.80
2	B	1201	UPG	O2B-PB-O1B	2.92	126.02	112.44
2	A	1201	UPG	O4-C4-C5	-2.65	120.59	125.16
2	B	1201	UPG	O2A-PA-O1A	2.42	123.68	112.44
2	B	1201	UPG	N3-C2-N1	2.03	117.54	114.89
2	A	1201	UPG	O2B-PB-O1B	2.00	121.75	112.44

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1201	UPG	C1'-O3B-PB-O1B
2	B	1201	UPG	C1'-O3B-PB-O2B
2	B	1201	UPG	C1'-O3B-PB-O1B
2	A	1201	UPG	PB-O3A-PA-O2A
2	B	1201	UPG	C2'-C1'-O3B-PB

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand UPG A 1201



Ligand UPG B 1201



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	431/498 (86%)	0.42	40 (9%) 14 15	9, 24, 75, 123	1 (0%)
1	B	431/498 (86%)	0.22	23 (5%) 32 34	9, 23, 57, 100	1 (0%)
All	All	862/996 (86%)	0.32	63 (7%) 21 22	9, 24, 69, 123	2 (0%)

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1040	THR	6.6
1	A	802	LEU	5.7
1	A	1042	VAL	5.3
1	B	1033	GLY	5.2
1	B	1032	LEU	5.2
1	A	1032	LEU	5.2
1	B	803	MET	4.6
1	A	1033	GLY	4.6
1	B	804	VAL	4.5
1	A	1039	LYS	4.3
1	B	1036	ALA	4.2
1	A	803	MET	4.1
1	B	802	LEU	4.1
1	A	811	VAL	4.0
1	A	1038	LYS	4.0
1	A	809	MET	3.9
1	A	799	LEU	3.9
1	A	968	LYS	3.8
1	B	1091	PHE	3.7
1	A	1091	PHE	3.4
1	B	1035	LYS	3.3
1	A	806	ASN	3.3
1	B	1031	ARG	3.3
1	A	1104	ILE	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	969	SER	3.1
1	A	800	MET	3.0
1	A	804	VAL	3.0
1	A	1051	ILE	2.9
1	B	1034	ASP	2.9
1	A	970	THR	2.9
1	B	800	MET	2.9
1	A	1018	ALA	2.9
1	B	806	ASN	2.8
1	A	979	GLU	2.8
1	A	1002	SER	2.8
1	B	1030	GLU	2.8
1	A	1103	ASP	2.8
1	A	1003	ASN	2.7
1	B	1111	LYS	2.6
1	B	1011	LEU	2.6
1	A	1041	GLU	2.6
1	A	1171	LYS	2.6
1	B	1038	LYS	2.5
1	A	1130	ILE	2.5
1	B	1051	ILE	2.5
1	A	805	GLU	2.4
1	B	801	SER	2.4
1	A	1043	ASP	2.4
1	B	805	GLU	2.3
1	B	1037	ALA	2.2
1	B	797	VAL	2.2
1	A	1127	THR	2.2
1	A	1016	MET	2.1
1	B	1043	ASP	2.1
1	A	1005	LYS	2.1
1	A	1111	LYS	2.1
1	A	1019	ASP	2.1
1	A	1031	ARG	2.1
1	A	737	ASN	2.1
1	A	801	SER	2.0
1	B	1129	LYS	2.0
1	A	814	LEU	2.0
1	A	1123	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	UPG	B	1201	36/36	0.95	0.07	13,24,44,46	0
2	UPG	A	1201	36/36	0.96	0.06	17,24,32,37	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around UPG B 1201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around UPG A 1201:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.