



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 11:03 PM UTC

PDB ID : 8GLG / pdb_00008glg
Title : Crystal Structure of Human CD1b in Complex with Phosphatidylethanolamine C34:1
Authors : Shahine, A.
Deposited on : 2023-03-22
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	300	2% 85% 8% 7%
2	B	101	5% 88% 11% .
3	C	7	57% 43%
4	D	4	100%

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 3851 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called T-cell surface glycoprotein CD1b.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	280	Total	C	N	O	S	0	8	0
			2232	1431	376	415	10			

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	279	GLY	-	expression tag	UNP P29016
A	280	SER	-	expression tag	UNP P29016
A	281	GLY	-	expression tag	UNP P29016
A	282	LEU	-	expression tag	UNP P29016
A	283	ASN	-	expression tag	UNP P29016
A	284	ASP	-	expression tag	UNP P29016
A	285	ILE	-	expression tag	UNP P29016
A	286	PHE	-	expression tag	UNP P29016
A	287	GLU	-	expression tag	UNP P29016
A	288	ALA	-	expression tag	UNP P29016
A	289	GLN	-	expression tag	UNP P29016
A	290	LYS	-	expression tag	UNP P29016
A	291	ILE	-	expression tag	UNP P29016
A	292	GLU	-	expression tag	UNP P29016
A	293	TRP	-	expression tag	UNP P29016
A	294	HIS	-	expression tag	UNP P29016
A	295	GLU	-	expression tag	UNP P29016
A	296	HIS	-	expression tag	UNP P29016
A	297	HIS	-	expression tag	UNP P29016
A	298	HIS	-	expression tag	UNP P29016
A	299	HIS	-	expression tag	UNP P29016
A	300	HIS	-	expression tag	UNP P29016
A	301	HIS	-	expression tag	UNP P29016


- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	5	0
			842	537	142	161	2			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	1	PRO	-	expression tag	UNP P61769
B	2	LYS	-	expression tag	UNP P61769

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)][alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	7	Total	C	N	O		0	0	0
			81	46	2	33				

- Molecule 4 is an oligosaccharide called alpha-L-fucopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)][alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	4	Total	C	N	O		0	0	0
			48	28	2	18				

- Molecule 5 is (1S)-2-{[(S)-(2-aminoethoxy)(hydroxy)phosphoryl]oxy}-1-[(octadecanoyloxy)methyl]ethyl (9Z)-octadec-9-enoate (CCD ID: L9Q) (formula: C₄₁H₈₀NO₈P) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			49	39	1	8	1		

- Molecule 6 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Ni	0	0
			1	1		

- Molecule 7 is tetracosyl octadecanoate (CCD ID: CUY) (formula: C₄₂H₈₄O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			35	33	2		


- Molecule 8 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	4	Total	Cl		0	0
			4	4			

- Molecule 9 is IODIDE ION (CCD ID: IOD) (formula: I).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	I		0	0
			1	1			
9	B	1	Total	I		0	0
			1	1			

- Molecule 10 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	C	O	0	0
			4	2	2		
10	B	1	Total	C	O	0	0
			4	2	2		
10	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	391	Total	O	0	0
			391	391		
11	B	150	Total	O	0	0
			150	150		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: T-cell surface glycoprotein CD1b




- Molecule 2: Beta-2-microglobulin



- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)][alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: alpha-L-fucopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)][alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.07Å 80.53Å 92.98Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.95 – 1.60 36.95 – 1.60	Depositor EDS
% Data completeness (in resolution range)	98.6 (36.95-1.60) 98.5 (36.95-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.21 (at 1.60Å)	Xtriage
Refinement program	BUSTER	Depositor
R, R_{free}	0.186 , 0.209 0.191 , 0.211	Depositor DCC
R_{free} test set	2854 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	17.4	Xtriage
Anisotropy	0.186	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 46.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3851	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, EDO, CL, L9Q, IOD, NAG, NI, FUC, MAN, CUY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	5/2306 (0.2%)	1.02	4/3133 (0.1%)
2	B	0.73	0/872	0.97	3/1188 (0.3%)
All	All	0.85	5/3178 (0.2%)	1.00	7/4321 (0.2%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	96	ILE	C-O	-6.52	1.17	1.24
1	A	31	TRP	C-O	-6.08	1.16	1.23
1	A	97	GLN	C-O	-5.80	1.16	1.23
1	A	222	ARG	C-O	-5.65	1.17	1.24
1	A	169	TYR	C-O	-5.25	1.18	1.24

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	43	ASP	CA-CB-CG	6.79	119.39	112.60
2	B	23	ASN	CA-CB-CG	6.20	118.80	112.60
1	A	68	GLU	CB-CG-CD	6.12	123.01	112.60
2	B	72	PHE	CA-CB-CG	5.57	119.37	113.80
2	B	35	SER	N-CA-C	5.13	117.78	111.82
1	A	282	LEU	CA-C-N	5.02	130.73	121.70
1	A	282	LEU	C-N-CA	5.02	130.73	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2232	0	2142	17	0
2	B	842	0	787	7	0
3	C	81	0	70	0	0
4	D	48	0	43	0	0
5	A	49	0	72	3	0
6	A	1	0	0	0	0
7	A	35	0	0	0	0
8	A	4	0	0	2	0
9	A	1	0	0	0	0
9	B	1	0	0	0	0
10	A	4	0	6	0	0
10	B	12	0	17	6	0
11	A	391	0	0	6	0
11	B	150	0	0	0	0
All	All	3851	0	3137	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:228:GLN:HG3	11:A:846:HOH:O	1.58	1.03
1:A:119:GLY:HA3	2:B:1:PRO:HA	1.40	1.01
8:A:604:CL:CL	11:A:868:HOH:O	2.21	0.94
8:A:605:CL:CL	11:A:846:HOH:O	2.28	0.87
2:B:67:LEU:HD23	10:B:203:EDO:H11	1.65	0.78
1:A:68:GLU:HG3	5:A:601:L9Q:N	2.09	0.67
2:B:34:PRO:CB	10:B:202:EDO:H11	2.28	0.64
1:A:228:GLN:CG	11:A:846:HOH:O	2.32	0.63
1:A:219:MET:CE	11:A:844:HOH:O	2.48	0.61
1:A:68:GLU:HG3	5:A:601:L9Q:HN	1.69	0.58
1:A:187:PRO:HB3	1:A:211:PHE:HB3	1.88	0.55
1:A:252:ALA:HB3	1:A:255:GLU:HG3	1.92	0.52
2:B:31:GLY:HA2	2:B:63[A]:SER:HB3	1.94	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:67:LEU:CD2	10:B:203:EDO:H11	2.39	0.49
1:A:219:MET:HE1	11:A:844:HOH:O	2.12	0.47
2:B:54:SER:HB3	10:B:203:EDO:H12	1.96	0.47
1:A:93:PRO:HB2	10:B:202:EDO:H21	1.97	0.46
2:B:34:PRO:HB2	10:B:202:EDO:H11	2.00	0.44
1:A:208:VAL:HG21	1:A:218[B]:VAL:HG11	2.00	0.44
1:A:66:LEU:HD13	5:A:601:L9Q:H46	2.00	0.44
1:A:235:ILE:HD13	1:A:245:LEU:HB2	2.01	0.43
1:A:208:VAL:HG21	1:A:218[B]:VAL:CG1	2.49	0.42
1:A:217:TRP:HB3	1:A:264:LYS:HB2	2.01	0.42
1:A:124:LEU:HD12	1:A:124:LEU:C	2.47	0.40
1:A:218[B]:VAL:HG23	1:A:218[B]:VAL:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	286/300 (95%)	283 (99%)	3 (1%)	0	100	100
2	B	103/101 (102%)	103 (100%)	0	0	100	100
All	All	389/401 (97%)	386 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/247 (96%)	235 (100%)	1 (0%)	84	75
2	B	94/96 (98%)	92 (98%)	2 (2%)	47	23
All	All	330/343 (96%)	327 (99%)	3 (1%)	70	55

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	269	GLU
2	B	3	ILE
2	B	6	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	GLN
1	A	184	GLN
1	A	283	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

11 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	C	1	1,3	14,14,15	0.41	0	17,19,21	0.97	1 (5%)
3	NAG	C	2	3	14,14,15	0.31	0	17,19,21	0.55	0
3	BMA	C	3	3	11,11,12	0.33	0	15,15,17	0.64	1 (6%)
3	MAN	C	4	3	11,11,12	0.35	0	15,15,17	0.63	0
3	MAN	C	5	3	11,11,12	0.40	0	15,15,17	0.91	1 (6%)
3	FUC	C	6	3	10,10,11	0.38	0	14,14,16	0.53	0
3	FUC	C	7	3	10,10,11	0.45	0	14,14,16	0.66	0
4	NAG	D	1	1,4	14,14,15	0.32	0	17,19,21	0.63	0
4	FUC	D	2	4	10,10,11	0.39	0	14,14,16	0.65	0
4	NAG	D	3	4	14,14,15	0.30	0	17,19,21	0.46	0
4	FUC	D	4	4	10,10,11	0.39	0	14,14,16	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
3	BMA	C	3	3	-	0/2/19/22	0/1/1/1
3	MAN	C	4	3	-	0/2/19/22	0/1/1/1
3	MAN	C	5	3	-	0/2/19/22	0/1/1/1
3	FUC	C	6	3	-	-	0/1/1/1
3	FUC	C	7	3	-	-	0/1/1/1
4	NAG	D	1	1,4	-	0/6/23/26	0/1/1/1
4	FUC	D	2	4	-	-	0/1/1/1
4	NAG	D	3	4	-	0/6/23/26	0/1/1/1
4	FUC	D	4	4	-	-	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	5	MAN	C1-O5-C5	3.20	116.48	112.19
3	C	1	NAG	C1-O5-C5	2.77	115.90	112.19
3	C	3	BMA	C1-O5-C5	2.02	114.89	112.19

There are no chirality outliers.


All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	1	NAG	C3-C2-N2-C7
3	C	1	NAG	C1-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 7 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	CUY	A	603	-	34,34,43	0.64	1 (2%)	34,34,43	0.69	1 (2%)
10	EDO	B	204	-	3,3,3	0.47	0	2,2,2	0.39	0
10	EDO	B	203	-	3,3,3	0.35	0	2,2,2	0.33	0
10	EDO	A	609	-	3,3,3	0.50	0	2,2,2	0.41	0
10	EDO	B	202	-	3,3,3	1.27	1 (33%)	2,2,2	0.61	0
5	L9Q	A	601	-	48,48,50	0.95	2 (4%)	51,53,55	1.13	5 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	CUY	A	603	-	-	16/33/33/42	-
10	EDO	B	204	-	-	1/1/1/1	-
10	EDO	B	203	-	-	0/1/1/1	-
10	EDO	A	609	-	-	0/1/1/1	-
10	EDO	B	202	-	-	1/1/1/1	-
5	L9Q	A	601	-	-	26/52/52/54	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	601	L9Q	O2-C31	2.65	1.41	1.34
7	A	603	CUY	OAQ-CAP	2.56	1.40	1.33
5	A	601	L9Q	O3-C11	2.16	1.39	1.33
10	B	202	EDO	O2-C2	-2.09	1.31	1.42

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	601	L9Q	O2-C31-C32	4.92	122.13	111.48
5	A	601	L9Q	O3-C11-C12	2.73	120.15	111.83
7	A	603	CUY	OAQ-CAP-CAO	2.54	119.58	111.83
5	A	601	L9Q	O2-C31-O31	-2.48	117.92	123.70
5	A	601	L9Q	C2-O2-C31	2.34	123.40	117.80
5	A	601	L9Q	O3-C11-O11	-2.24	118.02	123.63

There are no chirality outliers.

All (44) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	601	L9Q	C3-C2-O2-C31
5	A	601	L9Q	O31-C31-O2-C2
5	A	601	L9Q	C32-C31-O2-C2
5	A	601	L9Q	O4P-C4-C5-N
5	A	601	L9Q	C12-C11-O3-C3
5	A	601	L9Q	O11-C11-O3-C3
5	A	601	L9Q	C31-C32-C33-C34
7	A	603	CUY	OAQ-CAR-CAS-CAT
7	A	603	CUY	C1-CBD-CBE-CBF
7	A	603	CUY	CBD-CBE-CBF-CBG
5	A	601	L9Q	C17-C18-C19-C20
5	A	601	L9Q	C42-C43-C44-C45
5	A	601	L9Q	C41-C42-C43-C44
7	A	603	CUY	CBJ-CBK-CBL-CBM
5	A	601	L9Q	C35-C36-C37-C38
7	A	603	CUY	CAG-CAH-CAI-CAJ
5	A	601	L9Q	C20-C21-C22-C23
5	A	601	L9Q	C40-C41-C42-C43
7	A	603	CUY	CBF-CBG-CBH-CBI
5	A	601	L9Q	C16-C17-C18-C19
7	A	603	CUY	CAU-CAV-CAW-CBM
7	A	603	CUY	CAT-CAU-CAV-CAW
7	A	603	CUY	CAO-CAP-OAQ-CAR
7	A	603	CUY	CAR-CAS-CAT-CAU
7	A	603	CUY	CBG-CBH-CBI-CBJ
5	A	601	L9Q	C33-C34-C35-C36
7	A	603	CUY	CAF-CAG-CAH-CAI
7	A	603	CUY	OBO-CAP-OAQ-CAR
5	A	601	L9Q	C13-C14-C15-C16
5	A	601	L9Q	C34-C35-C36-C37
5	A	601	L9Q	C12-C13-C14-C15
5	A	601	L9Q	C14-C15-C16-C17
5	A	601	L9Q	C15-C16-C17-C18
7	A	603	CUY	CBH-CBI-CBJ-CBK
7	A	603	CUY	CAH-CAI-CAJ-CAK
5	A	601	L9Q	C36-C37-C38-C39
5	A	601	L9Q	C43-C44-C45-C46
5	A	601	L9Q	C2-C1-O3P-P
10	B	204	EDO	O1-C1-C2-O2
5	A	601	L9Q	C11-C12-C13-C14
7	A	603	CUY	CBI-CBJ-CBK-CBL
10	B	202	EDO	O1-C1-C2-O2
5	A	601	L9Q	C37-C38-C39-C40

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	A	601	L9Q	O3-C11-C12-C13


There are no ring outliers.

3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	B	203	EDO	3	0
10	B	202	EDO	3	0
5	A	601	L9Q	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	280/300 (93%)	-0.08	5 (1%) 67 71	7, 19, 39, 63	8 (2%)
2	B	100/101 (99%)	0.12	5 (5%) 34 35	7, 25, 49, 72	5 (5%)
All	All	380/401 (94%)	-0.03	10 (2%) 57 60	7, 20, 43, 72	13 (3%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	1	PRO	7.3
2	B	3	ILE	4.5
1	A	31	TRP	3.6
2	B	2	LYS	3.3
1	A	283	ASN	3.2
2	B	100	ASP	3.0
1	A	193	SER	2.6
2	B	73[A]	THR	2.6
1	A	152	GLN	2.3
1	A	279	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	FUC	D	2	10/11	0.58	0.15	60,66,67,67	0
4	NAG	D	3	14/15	0.62	0.16	54,57,60,62	0
4	FUC	D	4	10/11	0.78	0.13	50,53,55,55	0
4	NAG	D	1	14/15	0.85	0.11	39,45,49,55	0
3	FUC	C	6	10/11	0.90	0.10	23,28,31,31	0
3	FUC	C	7	10/11	0.92	0.08	20,24,26,26	0
3	MAN	C	4	11/12	0.93	0.08	23,25,29,29	0
3	NAG	C	2	14/15	0.94	0.07	16,18,20,25	0
3	MAN	C	5	11/12	0.95	0.06	16,17,20,21	0
3	NAG	C	1	14/15	0.96	0.06	14,15,17,18	0
3	BMA	C	3	11/12	0.96	0.06	16,17,20,20	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
10	EDO	A	609	4/4	0.75	0.15	50,52,54,55	0
10	EDO	B	204	4/4	0.79	0.16	40,42,43,44	0
5	L9Q	A	601	49/51	0.81	0.17	31,43,54,58	0
10	EDO	B	203	4/4	0.86	0.16	34,35,39,43	0
7	CUY	A	603	35/44	0.86	0.14	26,31,36,37	0

Continued on next page...




Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
9	IOD	B	201	1/1	0.91	0.09	30,30,30,30	1
10	EDO	B	202	4/4	0.94	0.16	12,17,27,33	0
8	CL	A	606	1/1	0.95	0.09	28,28,28,28	0
8	CL	A	607	1/1	0.96	0.11	12,12,12,12	0
6	NI	A	602	1/1	0.97	0.05	25,25,25,25	0
8	CL	A	605	1/1	0.98	0.05	21,21,21,21	0
9	IOD	A	608	1/1	0.99	0.03	15,15,15,15	1
8	CL	A	604	1/1	0.99	0.04	20,20,20,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around L9Q A 601:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.