



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 08:55 AM UTC

PDB ID : 3GMA / pdb_00003gma
Title : Glutaconyl-coA decarboxylase A subunit from Clostridium symbiosum co-crystallized with glutaryl-CoA
Authors : Kress, D.; Brugel, D.; Buckel, W.; Essen, L.-O.
Deposited on : 2009-03-13
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	588	5% 86% 9% 5%
1	B	588	7% 88% 8% • •

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 9170 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutaconyl-CoA decarboxylase subunit A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	558	Total	C	N	O	S	0	1	0
			4330	2751	730	821	28			
1	B	566	Total	C	N	O	S	0	1	0
			4376	2777	736	835	28			

- Molecule 2 is glutaryl-coenzyme A (CCD ID: GRA) (formula: $C_{26}H_{42}N_7O_{19}P_3S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P S	0	0
			56	26	7	19	3 1		
2	B	1	Total	C	N	O	P S	0	0
			56	26	7	19	3 1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	175	Total 175	O 175	0	0
3	B	177	Total 177	O 177	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutaconyl-CoA decarboxylase subunit A



- Molecule 1: Glutaconyl-CoA decarboxylase subunit A



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	156.47Å 156.47Å 141.40Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	68.52 – 2.60 68.52 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.5 (68.52-2.60) 99.8 (68.52-2.60)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.87 (at 2.58Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.189 , 0.209 0.187 , 0.183	Depositor DCC
R_{free} test set	674 reflections (1.09%)	wwPDB-VP
Wilson B-factor (Å ²)	38.2	Xtriage
Anisotropy	0.022	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 37.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.022 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9170	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GRA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	0/4428	0.82	0/6004
1	B	0.56	1/4474 (0.0%)	0.81	0/6069
All	All	0.55	1/8902 (0.0%)	0.81	0/12073

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	381	ASN	CA-C	-5.07	1.48	1.53

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4330	0	4275	30	0
1	B	4376	0	4294	29	0
2	A	56	0	37	2	0
2	B	56	0	37	2	0
3	A	175	0	0	1	0
3	B	177	0	0	2	0
All	All	9170	0	8643	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:374:ASN:HB2	3:B:706:HOH:O	1.52	1.07
1:A:27:ASN:HD21	1:A:318:MET:H	1.19	0.86
1:B:27:ASN:HD21	1:B:318:MET:H	1.21	0.85
1:A:150:CYS:H	1:A:193:ASN:ND2	1.81	0.77
1:B:150:CYS:H	1:B:193:ASN:ND2	1.83	0.76
1:A:67:ARG:HH12	1:A:149:ASN:HD22	1.35	0.74
1:B:67:ARG:HH12	1:B:149:ASN:HD22	1.36	0.73
1:A:273:GLN:HB3	3:A:723:HOH:O	1.93	0.67
1:A:367:ASN:ND2	1:A:416:ASP:H	1.94	0.64
1:B:367:ASN:ND2	1:B:416:ASP:H	1.95	0.63
1:A:150:CYS:H	1:A:193:ASN:HD21	1.45	0.63
1:A:63:SER:H	1:A:66:GLN:HE21	1.47	0.61
1:B:63:SER:H	1:B:66:GLN:HE21	1.47	0.60
1:B:150:CYS:H	1:B:193:ASN:HD21	1.49	0.59
1:B:367:ASN:HD22	1:B:416:ASP:H	1.48	0.59
1:B:195:ALA:HB1	2:B:589:GRA:H8B	1.84	0.58
1:A:367:ASN:HD22	1:A:416:ASP:H	1.50	0.58
1:A:426:ALA:HB1	1:A:431:LEU:HD22	1.85	0.57
1:B:426:ALA:HB1	1:B:431:LEU:HD22	1.86	0.57
1:B:63:SER:H	1:B:66:GLN:NE2	2.03	0.57
1:A:259:PRO:O	1:A:263:ASP:HB2	2.04	0.56
1:A:63:SER:H	1:A:66:GLN:NE2	2.03	0.56
1:A:72:ILE:CD1	1:A:77:TRP:HB2	2.36	0.55
1:A:195:ALA:HB1	2:A:589:GRA:H8B	1.89	0.55
1:B:72:ILE:CD1	1:B:77:TRP:HB2	2.38	0.53
1:A:199:TYR:HA	1:A:202:ILE:HG22	1.93	0.50
1:B:259:PRO:O	1:B:263:ASP:HB2	2.12	0.50
1:B:199:TYR:HA	1:B:202:ILE:HG22	1.92	0.50
1:B:375:TYR:CG	1:B:376:PRO:HA	2.49	0.48
1:A:375:TYR:CG	1:A:376:PRO:HA	2.48	0.48
1:B:67:ARG:NH1	1:B:149:ASN:HD22	2.09	0.48
1:A:187:VAL:HB	1:A:204:PRO:HG2	1.97	0.47
1:A:525:MET:HG2	1:A:529:LYS:HD2	1.96	0.46
1:B:154:GLU:HA	2:B:589:GRA:N1A	2.30	0.46
1:B:3:MET:N	3:B:704:HOH:O	2.50	0.45
1:A:203:SER:N	1:A:204:PRO:CD	2.79	0.45
1:B:67:ARG:HB3	1:B:147:LEU:HD13	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:94:THR:OG1	1:B:129:ASN:ND2	2.48	0.44
1:A:3:MET:HE2	1:B:270:GLU:HG2	2.00	0.44
1:A:299:VAL:HG13	1:A:571:MET:HE2	1.99	0.44
1:B:299:VAL:HG13	1:B:571:MET:HE2	2.00	0.44
1:B:455:ARG:HA	1:B:483:CYS:HA	2.00	0.43
1:A:270:GLU:HG2	1:B:3:MET:HE2	2.01	0.43
1:A:67:ARG:NH1	1:A:149:ASN:HD22	2.10	0.42
1:B:203:SER:N	1:B:204:PRO:CD	2.82	0.42
1:B:241:ILE:H	1:B:241:ILE:HG13	1.73	0.42
1:B:187:VAL:HB	1:B:204:PRO:HG2	2.02	0.41
1:A:416:ASP:CG	1:A:455:ARG:HB2	2.46	0.41
1:A:94:THR:OG1	1:A:129:ASN:ND2	2.53	0.41
1:A:154:GLU:HA	2:A:589:GRA:N1A	2.36	0.41
1:A:455:ARG:HA	1:A:483:CYS:HA	2.02	0.41
1:B:351:VAL:HG23	1:B:368:VAL:CG2	2.51	0.41
1:A:513:GLN:HA	1:A:516:ILE:HD13	2.03	0.41
1:B:371:LEU:HB3	1:B:373:MET:CE	2.51	0.40
1:A:380:GLN:O	1:A:381:ASN:C	2.63	0.40
1:A:416:ASP:OD1	1:A:455:ARG:HB2	2.21	0.40
1:A:6:MET:SD	1:A:572:HIS:HA	2.62	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	553/588 (94%)	537 (97%)	16 (3%)	0	100	100
1	B	561/588 (95%)	540 (96%)	21 (4%)	0	100	100
All	All	1114/1176 (95%)	1077 (97%)	37 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	465/491 (95%)	455 (98%)	10 (2%)	45	72
1	B	467/491 (95%)	455 (97%)	12 (3%)	40	68
All	All	932/982 (95%)	910 (98%)	22 (2%)	43	70

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	ILE
1	A	55	LYS
1	A	56	LEU
1	A	88	ASN
1	A	239	GLN
1	A	241	ILE
1	A	310	GLU
1	A	354	LEU
1	A	495	ASN
1	A	571	MET
1	B	50	ILE
1	B	55	LYS
1	B	56	LEU
1	B	88	ASN
1	B	241	ILE
1	B	307	LEU
1	B	310	GLU
1	B	354	LEU
1	B	371	LEU
1	B	381	ASN
1	B	495	ASN
1	B	571	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (29) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	GLN
1	A	27	ASN
1	A	57	ASN
1	A	66	GLN
1	A	81	ASN
1	A	89	ASN
1	A	98	ASN
1	A	149	ASN
1	A	175	ASN
1	A	193	ASN
1	A	306	GLN
1	A	367	ASN
1	A	513	GLN
1	B	11	GLN
1	B	27	ASN
1	B	57	ASN
1	B	66	GLN
1	B	81	ASN
1	B	89	ASN
1	B	98	ASN
1	B	149	ASN
1	B	175	ASN
1	B	193	ASN
1	B	200	HIS
1	B	306	GLN
1	B	367	ASN
1	B	380	GLN
1	B	436	GLN
1	B	513	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GRA	A	589	-	56,58,58	1.28	6 (10%)	79,85,85	1.93	19 (24%)
2	GRA	B	589	-	56,58,58	1.33	6 (10%)	79,85,85	2.02	17 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GRA	A	589	-	-	10/57/73/73	0/3/3/3
2	GRA	B	589	-	-	19/57/73/73	0/3/3/3

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	589	GRA	C9-C8	-5.10	1.33	1.52
2	A	589	GRA	C9-C8	-4.99	1.33	1.52
2	B	589	GRA	C7-S1	-3.50	1.68	1.76
2	A	589	GRA	C7-S1	-3.44	1.68	1.76
2	B	589	GRA	P1A-O3A	2.94	1.62	1.59
2	B	589	GRA	P2A-O3A	2.86	1.62	1.59
2	A	589	GRA	P1A-O3A	2.69	1.62	1.59
2	A	589	GRA	C8-C7	-2.68	1.48	1.50
2	B	589	GRA	C8-C7	-2.63	1.48	1.50
2	A	589	GRA	P2A-O3A	2.54	1.62	1.59
2	B	589	GRA	C5A-N7A	-2.29	1.34	1.39
2	A	589	GRA	C5A-N7A	-2.15	1.35	1.39

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	589	GRA	C2-N1-C1	7.48	135.98	122.55
2	B	589	GRA	C5A-C4A-N3A	-5.70	118.87	126.72
2	A	589	GRA	C8-C7-S1	5.54	120.00	113.40
2	A	589	GRA	C5A-C4A-N3A	-5.20	119.56	126.72
2	B	589	GRA	C9-C8-C7	5.06	123.44	112.27
2	B	589	GRA	N1A-C2A-N3A	-4.99	121.02	128.58
2	A	589	GRA	N1A-C2A-N3A	-4.99	121.03	128.58
2	A	589	GRA	O4-C7-C8	-4.23	119.10	123.98
2	B	589	GRA	C8-C7-S1	4.13	118.33	113.40
2	B	589	GRA	C6-C5-N2	-4.09	103.87	112.41
2	A	589	GRA	C9-C8-C7	4.06	121.24	112.27
2	B	589	GRA	N3A-C4A-N9A	3.80	133.63	127.17
2	B	589	GRA	C2A-N3A-C4A	3.63	120.71	111.83
2	A	589	GRA	C2A-N3A-C4A	3.48	120.34	111.83
2	B	589	GRA	C5-N2-C4	3.48	129.31	122.82
2	A	589	GRA	N3A-C4A-N9A	3.44	133.02	127.17
2	A	589	GRA	C6-C5-N2	-3.42	105.27	112.41
2	A	589	GRA	C10-C9-C8	3.34	119.18	112.20
2	A	589	GRA	C3-C2-N1	-3.17	105.26	112.00
2	A	589	GRA	C2-N1-C1	3.14	128.18	122.55
2	B	589	GRA	O4-C7-C8	-3.01	120.50	123.98
2	A	589	GRA	N9A-C8A-N7A	-3.01	109.67	113.94
2	A	589	GRA	C6-S1-C7	2.97	110.63	101.84
2	A	589	GRA	C5-N2-C4	2.97	128.34	122.82
2	B	589	GRA	N9A-C8A-N7A	-2.91	109.81	113.94
2	A	589	GRA	C9-C10-C11	-2.90	106.95	114.51
2	B	589	GRA	C10-C9-C8	2.82	118.10	112.20
2	B	589	GRA	C6-S1-C7	2.61	109.57	101.84
2	A	589	GRA	C2X-C1X-N9A	-2.59	106.86	113.30
2	B	589	GRA	C9-C10-C11	-2.54	107.88	114.51
2	B	589	GRA	C5A-N7A-C8A	2.46	107.32	103.45
2	B	589	GRA	C4A-C5A-N7A	-2.44	107.79	110.58
2	A	589	GRA	C5A-N7A-C8A	2.34	107.13	103.45
2	A	589	GRA	C4A-C5A-N7A	-2.34	107.91	110.58
2	B	589	GRA	C16-C12-C13	-2.05	104.84	108.22
2	A	589	GRA	C4A-N9A-C8A	2.03	107.87	105.74

There are no chirality outliers.

All (29) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	589	GRA	C16-C12-C13-O6A
2	A	589	GRA	C13-C12-C14-O1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	589	GRA	C13-C12-C14-O1
2	B	589	GRA	C13-C12-C14-C1
2	B	589	GRA	C16-C12-C14-O1
2	B	589	GRA	C16-C12-C14-C1
2	B	589	GRA	N1-C2-C3-C4
2	B	589	GRA	N2-C5-C6-S1
2	B	589	GRA	C3-C2-N1-C1
2	B	589	GRA	C15-C12-C14-O1
2	A	589	GRA	P2A-O3A-P1A-O2A
2	B	589	GRA	P1A-O3A-P2A-O5A
2	A	589	GRA	C15-C12-C13-O6A
2	B	589	GRA	C15-C12-C13-O6A
2	A	589	GRA	C13-O6A-P2A-O3A
2	B	589	GRA	C2-C3-C4-O3
2	B	589	GRA	P1A-O3A-P2A-O4A
2	B	589	GRA	C2-C3-C4-N2
2	A	589	GRA	C3X-O3X-P3X-O9A
2	B	589	GRA	C3X-O3X-P3X-O9A
2	A	589	GRA	C9-C10-C11-O5
2	A	589	GRA	C9-C10-C11-O6
2	B	589	GRA	C9-C10-C11-O6
2	B	589	GRA	C9-C10-C11-O5
2	B	589	GRA	C11-C10-C9-C8
2	B	589	GRA	C15-C12-C14-C1
2	A	589	GRA	C16-C12-C14-O1
2	A	589	GRA	P2A-O3A-P1A-O1A
2	B	589	GRA	P2A-O3A-P1A-O1A


There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	589	GRA	2	0
2	B	589	GRA	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	558/588 (94%)	-0.12	28 (5%)	34 28	15, 33, 67, 92	1 (0%)
1	B	566/588 (96%)	-0.00	42 (7%)	20 16	15, 33, 73, 96	1 (0%)
All	All	1124/1176 (95%)	-0.06	70 (6%)	26 21	15, 33, 70, 96	2 (0%)

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	238	GLU	7.2
1	B	239	GLN	6.7
1	B	241	ILE	6.0
1	A	241	ILE	5.9
1	B	505	ALA	5.8
1	A	236	ALA	5.8
1	B	238	GLU	5.3
1	A	246	GLU	5.2
1	A	243	ALA	5.1
1	A	240	ILE	5.1
1	B	245	ILE	5.0
1	B	237	ALA	4.9
1	A	237	ALA	4.9
1	A	245	ILE	4.7
1	B	236	ALA	4.4
1	B	243	ALA	4.4
1	A	501	LYS	4.3
1	A	244	GLN	4.2
1	B	515	ILE	4.2
1	B	221	GLY	4.2
1	B	503	VAL	4.1
1	B	516	ILE	4.0
1	A	242	ALA	3.9
1	B	235	GLU	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	250	LEU	3.8
1	B	242	ALA	3.8
1	B	504	LYS	3.8
1	B	50	ILE	3.8
1	A	515	ILE	3.6
1	B	513	GLN	3.6
1	B	240	ILE	3.6
1	B	246	GLU	3.6
1	B	251	LYS	3.5
1	A	516	ILE	3.5
1	A	514	PRO	3.4
1	B	502	LEU	3.3
1	A	239	GLN	3.2
1	A	250	LEU	3.2
1	B	234	ASP	3.1
1	A	499	SER	3.1
1	B	249	LYS	3.1
1	B	233	ASP	3.0
1	A	220	ALA	3.0
1	A	248	SER	3.0
1	A	500	ARG	2.9
1	A	247	ASN	2.8
1	B	374	ASN	2.8
1	B	514	PRO	2.8
1	B	500	ARG	2.8
1	B	248	SER	2.8
1	A	249	LYS	2.7
1	A	59	ARG	2.7
1	A	21	VAL	2.7
1	B	59	ARG	2.7
1	B	498	TYR	2.6
1	A	22	ASN	2.5
1	B	247	ASN	2.5
1	A	251	LYS	2.5
1	B	517	GLY	2.5
1	A	273	GLN	2.4
1	B	501	LYS	2.4
1	B	29	GLN	2.4
1	B	244	GLN	2.4
1	B	586	PHE	2.4
1	B	26	GLU	2.3
1	B	252	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	51	THR	2.2
1	B	220	ALA	2.2
1	B	22	ASN	2.1
1	A	491	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GRA	A	589	56/56	0.86	0.17	57,80,93,94	0
2	GRA	B	589	56/56	0.86	0.17	65,85,91,92	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around GRA A 589:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GRA B 589:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.