



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 09:07 PM UTC

PDB ID : 3GME / pdb_00003gme
Title : Crystal Structure of Polynucleotide Phosphorylase in complex with RNase E and manganese
Authors : Nurmohamed, S.; Luisi, B.L.
Deposited on : 2009-03-13
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	549	
2	D	41	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MN	A	551	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3854 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polyribonucleotide nucleotidyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	482	Total	C	N	O	S	0	0	0
			3560	2239	617	684	20			

- Molecule 2 is a protein called Ribonuclease E.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	21	Total	C	N	O	0	0	0
			134	80	30	24			

- Molecule 3 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Mn	0	0
			2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	150	Total	O	0	0
			150	150		
4	D	8	Total	O	0	0
			8	8		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Polyrribonucleotide nucleotidyltransferase



- Molecule 2: Ribonuclease E



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	158.57Å 158.57Å 156.12Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	25.91 – 2.40 25.91 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.3 (25.91-2.40) 99.3 (25.91-2.40)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.49 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.286 , 0.298 0.281 , 0.295	Depositor DCC
R_{free} test set	1490 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	47.0	Xtriage
Anisotropy	0.106	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 41.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3854	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	2/3615 (0.1%)	0.81	1/4909 (0.0%)
2	D	0.46	0/139	0.82	0/191
All	All	0.70	2/3754 (0.1%)	0.81	1/5100 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	65	GLU	C-O	-5.33	1.17	1.24
1	A	64	GLN	C-O	-5.13	1.17	1.23

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	191	SER	N-CA-C	5.83	117.65	108.96

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3560	0	3508	61	0
2	D	134	0	119	2	0
3	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	150	0	0	2	0
4	D	8	0	0	1	0
All	All	3854	0	3627	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:342:LEU:HD11	1:A:349:GLN:HE21	1.05	1.10
1:A:406:LEU:HD11	1:A:506:GLN:OE1	1.67	0.93
1:A:342:LEU:HD11	1:A:349:GLN:NE2	1.84	0.92
1:A:202:LEU:HD12	1:A:202:LEU:O	1.72	0.89
1:A:342:LEU:CD1	1:A:349:GLN:HE21	1.89	0.83
1:A:493:PHE:HZ	1:A:523:LEU:CD2	1.97	0.78
1:A:375:THR:HG22	4:A:628:HOH:O	1.82	0.78
1:A:406:LEU:CD1	1:A:506:GLN:OE1	2.32	0.77
1:A:396:PRO:HA	1:A:400:GLU:OE1	1.84	0.77
1:A:519:MET:O	1:A:523:LEU:HD23	1.84	0.76
1:A:544:PRO:O	1:A:545:ARG:C	2.30	0.74
1:A:470:LYS:HE2	1:A:472:GLY:O	1.90	0.70
1:A:308:ARG:HG2	1:A:313:GLU:OE1	1.90	0.70
1:A:493:PHE:CZ	1:A:523:LEU:CD2	2.76	0.69
1:A:493:PHE:CZ	1:A:523:LEU:HD22	2.28	0.68
1:A:188:MET:O	1:A:189:VAL:HG23	1.93	0.68
1:A:493:PHE:HZ	1:A:523:LEU:HD21	1.57	0.68
1:A:16:VAL:HG22	1:A:35:MET:HG3	1.79	0.63
1:A:456:PRO:HA	1:A:544:PRO:HG2	1.82	0.60
1:A:375:THR:CG2	4:A:628:HOH:O	2.45	0.59
1:A:202:LEU:HD12	1:A:202:LEU:C	2.17	0.59
1:A:466:MET:HB2	1:A:493:PHE:HB3	1.84	0.58
1:A:397:LYS:O	1:A:398:ARG:C	2.46	0.57
1:A:181:GLY:HA2	1:A:187:LEU:HD12	1.87	0.56
1:A:144:PRO:HD3	1:A:231:TRP:HB2	1.88	0.55
1:A:325:ARG:HD2	1:A:346:GLY:HA3	1.87	0.55
1:A:477:VAL:HB	1:A:525:GLN:OE1	2.06	0.55
1:A:188:MET:O	1:A:189:VAL:CG2	2.55	0.54
2:D:1056:ARG:HD2	4:D:118:HOH:O	2.09	0.53
1:A:33:VAL:HG21	1:A:133:GLY:CA	2.40	0.51
1:A:493:PHE:CZ	1:A:523:LEU:HD21	2.43	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:478:LEU:HD12	1:A:478:LEU:N	2.26	0.51
1:A:188:MET:CG	1:A:189:VAL:N	2.73	0.51
1:A:188:MET:HG2	1:A:189:VAL:N	2.25	0.51
1:A:534:LEU:O	1:A:538:GLU:HG3	2.11	0.51
1:A:397:LYS:O	1:A:399:ARG:N	2.45	0.50
1:A:64:GLN:HE21	1:A:116:THR:HB	1.77	0.49
1:A:331:THR:HB	2:D:1048:HIS:HB2	1.94	0.49
1:A:322:ASP:HA	1:A:525:GLN:HE21	1.78	0.48
1:A:342:LEU:HD11	1:A:349:GLN:CG	2.43	0.48
1:A:23:MET:HE1	1:A:32:MET:HE2	1.95	0.47
1:A:33:VAL:HG21	1:A:133:GLY:HA2	1.96	0.47
1:A:2:LEU:HD12	1:A:2:LEU:N	2.32	0.45
1:A:478:LEU:N	1:A:478:LEU:CD1	2.80	0.45
1:A:20:THR:OG1	1:A:21:GLY:N	2.50	0.43
1:A:351:LEU:HB3	1:A:430:GLU:HB2	2.00	0.43
1:A:519:MET:O	1:A:523:LEU:CD2	2.63	0.43
1:A:28:THR:HG21	1:A:143:ILE:HD11	2.01	0.43
1:A:44:VAL:HG22	1:A:113:VAL:HG22	2.01	0.43
1:A:491:MET:HE3	1:A:493:PHE:CD1	2.54	0.42
1:A:2:LEU:HB3	1:A:22:MET:HG3	2.01	0.42
1:A:91:ILE:HA	1:A:94:LEU:HD12	2.00	0.42
1:A:189:VAL:HB	1:A:507:MET:HB2	2.01	0.42
1:A:316:ILE:H	1:A:316:ILE:HG13	1.72	0.42
1:A:502:ILE:HG12	1:A:530:ARG:HH22	1.85	0.42
1:A:543:ALA:HA	1:A:544:PRO:HD3	1.88	0.42
1:A:322:ASP:HB3	1:A:525:GLN:NE2	2.35	0.42
1:A:406:LEU:CG	1:A:506:GLN:OE1	2.67	0.42
1:A:4:PRO:HB3	1:A:19:GLU:OE2	2.20	0.41
1:A:509:ILE:HG21	1:A:514:ILE:HG21	2.02	0.41
1:A:3:ASN:HA	1:A:4:PRO:HD3	1.93	0.41
1:A:67:THR:HG21	1:A:74:PRO:HG3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	472/549 (86%)	447 (95%)	24 (5%)	1 (0%)	43	58
2	D	19/41 (46%)	18 (95%)	1 (5%)	0	100	100
All	All	491/590 (83%)	465 (95%)	25 (5%)	1 (0%)	43	58

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	52	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	369/446 (83%)	364 (99%)	5 (1%)	59	79
2	D	10/28 (36%)	10 (100%)	0	100	100
All	All	379/474 (80%)	374 (99%)	5 (1%)	61	80

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	202	LEU
1	A	342	LEU
1	A	406	LEU
1	A	464	ILE
1	A	518	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	64	GLN
1	A	112	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	167	GLN
1	A	213	GLN
1	A	220	ASN
1	A	349	GLN
1	A	487	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	482/549 (87%)	1.34	117 (24%) 2 1	24, 45, 90, 112	1 (0%)
2	D	21/41 (51%)	1.88	5 (23%) 2 1	36, 52, 71, 80	0
All	All	503/590 (85%)	1.37	122 (24%) 2 1	24, 46, 90, 112	1 (0%)

All (122) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	159	ASP	8.7
1	A	267	TYR	6.3
1	A	261	THR	5.5
1	A	241	ALA	5.5
1	A	262	ASP	5.4
2	D	1039	GLY	5.1
1	A	396	PRO	5.1
1	A	242	LEU	4.9
1	A	297	LEU	4.8
1	A	238	VAL	4.6
1	A	21	GLY	4.5
1	A	514	ILE	4.4
1	A	543	ALA	4.3
1	A	244	ALA	4.2
1	A	265	GLU	4.1
1	A	263	LYS	4.0
1	A	394	GLY	4.0
1	A	235	PRO	4.0
1	A	55	ASP	3.9
1	A	493	PHE	3.8
1	A	314	PRO	3.8
1	A	266	ARG	3.7
2	D	1059	PRO	3.7
1	A	246	VAL	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	237	PRO	3.6
1	A	245	ARG	3.6
1	A	398	ARG	3.6
1	A	12	GLY	3.6
1	A	264	GLN	3.6
2	D	1040	ALA	3.5
1	A	490	ASP	3.4
1	A	397	LYS	3.4
1	A	301	GLU	3.4
1	A	488	LEU	3.4
1	A	489	GLY	3.3
1	A	44	VAL	3.3
1	A	240	GLU	3.2
1	A	232	ASP	3.2
1	A	230	ARG	3.1
1	A	142	GLY	3.1
1	A	333	VAL	3.1
1	A	331	THR	3.1
1	A	393	VAL	3.1
1	A	224	LYS	3.0
1	A	420	LYS	3.0
1	A	464	ILE	3.0
1	A	52	PRO	3.0
1	A	321	LYS	3.0
2	D	1058	GLN	3.0
1	A	188	MET	2.9
2	D	1045	THR	2.9
1	A	51	LYS	2.9
1	A	544	PRO	2.9
1	A	53	GLY	2.8
1	A	316	ILE	2.8
1	A	309	VAL	2.8
1	A	140	LEU	2.8
1	A	542	ASN	2.8
1	A	515	THR	2.8
1	A	491	MET	2.8
1	A	300	ILE	2.8
1	A	46	GLY	2.7
1	A	395	SER	2.7
1	A	392	MET	2.6
1	A	312	GLY	2.6
1	A	108	VAL	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	507	MET	2.6
1	A	476	VAL	2.6
1	A	234	GLN	2.6
1	A	511	ILE	2.5
1	A	541	ILE	2.5
1	A	107	PHE	2.5
1	A	223	VAL	2.5
1	A	499	ARG	2.5
1	A	513	GLY	2.5
1	A	337	THR	2.5
1	A	518	ILE	2.5
1	A	226	ALA	2.5
1	A	492	ASP	2.4
1	A	70	ALA	2.4
1	A	299	ALA	2.4
1	A	545	ARG	2.4
1	A	323	MET	2.4
1	A	474	ASN	2.4
1	A	438	SER	2.4
1	A	313	GLU	2.4
1	A	2	LEU	2.4
1	A	54	GLN	2.4
1	A	45	VAL	2.4
1	A	387	VAL	2.3
1	A	110	GLU	2.3
1	A	487	HIS	2.3
1	A	1	MET	2.3
1	A	77	PHE	2.3
1	A	311	ALA	2.3
1	A	540	ALA	2.3
1	A	106	GLY	2.3
1	A	217	GLN	2.3
1	A	233	TRP	2.3
1	A	303	ASN	2.2
1	A	345	ARG	2.3
1	A	322	ASP	2.2
1	A	418	MET	2.2
1	A	172	GLU	2.2
1	A	471	GLU	2.2
1	A	243	ASN	2.2
1	A	475	TYR	2.2
1	A	318	GLY	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	538	GLU	2.2
1	A	56	PHE	2.1
1	A	412	LEU	2.1
1	A	437	SER	2.1
1	A	187	LEU	2.1
1	A	468	LEU	2.1
1	A	298	HIS	2.1
1	A	358	THR	2.1
1	A	390	THR	2.1
1	A	419	ASP	2.1
1	A	424	THR	2.0
1	A	194	GLU	2.0
1	A	221	GLU	2.0
1	A	406	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MN	A	550	1/1	-0.09	0.26	228,228,228,228	0
3	MN	A	551	1/1	0.62	0.64	82,82,82,82	0

6.5 Other polymers [i](#)

There are no such residues in this entry.