



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:09 PM UTC

PDB ID : 3GMP / pdb_00003gmp
Title : Structure of mouse CD1d in complex with PBS-25
Authors : Schiefner, A.; Wilson, I.A.
Deposited on : 2009-03-14
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	287	4% 77% 14% • 7%
2	B	99	2% 86% 12% •
3	C	5	80% 20%
4	D	6	100%

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 3591 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called T-cell surface glycoprotein CD1d1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	268	Total	C	N	O	S	0	7	0
			2219	1410	385	411	13			


There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	201	HIS	ASP	SEE REMARK 999	UNP P11609
A	280	GLY	-	expression tag	UNP P11609
A	281	SER	-	expression tag	UNP P11609
A	282	HIS	-	expression tag	UNP P11609
A	283	HIS	-	expression tag	UNP P11609
A	284	HIS	-	expression tag	UNP P11609
A	285	HIS	-	expression tag	UNP P11609
A	286	HIS	-	expression tag	UNP P11609
A	287	HIS	-	expression tag	UNP P11609

- Molecule 2 is a protein called Beta-2 microglobulin.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C	N	O	S	0	2	0
			833	531	140	155	7			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	6	Total	C	N	O	0	0	0
			71	40	2	29			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is (2S,3S,4R)-N-OCTANOYL-1-[(ALPHA-D-GALACTOPYRANOSYL)OXY]-2-AMINO-OCTADECANE-3,4-DIOL (CCD ID: PBS) (formula: C₃₂H₆₃NO₉).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			42	32	1	9		

- Molecule 7 is PALMITIC ACID (CCD ID: PLM) (formula: $C_{16}H_{32}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			18	16	2		

- Molecule 8 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			4	2	2		
8	A	1	Total	C	O	0	0
			4	2	2		
8	A	1	Total	C	O	0	1
			8	4	4		
8	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	218	Total	O	0	5
			223	223		
9	B	89	Total	O	0	1
			90	90		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: T-cell surface glycoprotein CD1d1




- Molecule 2: Beta-2 microglobulin



- Molecule 3: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	41.71Å 97.73Å 55.49Å 90.00° 106.52° 90.00°	Depositor
Resolution (Å)	27.78 – 1.70 27.78 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.6 (27.78-1.70) 98.9 (27.78-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.03	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.94 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.5.0066	Depositor
R, R_{free}	0.176 , 0.216 (Not available) , 0.234	Depositor DCC
R_{free} test set	2361 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	25.1	Xtriage
Anisotropy	0.020	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 38.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3591	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, BMA, MAN, PLM, FUC, NAG, PBS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.21	3/2282 (0.1%)	1.13	6/3098 (0.2%)
2	B	1.33	5/859 (0.6%)	1.24	2/1164 (0.2%)
All	All	1.24	8/3141 (0.3%)	1.16	8/4262 (0.2%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	54	MET	CG-SD	-8.31	1.59	1.80
1	A	13	LEU	C-O	6.24	1.31	1.24
2	B	54	MET	CB-CG	6.05	1.70	1.52
2	B	79	ALA	N-CA	-5.99	1.38	1.45
2	B	48	LYS	N-CA	5.61	1.52	1.46
1	A	135	THR	CA-CB	5.18	1.61	1.53
2	B	81	ARG	N-CA	5.12	1.52	1.46
1	A	38	HIS	C-O	5.08	1.30	1.23

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	204	ARG	N-CA-C	8.84	120.35	107.88
2	B	47	PRO	CA-C-N	-7.89	109.55	122.65
2	B	47	PRO	C-N-CA	-7.89	109.55	122.65
1	A	94	TYR	N-CA-C	-6.61	100.48	110.39
1	A	216	LYS	CA-C-N	-5.38	114.22	119.76
1	A	216	LYS	C-N-CA	-5.38	114.22	119.76
1	A	109	GLY	N-CA-C	-5.32	102.12	111.80
1	A	166	ASP	N-CA-C	5.19	119.08	112.12

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2219	0	2111	45	3
2	B	833	0	806	11	1
3	C	61	0	52	0	1
4	D	71	0	61	0	0
5	A	14	0	13	3	0
6	A	42	0	63	1	0
7	A	18	0	31	0	0
8	A	16	0	24	2	0
8	B	4	0	6	0	0
9	A	223	0	0	4	2
9	B	90	0	0	4	1
All	All	3591	0	3167	55	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:203:HIS:N	1:A:255:ALA:HB2	1.60	1.16
1:A:257:GLU:HG2	1:A:257:GLU:O	1.41	1.11
1:A:59:SER:OG	1:A:62:GLN:HG3	1.48	1.10
1:A:107:TYR:HB3	1:A:108:PRO:HD2	1.18	1.09
1:A:234:ARG:HG2	1:A:234:ARG:HH11	1.10	1.08
1:A:60:ASN:O	1:A:64:GLU:HG2	1.66	0.95
1:A:107:TYR:CB	1:A:108:PRO:HD2	2.00	0.89
1:A:22:SER:CB	5:A:288:NAG:H82	2.02	0.89
1:A:257:GLU:O	1:A:257:GLU:CG	2.23	0.85
1:A:61:GLN:HE22	1:A:65:LYS:HE3	1.44	0.82
1:A:234:ARG:HG2	1:A:234:ARG:NH1	1.90	0.79
1:A:248:GLN:OE1	2:B:99:MET:HG2	1.84	0.78
1:A:22:SER:HB2	5:A:288:NAG:H82	1.68	0.76
1:A:118:VAL:HG11	6:A:300:PBS:H132	1.69	0.75
1:A:203:HIS:N	1:A:255:ALA:CB	2.47	0.74
1:A:248:GLN:HG3	8:A:303:EDO:H21	1.70	0.74

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:107:TYR:HB3	1:A:108:PRO:CD	2.09	0.74
1:A:236:ASP:OD2	9:A:519:HOH:O	2.06	0.73
1:A:234:ARG:HH11	1:A:234:ARG:CG	1.98	0.71
1:A:61:GLN:HE22	1:A:65:LYS:CE	2.05	0.68
1:A:22:SER:HB3	5:A:288:NAG:H82	1.76	0.66
1:A:61:GLN:NE2	1:A:65:LYS:HE3	2.11	0.66
2:B:1:ILE:HG13	2:B:2:GLN:N	2.13	0.64
2:B:2:GLN:HG2	9:B:268:HOH:O	2.02	0.59
2:B:2:GLN:NE2	9:B:268:HOH:O	1.97	0.59
2:B:7:ILE:HD12	2:B:91:LYS:HD3	1.84	0.59
1:A:107:TYR:CB	1:A:108:PRO:CD	2.75	0.56
1:A:234:ARG:NH1	1:A:234:ARG:CG	2.65	0.55
1:A:195:SER:OG	1:A:204:ARG:NH1	2.41	0.53
1:A:100[A]:LEU:HD21	1:A:102:ALA:HB2	1.90	0.53
1:A:100[A]:LEU:C	1:A:100[A]:LEU:HD23	2.35	0.52
1:A:234:ARG:HD2	9:A:451:HOH:O	2.09	0.52
1:A:248:GLN:HG3	8:A:303:EDO:C2	2.39	0.51
2:B:7:ILE:CD1	2:B:91:LYS:HD3	2.40	0.51
1:A:173:ARG:HG2	1:A:173:ARG:HH11	1.76	0.51
1:A:61:GLN:NE2	1:A:65:LYS:CE	2.73	0.50
1:A:236:ASP:CG	9:A:519:HOH:O	2.55	0.49
1:A:134[B]:GLN:HG2	1:A:135:THR:O	2.14	0.48
2:B:74:GLU:HA	2:B:74:GLU:OE1	2.14	0.48
1:A:127:ARG:HE	1:A:134[B]:GLN:CD	2.22	0.48
1:A:59:SER:HG	1:A:62:GLN:HG3	1.71	0.47
1:A:168:CYS:HB3	1:A:169:PRO:CD	2.45	0.47
1:A:20:ASN:HA	1:A:94:TYR:CD1	2.51	0.46
2:B:47:PRO:O	9:B:261:HOH:O	2.21	0.45
1:A:230:GLN:HA	1:A:230:GLN:OE1	2.17	0.45
1:A:168:CYS:O	1:A:172:VAL:HG23	2.19	0.43
1:A:69:MET:HE2	1:A:69:MET:HB2	1.69	0.43
2:B:32:PRO:HB2	2:B:33:PRO:HD2	2.00	0.43
2:B:2:GLN:CG	9:B:268:HOH:O	2.64	0.42
1:A:234:ARG:NH1	1:A:235:GLY:O	2.53	0.41
1:A:145:LEU:HB3	1:A:146:PRO:HD3	2.02	0.41
1:A:43:ASP:OD2	9:A:445:HOH:O	2.21	0.40
1:A:277:LEU:HD23	1:A:277:LEU:HA	1.80	0.40
2:B:10:TYR:HB3	2:B:99:MET:HE2	2.02	0.40
1:A:216:LYS:N	1:A:217:PRO:CD	2.85	0.40

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:1:ILE:CD1	9:B:392:HOH:O[1_655]	1.51	0.69
1:A:79:ARG:NH1	1:A:230:GLN:NE2[2_555]	2.07	0.13
1:A:190:VAL:CG2	3:C:4:MAN:O3[2_445]	2.10	0.10
9:A:433:HOH:O	9:A:448:HOH:O[1_655]	2.10	0.10
1:A:227[A]:GLN:OE1	9:A:337:HOH:O[2_545]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	271/287 (94%)	266 (98%)	5 (2%)	0	100	100
2	B	99/99 (100%)	97 (98%)	2 (2%)	0	100	100
All	All	370/386 (96%)	363 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	241/250 (96%)	235 (98%)	6 (2%)	42	24
2	B	95/93 (102%)	94 (99%)	1 (1%)	65	54
All	All	336/343 (98%)	329 (98%)	7 (2%)	47	30

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	MET
1	A	61	GLN
1	A	113	GLU
1	A	234	ARG
1	A	257	GLU
1	A	281	SER
2	B	1	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	61	GLN
1	A	186	GLN
1	A	203	HIS
2	B	17	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

11 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	C	1	3,1	14,14,15	0.68	0	17,19,21	2.02	6 (35%)
3	NAG	C	2	3	14,14,15	0.78	0	17,19,21	1.34	1 (5%)
3	BMA	C	3	3	11,11,12	0.50	0	15,15,17	1.81	4 (26%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MAN	C	4	3	11,11,12	0.56	0	15,15,17	1.33	2 (13%)
3	MAN	C	5	3	11,11,12	0.55	0	15,15,17	1.73	3 (20%)
4	NAG	D	1	1,4	14,14,15	0.98	0	17,19,21	1.28	2 (11%)
4	NAG	D	2	4	14,14,15	0.61	0	17,19,21	1.68	3 (17%)
4	BMA	D	3	4	11,11,12	0.89	0	15,15,17	1.78	5 (33%)
4	MAN	D	4	4	11,11,12	0.62	0	15,15,17	1.68	3 (20%)
4	MAN	D	5	4	11,11,12	0.69	0	15,15,17	1.45	2 (13%)
4	FUC	D	6	4	10,10,11	0.70	0	14,14,16	1.48	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	2/6/23/26	0/1/1/1
3	BMA	C	3	3	-	2/2/19/22	0/1/1/1
3	MAN	C	4	3	-	0/2/19/22	0/1/1/1
3	MAN	C	5	3	-	1/2/19/22	0/1/1/1
4	NAG	D	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	D	2	4	-	0/6/23/26	0/1/1/1
4	BMA	D	3	4	-	0/2/19/22	0/1/1/1
4	MAN	D	4	4	-	0/2/19/22	0/1/1/1
4	MAN	D	5	4	-	0/2/19/22	0/1/1/1
4	FUC	D	6	4	-	-	0/1/1/1

There are no bond length outliers.

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	3	BMA	C3-C4-C5	-4.90	101.34	110.23
4	D	2	NAG	C2-N2-C7	-4.50	116.87	122.90
3	C	1	NAG	C1-C2-N2	4.16	116.99	110.43
3	C	5	MAN	C2-C3-C4	-3.81	104.15	110.86
3	C	5	MAN	O5-C5-C6	3.74	114.95	107.66
4	D	3	BMA	C1-O5-C5	3.67	117.10	112.19
4	D	4	MAN	O2-C2-C1	-3.65	100.87	109.22
3	C	1	NAG	C4-C3-C2	3.63	116.33	111.02

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	5	MAN	O3-C3-C4	-3.25	102.72	110.38
4	D	2	NAG	O5-C1-C2	-3.18	106.37	111.29
3	C	2	NAG	C3-C4-C5	-3.10	104.61	110.23
4	D	6	FUC	O5-C1-C2	-3.09	103.41	110.79
3	C	3	BMA	C1-C2-C3	3.07	114.12	109.64
3	C	1	NAG	C3-C4-C5	-3.05	104.69	110.23
4	D	1	NAG	C1-C2-N2	2.99	115.15	110.43
3	C	4	MAN	C6-C5-C4	-2.97	105.72	113.02
4	D	4	MAN	C3-C4-C5	-2.92	104.94	110.23
4	D	3	BMA	O2-C2-C3	-2.72	104.51	110.15
3	C	1	NAG	O5-C5-C6	2.69	112.89	107.66
3	C	1	NAG	O7-C7-C8	-2.61	117.41	122.05
3	C	1	NAG	C8-C7-N2	2.54	120.33	116.12
4	D	3	BMA	O6-C6-C5	-2.49	102.85	111.33
4	D	4	MAN	C1-C2-C3	-2.43	106.10	109.64
4	D	2	NAG	C8-C7-N2	2.43	120.15	116.12
4	D	5	MAN	C1-O5-C5	2.35	115.34	112.19
4	D	3	BMA	O3-C3-C4	-2.24	105.10	110.38
3	C	5	MAN	C3-C4-C5	-2.23	106.18	110.23
4	D	3	BMA	O2-C2-C1	-2.20	104.18	109.22
3	C	3	BMA	O5-C5-C4	-2.20	105.48	110.83
3	C	3	BMA	O5-C5-C6	2.10	111.75	107.66
4	D	1	NAG	O4-C4-C5	-2.03	104.32	109.32
3	C	4	MAN	O5-C5-C6	2.01	111.58	107.66

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	3	BMA	O5-C5-C6-O6
3	C	3	BMA	C4-C5-C6-O6
3	C	2	NAG	C8-C7-N2-C2
3	C	2	NAG	O7-C7-N2-C2
3	C	5	MAN	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	4	MAN	0	1

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	EDO	A	302	-	3,3,3	0.45	0	2,2,2	0.66	0
7	PLM	A	301	-	17,17,17	0.80	0	17,17,17	1.18	2 (11%)
8	EDO	A	304[A]	-	3,3,3	0.37	0	2,2,2	0.77	0
5	NAG	A	288	1	14,14,15	0.93	1 (7%)	17,19,21	1.52	3 (17%)
8	EDO	A	303	-	3,3,3	0.54	0	2,2,2	0.63	0
8	EDO	A	304[B]	-	3,3,3	0.45	0	2,2,2	0.05	0
8	EDO	B	100	-	3,3,3	0.82	0	2,2,2	0.45	0
6	PBS	A	300	-	42,42,42	0.90	3 (7%)	48,51,51	1.56	10 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	EDO	A	302	-	-	1/1/1/1	-
7	PLM	A	301	-	-	8/15/15/15	-
8	EDO	A	304[A]	-	-	1/1/1/1	-
5	NAG	A	288	1	-	0/6/23/26	0/1/1/1
8	EDO	A	303	-	-	0/1/1/1	-
8	EDO	A	304[B]	-	-	1/1/1/1	-
8	EDO	B	100	-	-	0/1/1/1	-
6	PBS	A	300	-	-	9/40/60/60	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	300	PBS	C-C1	3.13	1.57	1.51
6	A	300	PBS	OC1-CG1	2.87	1.45	1.40
6	A	300	PBS	O2-C18	2.09	1.27	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	288	NAG	C1-C2	2.06	1.55	1.52

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	300	PBS	C-OC1-CG1	3.50	121.31	113.80
6	A	300	PBS	OG-CG-CG4	3.44	115.91	109.70
5	A	288	NAG	O5-C5-C4	-3.41	102.53	110.83
7	A	301	PLM	O1-C1-O2	-3.37	114.68	123.33
5	A	288	NAG	C4-C3-C2	3.05	115.49	111.02
6	A	300	PBS	C1-N-C18	-2.98	118.43	123.40
6	A	300	PBS	C7-C8-C9	-2.86	99.90	114.37
7	A	301	PLM	O1-C1-C2	2.68	122.47	114.00
6	A	300	PBS	C21-C20-C19	-2.66	103.36	113.13
6	A	300	PBS	O2-C18-C19	2.64	126.80	122.02
6	A	300	PBS	C8-C9-C10	-2.36	102.43	114.37
6	A	300	PBS	C19-C18-N	-2.33	111.76	115.86
6	A	300	PBS	CG1-CG2-CG3	2.21	114.66	110.01
6	A	300	PBS	CG3-CG4-CG	-2.11	106.40	110.23
5	A	288	NAG	C1-O5-C5	-2.01	109.49	112.19

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	301	PLM	CB-CC-CD-CE
7	A	301	PLM	C8-C9-CA-CB
7	A	301	PLM	C5-C6-C7-C8
6	A	300	PBS	C10-C11-C12-C13
7	A	301	PLM	C9-CA-CB-CC
6	A	300	PBS	C11-C12-C13-C14
7	A	301	PLM	CC-CD-CE-CF
6	A	300	PBS	C13-C14-C15-C16
6	A	300	PBS	C12-C13-C14-C15
6	A	300	PBS	C11-C10-C9-C8
6	A	300	PBS	C14-C15-C16-C17
6	A	300	PBS	C9-C10-C11-C12
8	A	302	EDO	O1-C1-C2-O2
7	A	301	PLM	C2-C3-C4-C5
6	A	300	PBS	C7-C8-C9-C10
7	A	301	PLM	O2-C1-C2-C3
8	A	304[A]	EDO	O1-C1-C2-O2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
8	A	304[B]	EDO	O1-C1-C2-O2
7	A	301	PLM	O1-C1-C2-C3
6	A	300	PBS	OG-CG-CG5-OG6


There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	288	NAG	3	0
8	A	303	EDO	2	0
6	A	300	PBS	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	268/287 (93%)	0.25	11 (4%) 41 45	8, 20, 39, 53	7 (2%)
2	B	99/99 (100%)	-0.06	2 (2%) 65 69	8, 19, 29, 36	2 (2%)
All	All	367/386 (95%)	0.17	13 (3%) 47 52	8, 19, 36, 53	9 (2%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	280	GLY	3.5
1	A	255	ALA	3.2
1	A	94	TYR	3.0
1	A	108	PRO	2.9
2	B	1	ILE	2.5
1	A	281	SER	2.4
2	B	48	LYS	2.4
1	A	110	ASN	2.3
1	A	109	GLY	2.3
1	A	256	GLY	2.2
1	A	93	ASP	2.1
1	A	271	GLY	2.1
1	A	107	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BMA	C	3	11/12	0.57	0.15	52,57,61,63	0
3	MAN	C	5	11/12	0.61	0.16	60,63,65,67	0
4	MAN	D	4	11/12	0.65	0.14	52,55,56,58	0
3	NAG	C	2	14/15	0.71	0.14	49,52,54,57	0
4	FUC	D	6	10/11	0.84	0.12	33,36,39,41	0
3	MAN	C	4	11/12	0.85	0.11	33,40,47,49	0
4	BMA	D	3	11/12	0.88	0.10	32,34,41,46	0
4	NAG	D	2	14/15	0.90	0.09	30,33,36,38	0
4	MAN	D	5	11/12	0.91	0.09	29,32,38,43	0
3	NAG	C	1	14/15	0.92	0.10	28,36,41,42	0
4	NAG	D	1	14/15	0.94	0.07	24,26,28,29	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	NAG	A	288	14/15	0.75	0.12	52,59,67,69	0
7	PLM	A	301	18/18	0.86	0.12	36,43,47,48	0
8	EDO	A	304[A]	4/4	0.87	0.12	27,29,29,31	4
8	EDO	A	304[B]	4/4	0.87	0.12	25,26,26,27	4
6	PBS	A	300	42/42	0.92	0.09	23,30,39,45	0
8	EDO	A	303	4/4	0.92	0.10	25,31,34,41	0
8	EDO	A	302	4/4	0.94	0.07	21,22,25,30	0
8	EDO	B	100	4/4	0.95	0.09	20,22,26,30	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around PBS A 300:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.