



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 01:50 AM UTC

PDB ID : 4GMP / pdb_00004gmp
Title : Crystal structure of enterovirus 71 strain 1095 procapsid
Authors : Yoder, J.D.; Hafenstein, S.
Deposited on : 2012-08-16
Resolution : 3.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1270 (4.10-3.70)
Clashscore	190562	1034 (4.08-3.72)
Ramachandran outliers	187476	1251 (4.10-3.70)
Sidechain outliers	187428	1243 (4.10-3.70)
RSRZ outliers	180081	1269 (4.10-3.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	0	323	44% 27% 39% 7% 27%
2	1	297	22% 30% 36% 7% 27%
3	3	242	41% 41% 41% 16%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5395 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called capsid protein VP0.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	0	237	Total	C	N	O	S	0	0	0
			1833	1179	301	345	8			

- Molecule 2 is a protein called capsid protein VP1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	1	218	Total	C	N	O	S	0	0	0
			1717	1101	289	316	11			

- Molecule 3 is a protein called capsid protein VP3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	3	240	Total	C	N	O	S	0	0	0
			1845	1188	304	342	11			



● Molecule 3: capsid protein VP3



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 ₂ 3 ₂	Depositor
Cell constants a, b, c, α , β , γ	350.25Å 350.25Å 350.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.53 – 3.90 49.53 – 3.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (49.53-3.90) 99.8 (49.53-3.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.81 (at 3.88Å)	Xtriage
Refinement program	CNS 1.3	Depositor
R, R_{free}	0.269 , 0.285 0.262 , 0.280	Depositor DCC
R_{free} test set	3398 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	109.8	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 102.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.37	EDS
Total number of atoms	5395	wwPDB-VP
Average B, all atoms (Å ²)	125.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	0	0.56	0/1888	1.05	11/2591 (0.4%)
2	1	0.61	0/1769	1.03	7/2411 (0.3%)
3	3	0.58	0/1897	1.09	18/2596 (0.7%)
All	All	0.59	0/5554	1.06	36/7598 (0.5%)

There are no bond length outliers.

The worst 5 of 36 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	0	161	PHE	N-CA-C	-8.57	102.26	112.89
3	3	45	GLU	N-CA-C	-8.28	102.21	111.07
3	3	18	ASP	N-CA-C	7.61	120.62	110.88
1	0	294	ASP	N-CA-C	-7.26	99.71	110.23
3	3	74	SER	N-CA-C	7.11	117.98	108.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	0	1833	0	1774	134	3
2	1	1717	0	1674	135	8
3	3	1845	0	1822	141	1
All	All	5395	0	5270	366	9

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 34.

The worst 5 of 366 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:3:20:VAL:HG22	3:3:21:SER:H	1.17	1.06
3:3:206:ILE:HD12	3:3:206:ILE:H	1.23	1.01
1:0:233:TYR:HA	3:3:66:MET:HE3	1.43	0.99
2:1:229:MET:HE2	2:1:231:GLY:H	1.28	0.98
3:3:109:THR:HB	3:3:228:LEU:HB3	1.52	0.91

The worst 5 of 9 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:0:208:THR:CG2	2:1:283:SER:OG[13_454]	1.16	1.04
1:0:208:THR:CG2	2:1:283:SER:CB[13_454]	1.57	0.63
2:1:282:ASN:ND2	2:1:283:SER:N[13_454]	1.68	0.52
3:3:63:THR:CG2	3:3:63:THR:CG2[13_454]	1.71	0.49
2:1:282:ASN:O	2:1:282:ASN:CB[13_454]	1.75	0.45

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	0	235/323 (73%)	175 (74%)	42 (18%)	18 (8%)	1	12
2	1	214/297 (72%)	171 (80%)	27 (13%)	16 (8%)	1	12
3	3	238/242 (98%)	187 (79%)	34 (14%)	17 (7%)	1	13
All	All	687/862 (80%)	533 (78%)	103 (15%)	51 (7%)	1	12

5 of 51 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	0	97	ALA
1	0	112	SER
1	0	127	VAL
2	1	169	LEU
2	1	196	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	0	201/272 (74%)	188 (94%)	13 (6%)	15	42
2	1	186/250 (74%)	167 (90%)	19 (10%)	7	26
3	3	200/202 (99%)	177 (88%)	23 (12%)	5	22
All	All	587/724 (81%)	532 (91%)	55 (9%)	8	29

5 of 55 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	1	228	ASN
3	3	51	THR
3	3	230	LYS
3	3	204	VAL
2	1	237	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
2	1	172	GLN
2	1	176	ASN
3	3	176	HIS
2	1	278	ASN
3	3	110	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	0	237/323 (73%)	3.01	142 (59%) 0 1	89, 133, 217, 234	0
2	1	218/297 (73%)	1.75	64 (29%) 1 3	84, 110, 155, 200	0
3	3	240/242 (99%)	2.06	99 (41%) 0 1	80, 112, 189, 230	0
All	All	695/862 (80%)	2.29	305 (43%) 0 1	80, 114, 211, 234	0

The worst 5 of 305 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	0	97	ALA	16.0
1	0	96	GLU	13.4
3	3	179	ALA	10.9
1	0	126	ASP	9.2
1	0	130	ASN	9.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.