



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:49 PM UTC

PDB ID : 5GML / pdb_00005gml
Title : Crystal Structure of GRASP Domain of GRASP55 with N terminal extra residues
Authors : Shi, N.; Shi, X.; Morelli, X.; Betzi, S.; Huang, X.
Deposited on : 2016-07-14
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1091 (2.54-2.54)
Clashscore	190562	1120 (2.54-2.54)
Ramachandran outliers	187476	1106 (2.54-2.54)
Sidechain outliers	187428	1106 (2.54-2.54)
RSRZ outliers	180081	1091 (2.54-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	235	22% 74% 20% • 5%
1	B	235	25% 77% 17% • 5%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 3499 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Golgi reassembly-stacking protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	223	Total	C	N	O	S	0	0	0
			1713	1080	294	334	5			
1	B	223	Total	C	N	O	S	0	0	0
			1713	1080	294	334	5			

There are 56 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-26	MET	-	expression tag	UNP Q99JX3
A	-25	SER	-	expression tag	UNP Q99JX3
A	-24	TYR	-	expression tag	UNP Q99JX3
A	-23	TYR	-	expression tag	UNP Q99JX3
A	-22	HIS	-	expression tag	UNP Q99JX3
A	-21	HIS	-	expression tag	UNP Q99JX3
A	-20	HIS	-	expression tag	UNP Q99JX3
A	-19	HIS	-	expression tag	UNP Q99JX3
A	-18	HIS	-	expression tag	UNP Q99JX3
A	-17	HIS	-	expression tag	UNP Q99JX3
A	-16	LEU	-	expression tag	UNP Q99JX3
A	-15	GLU	-	expression tag	UNP Q99JX3
A	-14	SER	-	expression tag	UNP Q99JX3
A	-13	THR	-	expression tag	UNP Q99JX3
A	-12	SER	-	expression tag	UNP Q99JX3
A	-11	LEU	-	expression tag	UNP Q99JX3
A	-10	TYR	-	expression tag	UNP Q99JX3
A	-9	LYS	-	expression tag	UNP Q99JX3
A	-8	LYS	-	expression tag	UNP Q99JX3
A	-7	ALA	-	expression tag	UNP Q99JX3
A	-6	GLY	-	expression tag	UNP Q99JX3
A	-5	PHE	-	expression tag	UNP Q99JX3
A	-4	LEU	-	expression tag	UNP Q99JX3
A	-3	VAL	-	expression tag	UNP Q99JX3
A	-2	PRO	-	expression tag	UNP Q99JX3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	ARG	-	expression tag	UNP Q99JX3
A	0	GLY	-	expression tag	UNP Q99JX3
A	1	SER	-	expression tag	UNP Q99JX3
B	-26	MET	-	expression tag	UNP Q99JX3
B	-25	SER	-	expression tag	UNP Q99JX3
B	-24	TYR	-	expression tag	UNP Q99JX3
B	-23	TYR	-	expression tag	UNP Q99JX3
B	-22	HIS	-	expression tag	UNP Q99JX3
B	-21	HIS	-	expression tag	UNP Q99JX3
B	-20	HIS	-	expression tag	UNP Q99JX3
B	-19	HIS	-	expression tag	UNP Q99JX3
B	-18	HIS	-	expression tag	UNP Q99JX3
B	-17	HIS	-	expression tag	UNP Q99JX3
B	-16	LEU	-	expression tag	UNP Q99JX3
B	-15	GLU	-	expression tag	UNP Q99JX3
B	-14	SER	-	expression tag	UNP Q99JX3
B	-13	THR	-	expression tag	UNP Q99JX3
B	-12	SER	-	expression tag	UNP Q99JX3
B	-11	LEU	-	expression tag	UNP Q99JX3
B	-10	TYR	-	expression tag	UNP Q99JX3
B	-9	LYS	-	expression tag	UNP Q99JX3
B	-8	LYS	-	expression tag	UNP Q99JX3
B	-7	ALA	-	expression tag	UNP Q99JX3
B	-6	GLY	-	expression tag	UNP Q99JX3
B	-5	PHE	-	expression tag	UNP Q99JX3
B	-4	LEU	-	expression tag	UNP Q99JX3
B	-3	VAL	-	expression tag	UNP Q99JX3
B	-2	PRO	-	expression tag	UNP Q99JX3
B	-1	ARG	-	expression tag	UNP Q99JX3
B	0	GLY	-	expression tag	UNP Q99JX3
B	1	SER	-	expression tag	UNP Q99JX3


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	36	Total O 36 36	0	0
2	B	37	Total O 37 37	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Golgi reassembly-stacking protein 2



• Molecule 1: Golgi reassembly-stacking protein 2



4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	132.31Å 132.31Å 221.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	93.56 – 2.55 93.56 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.4 (93.56-2.55) 94.2 (93.56-2.55)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.22 (at 2.55Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.261 , 0.311 0.273 , 0.317	Depositor DCC
R_{free} test set	2000 reflections (6.16%)	wwPDB-VP
Wilson B-factor (Å ²)	45.2	Xtriage
Anisotropy	0.133	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 64.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	3499	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 39.89 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.9694e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/1752	0.96	3/2377 (0.1%)
1	B	0.47	0/1752	0.91	7/2377 (0.3%)
All	All	0.49	0/3504	0.94	10/4754 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	152	LEU	N-CA-C	-8.67	104.74	114.62
1	B	9	ILE	N-CA-C	7.23	115.06	107.76
1	B	179	THR	CA-C-N	6.49	126.52	119.90
1	B	179	THR	C-N-CA	6.49	126.52	119.90
1	B	202	ILE	CA-C-N	5.77	125.72	119.78
1	B	202	ILE	C-N-CA	5.77	125.72	119.78
1	B	142	VAL	N-CA-C	-5.72	102.20	109.58
1	B	152	LEU	CA-CB-CG	5.09	134.13	116.30
1	A	202	ILE	CA-C-N	5.03	124.96	119.78
1	A	202	ILE	C-N-CA	5.03	124.96	119.78

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	49	ASN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1713	0	1673	40	0
1	B	1713	0	1673	28	0
2	A	36	0	0	3	0
2	B	37	0	0	7	0
All	All	3499	0	3346	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:50:LYS:HE3	1:A:55:LEU:H	1.20	1.06
1:A:7:VAL:HG23	1:A:8:GLU:H	1.36	0.88
1:A:17:TYR:OH	1:A:50:LYS:NZ	2.17	0.76
1:B:6:SER:OG	1:B:7:VAL:N	2.16	0.75
1:A:6:SER:OG	1:A:7:VAL:N	2.14	0.75
1:A:63:VAL:HG13	1:A:64:GLU:HG2	1.73	0.69
1:A:159:LYS:HG2	1:A:160:PRO:HD2	1.74	0.68
1:B:171:ASP:OD1	2:B:301:HOH:O	2.11	0.68
1:A:50:LYS:HE3	1:A:55:LEU:N	2.03	0.66
1:B:159:LYS:NZ	2:B:306:HOH:O	2.27	0.66
1:A:50:LYS:HG2	1:A:52:ASN:H	1.62	0.64
1:A:201:ARG:NH1	2:A:304:HOH:O	2.30	0.64
1:B:154:GLU:OE1	1:B:154:GLU:N	2.29	0.63
1:A:91:GLY:N	2:A:302:HOH:O	2.26	0.63
1:A:112:VAL:O	2:A:301:HOH:O	2.16	0.63
1:B:142:VAL:HG22	1:B:143:MET:HG2	1.82	0.61
1:A:5:GLN:CD	1:A:6:SER:HB2	2.27	0.59
1:B:-2:PRO:O	1:B:1:SER:OG	2.16	0.59
1:A:155:THR:HG23	1:A:156:HIS:ND1	2.20	0.57
1:A:50:LYS:NZ	1:A:52:ASN:O	2.25	0.56
1:B:63:VAL:HG23	1:B:64:GLU:HG2	1.85	0.56
1:B:148:ASP:OD1	2:B:302:HOH:O	2.17	0.56
1:B:201:ARG:HD2	2:B:325:HOH:O	2.06	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:50:LYS:HE2	1:A:52:ASN:OD1	2.06	0.55
1:B:101:ARG:NH1	2:B:307:HOH:O	2.31	0.55
1:A:154:GLU:OE1	1:A:154:GLU:N	2.30	0.54
1:A:147:GLU:HA	1:A:147:GLU:OE1	2.07	0.54
1:A:113:TRP:CH2	1:A:148:ASP:HA	2.43	0.52
1:A:90:TRP:CH2	1:A:92:GLY:HA3	2.45	0.52
1:B:140:ASP:HB2	1:B:162:LYS:HB2	1.92	0.52
1:A:5:GLN:NE2	1:A:6:SER:HB2	2.24	0.51
1:B:120:SER:O	2:B:304:HOH:O	2.19	0.51
1:A:69:MET:HE3	1:A:71:ILE:HD11	1.93	0.51
1:A:110:GLU:HA	1:A:201:ARG:NH1	2.26	0.51
1:B:162:LYS:HZ2	1:B:177:ILE:HD13	1.75	0.51
1:A:149:LEU:HD13	1:A:153:ILE:HD11	1.92	0.51
1:B:90:TRP:CH2	1:B:92:GLY:HA3	2.46	0.50
1:B:162:LYS:NZ	1:B:177:ILE:HD13	2.25	0.50
1:B:37:PHE:CD1	1:B:75:LYS:HB2	2.47	0.50
1:A:7:VAL:CG2	1:A:8:GLU:H	2.17	0.50
1:B:184:TRP:CZ2	1:B:186:GLY:HA3	2.47	0.50
1:A:110:GLU:HA	1:A:201:ARG:HH12	1.76	0.49
1:B:161:LEU:O	1:B:177:ILE:HD12	2.13	0.49
1:A:142:VAL:HA	1:A:143:MET:C	2.38	0.48
1:A:17:TYR:HE2	1:A:50:LYS:HD3	1.79	0.47
1:A:129:LEU:O	1:A:130:ARG:NH1	2.38	0.47
1:A:145:GLU:HB3	1:A:146:SER:H	1.42	0.47
1:A:206:PRO:HB3	1:B:130:ARG:HG3	1.97	0.46
1:A:151:SER:O	1:A:151:SER:OG	2.33	0.46
1:B:1:SER:HB3	1:B:201:ARG:NH1	2.31	0.45
1:A:7:VAL:HG23	1:A:8:GLU:N	2.16	0.44
1:A:122:SER:HB2	1:A:191:GLY:HA3	1.99	0.44
1:B:178:ILE:O	1:B:180:PRO:HD3	2.18	0.44
1:B:1:SER:HB3	1:B:201:ARG:HH12	1.82	0.44
1:A:184:TRP:CZ2	1:A:186:GLY:HA3	2.53	0.44
1:B:186:GLY:O	2:B:305:HOH:O	2.21	0.44
1:B:140:ASP:HB2	1:B:162:LYS:CB	2.48	0.43
1:A:149:LEU:CD1	1:A:153:ILE:HD11	2.48	0.43
1:A:50:LYS:HZ3	1:A:52:ASN:C	2.18	0.43
1:B:130:ARG:HA	1:B:130:ARG:HD3	1.87	0.42
1:B:149:LEU:O	1:B:152:LEU:HD13	2.20	0.41
1:A:1:SER:O	1:A:3:SER:N	2.53	0.41
1:A:148:ASP:N	1:A:148:ASP:OD1	2.53	0.41
1:B:33:LEU:HD12	1:B:98:VAL:HG21	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:160:PRO:HB2	1:B:177:ILE:HD11	2.02	0.41
1:A:50:LYS:HE2	1:A:54:THR:H	1.86	0.41
1:A:130:ARG:HA	1:A:130:ARG:HD3	1.86	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	221/235 (94%)	198 (90%)	21 (10%)	2 (1%)	14	20
1	B	221/235 (94%)	204 (92%)	17 (8%)	0	100	100
All	All	442/470 (94%)	402 (91%)	38 (9%)	2 (0%)	24	34

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2	GLY
1	A	7	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/200 (94%)	183 (97%)	5 (3%)	39	58

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	188/200 (94%)	179 (95%)	9 (5%)	23	35
All	All	376/400 (94%)	362 (96%)	14 (4%)	30	45

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	GLU
1	A	43	ILE
1	A	67	VAL
1	A	147	GLU
1	A	152	LEU
1	B	24	GLU
1	B	30	ARG
1	B	67	VAL
1	B	106	ASP
1	B	142	VAL
1	B	145	GLU
1	B	147	GLU
1	B	152	LEU
1	B	153	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	GLN
1	A	88	ASN
1	A	111	ASN
1	B	23	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	223/235 (94%)	1.23	51 (22%) 2 2	28, 55, 125, 171	0
1	B	223/235 (94%)	1.39	58 (26%) 1 1	34, 58, 137, 189	0
All	All	446/470 (94%)	1.31	109 (24%) 2 1	28, 57, 136, 189	0

All (109) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	158	ALA	10.4
1	A	142	VAL	8.7
1	A	7	VAL	8.3
1	B	7	VAL	7.7
1	A	146	SER	7.7
1	A	140	ASP	7.2
1	B	142	VAL	6.7
1	B	154	GLU	6.5
1	A	107	GLY	6.3
1	A	-14	SER	6.2
1	B	106	ASP	6.2
1	A	141	THR	6.2
1	B	6	SER	6.2
1	B	159	LYS	6.0
1	A	147	GLU	5.9
1	B	144	ASN	5.8
1	B	146	SER	5.8
1	B	-14	SER	5.6
1	B	141	THR	5.5
1	B	152	LEU	5.3
1	A	151	SER	5.2
1	B	4	SER	5.0
1	A	6	SER	5.0
1	A	5	GLN	5.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	159	LYS	4.9
1	B	151	SER	4.9
1	B	145	GLU	4.9
1	A	2	GLY	4.7
1	B	140	ASP	4.6
1	A	155	THR	4.5
1	A	144	ASN	4.5
1	A	106	ASP	4.5
1	B	3	SER	4.4
1	A	148	ASP	4.4
1	B	148	ASP	4.3
1	B	13	GLY	4.2
1	B	162	LYS	4.1
1	A	158	ALA	4.1
1	B	107	GLY	4.0
1	B	155	THR	4.0
1	B	150	PHE	3.9
1	A	143	MET	3.9
1	B	143	MET	3.9
1	B	14	THR	3.9
1	A	51	ASP	3.9
1	B	160	PRO	3.8
1	B	5	GLN	3.8
1	B	156	HIS	3.7
1	A	145	GLU	3.7
1	B	153	ILE	3.7
1	A	49	ASN	3.7
1	B	147	GLU	3.6
1	A	12	GLY	3.6
1	A	56	LYS	3.4
1	A	1	SER	3.4
1	A	157	GLU	3.4
1	A	24	GLU	3.3
1	B	8	GLU	3.3
1	B	15	GLU	3.3
1	B	177	ILE	3.3
1	B	179	THR	3.2
1	A	3	SER	3.2
1	A	182	SER	3.2
1	A	14	THR	3.1
1	B	50	LYS	3.1
1	A	4	SER	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	13	GLY	3.1
1	A	15	GLU	3.0
1	A	162	LYS	3.0
1	B	12	GLY	3.0
1	B	63	VAL	3.0
1	A	63	VAL	3.0
1	B	-13	THR	3.0
1	B	139	ALA	3.0
1	A	154	GLU	2.9
1	A	105	PHE	2.9
1	B	185	GLY	2.9
1	A	50	LYS	2.9
1	B	149	LEU	2.7
1	A	8	GLU	2.7
1	A	150	PHE	2.6
1	B	157	GLU	2.6
1	A	11	GLY	2.6
1	A	160	PRO	2.6
1	B	10	PRO	2.6
1	B	91	GLY	2.5
1	B	24	GLU	2.5
1	A	93	GLN	2.5
1	B	93	GLN	2.4
1	A	113	TRP	2.4
1	A	189	SER	2.4
1	B	102	PHE	2.4
1	A	90	TRP	2.4
1	B	110	GLU	2.3
1	B	11	GLY	2.3
1	B	0	GLY	2.3
1	B	9	ILE	2.3
1	B	175	GLU	2.3
1	A	52	ASN	2.2
1	B	108	ALA	2.2
1	A	29	HIS	2.1
1	B	109	ASN	2.1
1	B	2	GLY	2.1
1	A	110	GLU	2.1
1	A	152	LEU	2.1
1	B	30	ARG	2.1
1	B	56	LYS	2.1
1	B	82	ALA	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	139	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.