



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 07:02 AM UTC

PDB ID : 6GMN / pdb_00006gmn
Title : pVHL:EloB:EloC in complex with methyl 4H-furo[3,2-b]pyrrole-5-carboxylate
Authors : Van Molle, I.; Lucas, X.; Ciulli, A.
Deposited on : 2018-05-27
Resolution : 1.94 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	104	7% 87% 12% .
2	B	97	8% 82% 7% 10%
2	E	97	9% 79% 10% 10%
2	H	97	10% 78% 10% 11%
2	K	97	6% 73% 14% .. 10%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	162	
3	F	162	
3	L	162	
4	D	104	
4	G	104	
4	J	104	
5	I	162	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 11441 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Elongin-B.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	103	Total	As	C	N	O	S	0	0	0
			800	1	509	136	150	4			

- Molecule 2 is a protein called Elongin-C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	87	Total	C	N	O	S	0	0	0
			688	445	110	127	6			
2	E	87	Total	C	N	O	S	0	1	0
			695	448	112	129	6			
2	H	86	Total	C	N	O	S	0	0	0
			687	443	110	128	6			
2	K	87	Total	C	N	O	S	0	0	0
			696	449	111	129	7			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	16	MET	-	initiating methionine	UNP Q15369
E	16	MET	-	initiating methionine	UNP Q15369
H	16	MET	-	initiating methionine	UNP Q15369
K	16	MET	-	initiating methionine	UNP Q15369

- Molecule 3 is a protein called von Hippel-Lindau disease tumor suppressor.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	137	Total	As	C	N	O	S	0	0	0
			1099	1	702	195	199	2			
3	F	142	Total	As	C	N	O	S	0	3	0
			1165	1	742	210	210	2			
3	L	144	Total	As	C	N	O	S	0	1	0
			1173	1	746	212	212	2			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	52	GLY	-	expression tag	UNP P40337
C	53	SER	-	expression tag	UNP P40337
F	52	GLY	-	expression tag	UNP P40337
F	53	SER	-	expression tag	UNP P40337
L	52	GLY	-	expression tag	UNP P40337
L	53	SER	-	expression tag	UNP P40337

- Molecule 4 is a protein called Elongin-B.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	D	100	Total	As	C	N	O	S	0	0	0
			779	2	496	133	145	3			
4	G	103	Total	As	C	N	O	S	0	0	0
			809	2	513	135	154	5			
4	J	103	Total	As	C	N	O	S	0	0	0
			809	2	514	136	152	5			

- Molecule 5 is a protein called von Hippel-Lindau disease tumor suppressor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	I	145	Total	C	N	O	S	0	0	0
			1158	737	208	211	2			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
I	52	GLY	-	expression tag	UNP P40337
I	53	SER	-	expression tag	UNP P40337

- Molecule 6 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		
6	C	1	Total	C	O	0	0
			4	2	2		
6	F	1	Total	C	O	0	0
			4	2	2		
6	I	1	Total	C	O	0	0
			4	2	2		
6	L	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is methyl 4 {H}-furo[3,2-b]pyrrole-5-carboxylate (CCD ID: F4E) (formula: C₈H₇NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	B	1	Total	C	N	O	0	0
			12	8	1	3		
7	K	1	Total	C	N	O	0	0
			12	8	1	3		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	103	Total	O	0	0
			103	103		
8	B	52	Total	O	0	0
			52	52		
8	C	59	Total	O	0	0
			59	59		
8	D	45	Total	O	0	0
			45	45		
8	E	46	Total	O	0	0
			46	46		
8	F	54	Total	O	0	0
			54	54		
8	G	59	Total	O	0	0
			59	59		
8	H	46	Total	O	0	0
			46	46		
8	I	78	Total	O	0	0
			78	78		
8	J	119	Total	O	0	0
			119	119		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	K	66	Total 66	O 66	0	0
8	L	112	Total 112	O 112	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Elongin-B





- Molecule 2: Elongin-C




- Molecule 2: Elongin-C





- Molecule 2: Elongin-C




- Molecule 2: Elongin-C




- Molecule 3: von Hippel-Lindau disease tumor suppressor




- Molecule 3: von Hippel-Lindau disease tumor suppressor




- Molecule 3: von Hippel-Lindau disease tumor suppressor




- Molecule 4: Elongin-B




- Molecule 4: Elongin-B





- Molecule 4: Elongin-B



- Molecule 5: von Hippel-Lindau disease tumor suppressor





4 Data and refinement statistics

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	93.71Å 93.71Å 364.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.47 – 1.94 46.47 – 1.94	Depositor EDS
% Data completeness (in resolution range)	99.2 (46.47-1.94) 99.2 (46.47-1.94)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.27 (at 1.94Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.203 , 0.244 0.204 , 0.245	Depositor DCC
R_{free} test set	6006 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	36.5	Xtriage
Anisotropy	0.350	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 49.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11441	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 30.51 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.2911e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, CAS, F4E

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.42	0/806	0.65	0/1090
2	B	0.42	0/702	0.59	0/948
2	E	0.37	0/709	0.62	1/956 (0.1%)
2	H	0.38	0/701	0.57	0/945
2	K	0.43	0/710	0.67	2/957 (0.2%)
3	C	0.44	0/1118	0.68	3/1527 (0.2%)
3	F	0.48	1/1186 (0.1%)	0.90	10/1622 (0.6%)
3	L	0.39	0/1193	0.58	0/1628
4	D	0.34	0/773	0.74	4/1042 (0.4%)
4	G	0.39	0/805	0.55	0/1087
4	J	0.43	0/805	0.64	0/1086
5	I	0.42	1/1187 (0.1%)	0.60	0/1624
All	All	0.41	2/10695 (0.0%)	0.66	20/14512 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	C	0	1
3	F	0	1
4	G	0	1
All	All	0	3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	I	77	CYS	CB-SG	5.58	1.99	1.81
3	F	134	GLU	CG-CD	-5.06	1.39	1.52

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	79	ARG	CB-CG-CD	-11.25	85.42	111.30
3	F	134	GLU	CB-CA-C	8.77	124.59	109.24
4	D	100	PRO	N-CA-CB	8.46	112.14	103.25
3	F	134	GLU	CA-CB-CG	8.20	130.49	114.10
3	F	79	ARG	CD-NE-CZ	7.70	135.18	124.40

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	C	185	TYR	Sidechain
3	F	134	GLU	Sidechain
4	G	60	CAS	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	800	0	793	11	0
2	B	688	0	683	8	0
2	E	695	0	687	9	0
2	H	687	0	688	14	0
2	K	696	0	699	20	0
3	C	1099	0	1063	13	0
3	F	1165	0	1129	10	0
3	L	1173	0	1149	8	1
4	D	779	0	764	11	0
4	G	809	0	793	15	0
4	J	809	0	800	7	0
5	I	1158	0	1139	15	0
6	A	4	0	3	1	0
6	C	4	0	3	0	0
6	F	4	0	3	0	0
6	I	4	0	3	0	0
6	L	4	0	3	0	0
7	B	12	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	K	12	0	0	0	0
8	A	103	0	0	1	0
8	B	52	0	0	0	0
8	C	59	0	0	2	0
8	D	45	0	0	1	0
8	E	46	0	0	2	0
8	F	54	0	0	1	0
8	G	59	0	0	0	0
8	H	46	0	0	2	0
8	I	78	0	0	3	0
8	J	119	0	0	1	0
8	K	66	0	0	3	0
8	L	112	0	0	0	1
All	All	11441	0	10402	118	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 118 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:J:72:PRO:HD2	2:K:75:MET:HE3	1.32	1.12
4:G:25:PHE:CZ	4:G:29:ARG:NH1	2.19	1.11
1:A:72:PRO:HD2	2:B:75:MET:HE3	1.33	1.07
4:G:25:PHE:HZ	4:G:29:ARG:NH1	1.60	0.99
3:F:182:ARG:O	3:F:186:GLU:OE1	1.83	0.94

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:195:GLN:OE1	8:L:431:HOH:O[5_555]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	100/104 (96%)	95 (95%)	4 (4%)	1 (1%)	12	5
2	B	83/97 (86%)	82 (99%)	1 (1%)	0	100	100
2	E	84/97 (87%)	82 (98%)	1 (1%)	1 (1%)	10	3
2	H	82/97 (84%)	82 (100%)	0	0	100	100
2	K	83/97 (86%)	81 (98%)	0	2 (2%)	4	0
3	C	132/162 (82%)	127 (96%)	4 (3%)	1 (1%)	16	7
3	F	142/162 (88%)	137 (96%)	5 (4%)	0	100	100
3	L	142/162 (88%)	135 (95%)	6 (4%)	1 (1%)	18	9
4	D	94/104 (90%)	87 (93%)	3 (3%)	4 (4%)	2	0
4	G	99/104 (95%)	95 (96%)	4 (4%)	0	100	100
4	J	99/104 (95%)	96 (97%)	3 (3%)	0	100	100
5	I	143/162 (88%)	139 (97%)	4 (3%)	0	100	100
All	All	1283/1452 (88%)	1238 (96%)	35 (3%)	10 (1%)	16	7

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	84	THR
4	D	98	GLU
4	D	100	PRO
2	K	88	THR
2	E	58	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/91 (93%)	85 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	76/86 (88%)	75 (99%)	1 (1%)	61	54
2	E	77/86 (90%)	77 (100%)	0	100	100
2	H	78/86 (91%)	77 (99%)	1 (1%)	61	54
2	K	79/86 (92%)	79 (100%)	0	100	100
3	C	121/147 (82%)	121 (100%)	0	100	100
3	F	128/147 (87%)	126 (98%)	2 (2%)	55	46
3	L	130/147 (88%)	129 (99%)	1 (1%)	73	71
4	D	81/90 (90%)	81 (100%)	0	100	100
4	G	86/90 (96%)	86 (100%)	0	100	100
4	J	86/90 (96%)	86 (100%)	0	100	100
5	I	130/148 (88%)	129 (99%)	1 (1%)	73	71
All	All	1157/1294 (89%)	1151 (100%)	6 (0%)	84	82

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	72	LYS
5	I	159	LYS
3	L	188	LEU
3	F	120[A]	ARG
2	B	72	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
5	I	191	HIS
2	K	61	ASN
3	L	115	HIS
2	K	85	ASN
2	E	61	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

10 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	CAS	D	89	4	5,8,9	1.15	0	1,9,11	0.77	0
4	CAS	D	60	4	5,8,9	0.96	0	1,9,11	0.38	0
1	CAS	A	60	1	5,8,9	0.96	0	1,9,11	0.90	0
3	CAS	L	77	3	5,8,9	0.74	0	1,9,11	2.06	1 (100%)
4	CAS	J	60	4	5,8,9	0.90	0	1,9,11	0.14	0
3	CAS	F	77	3	5,8,9	0.85	0	1,9,11	1.28	0
4	CAS	G	60	4	5,8,9	0.94	0	1,9,11	0.72	0
4	CAS	G	89	4	5,8,9	1.02	0	1,9,11	0.67	0
3	CAS	C	77	3	5,8,9	1.08	0	1,9,11	1.47	0
4	CAS	J	89	4	5,8,9	1.11	0	1,9,11	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CAS	D	89	4	-	0/0/7/9	-
4	CAS	D	60	4	-	0/0/7/9	-
1	CAS	A	60	1	-	0/0/7/9	-
3	CAS	L	77	3	-	0/0/7/9	-
4	CAS	J	60	4	-	0/0/7/9	-
3	CAS	F	77	3	-	0/0/7/9	-
4	CAS	G	60	4	-	0/0/7/9	-
4	CAS	G	89	4	-	0/0/7/9	-
3	CAS	C	77	3	-	0/0/7/9	-
4	CAS	J	89	4	-	0/0/7/9	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	77	CAS	CB-CA-C	2.06	116.38	110.80

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	60	CAS	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	ACT	L	301	-	3,3,3	1.82	1 (33%)	3,3,3	1.41	0
7	F4E	B	201	-	12,13,13	1.59	3 (25%)	15,18,18	2.27	6 (40%)
6	ACT	A	201	-	3,3,3	1.52	1 (33%)	3,3,3	1.48	0
6	ACT	F	301	-	3,3,3	1.49	1 (33%)	3,3,3	1.31	0
6	ACT	I	301	-	3,3,3	1.55	1 (33%)	3,3,3	1.35	0
7	F4E	K	201	-	12,13,13	1.55	2 (16%)	15,18,18	2.39	7 (46%)
6	ACT	C	301	-	3,3,3	1.72	1 (33%)	3,3,3	1.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	F4E	K	201	-	-	5/6/6/6	0/2/2/2
7	F4E	B	201	-	-	4/6/6/6	0/2/2/2

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	B	201	F4E	O02-C01	-3.10	1.38	1.45
7	K	201	F4E	O02-C01	-3.04	1.38	1.45
6	C	301	ACT	CH3-C	2.42	1.58	1.49
6	L	301	ACT	CH3-C	2.35	1.58	1.49
6	A	201	ACT	CH3-C	2.27	1.58	1.49

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	K	201	F4E	O02-C03-C05	6.33	119.87	111.97
7	B	201	F4E	O02-C03-C05	5.54	118.89	111.97
7	B	201	F4E	C06-C05-C03	-3.42	126.09	131.36
7	K	201	F4E	C06-C05-C03	-3.22	126.39	131.36
7	B	201	F4E	C01-O02-C03	2.71	120.88	115.85

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	B	201	F4E	O02-C03-C05-N12
7	B	201	F4E	O02-C03-C05-C06
7	B	201	F4E	O04-C03-C05-N12
7	B	201	F4E	O04-C03-C05-C06
7	K	201	F4E	O02-C03-C05-N12

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	201	ACT	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	102/104 (98%)	0.28	7 (6%) 23 26	27, 38, 72, 103	1 (0%)
2	B	87/97 (89%)	0.45	8 (9%) 14 16	29, 40, 88, 102	0
2	E	87/97 (89%)	0.78	9 (10%) 12 13	22, 48, 86, 97	2 (2%)
2	H	86/97 (88%)	0.77	10 (11%) 9 10	33, 47, 91, 115	0
2	K	87/97 (89%)	0.35	6 (6%) 23 26	27, 39, 80, 95	0
3	C	136/162 (83%)	0.60	12 (8%) 15 17	30, 46, 79, 116	0
3	F	141/162 (87%)	0.63	10 (7%) 22 25	18, 48, 81, 119	3 (2%)
3	L	143/162 (88%)	0.22	6 (4%) 40 46	14, 37, 72, 90	1 (0%)
4	D	98/104 (94%)	0.96	11 (11%) 10 11	31, 50, 84, 102	0
4	G	101/104 (97%)	0.54	7 (6%) 23 26	32, 48, 76, 80	0
4	J	101/104 (97%)	0.05	2 (1%) 65 72	26, 35, 61, 73	0
5	I	145/162 (89%)	0.54	14 (9%) 13 15	30, 43, 73, 96	0
All	All	1314/1452 (90%)	0.51	102 (7%) 19 21	14, 44, 80, 119	7 (0%)

The worst 5 of 102 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	201	LEU	6.9
4	D	102	VAL	6.5
3	C	185	TYR	5.9
2	H	110	LEU	5.2
3	L	142	VAL	4.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CAS	D	89	9/10	0.84	0.15	44,54,124,125	9
4	CAS	G	89	9/10	0.89	0.15	29,36,125,126	0
1	CAS	A	60	9/10	0.90	0.18	26,30,103,108	0
4	CAS	D	60	9/10	0.92	0.15	26,35,85,92	0
4	CAS	J	60	9/10	0.92	0.17	28,33,94,94	9
3	CAS	F	77	9/10	0.93	0.12	33,37,82,94	9
3	CAS	L	77	9/10	0.93	0.15	24,31,102,115	0
4	CAS	J	89	9/10	0.94	0.12	26,31,94,102	0
3	CAS	C	77	9/10	0.94	0.12	32,37,85,104	9
4	CAS	G	60	9/10	0.97	0.10	38,43,76,87	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	F4E	K	201	12/12	0.66	0.24	108,115,118,119	0
7	F4E	B	201	12/12	0.74	0.23	90,103,107,111	0
6	ACT	I	301	4/4	0.74	0.17	56,61,65,66	0
6	ACT	C	301	4/4	0.80	0.17	61,62,71,72	0
6	ACT	F	301	4/4	0.83	0.14	52,61,61,67	0
6	ACT	A	201	4/4	0.84	0.14	51,64,64,78	0
6	ACT	L	301	4/4	0.85	0.18	42,70,71,75	0

6.5 Other polymers [i](#)

There are no such residues in this entry.