



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 04:12 PM UTC

PDB ID : 3GNV / pdb_00003gnv
Title : HCV NS5B polymerase in complex with 1,5 benzodiazepine inhibitor 1b
Authors : De Bondt, H.; Nyanguile, O.
Deposited on : 2009-03-18
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	581	
1	B	581	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8957 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA-directed RNA polymerase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	563	Total	C	N	O	S	20	0	0
			4384	2760	775	817	32			
1	B	553	Total	C	N	O	S	5	0	0
			4311	2716	762	801	32			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	expression tag	UNP O92972
A	-1	ALA	-	expression tag	UNP O92972
A	0	SER	-	expression tag	UNP O92972
A	571	LEU	-	expression tag	UNP O92972
A	572	GLU	-	expression tag	UNP O92972
A	573	HIS	-	expression tag	UNP O92972
A	574	HIS	-	expression tag	UNP O92972
A	575	HIS	-	expression tag	UNP O92972
A	576	HIS	-	expression tag	UNP O92972
A	577	HIS	-	expression tag	UNP O92972
A	578	HIS	-	expression tag	UNP O92972
B	-2	MET	-	expression tag	UNP O92972
B	-1	ALA	-	expression tag	UNP O92972
B	0	SER	-	expression tag	UNP O92972
B	571	LEU	-	expression tag	UNP O92972
B	572	GLU	-	expression tag	UNP O92972
B	573	HIS	-	expression tag	UNP O92972
B	574	HIS	-	expression tag	UNP O92972
B	575	HIS	-	expression tag	UNP O92972
B	576	HIS	-	expression tag	UNP O92972
B	577	HIS	-	expression tag	UNP O92972
B	578	HIS	-	expression tag	UNP O92972


- Molecule 2 is (11R)-10-acetyl-11-[4-(benzyloxy)-2-chlorophenyl]-6-hydroxy-3,3-dimethyl-2,3,4,5,10,11-hexahydro-1H-dibenzo[b,e][1,4]diazepin-1-one (CCD ID: XNZ) (formula:

$C_{30}H_{29}ClN_2O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Cl	N	O	0	0
			37	30	1	2	4		
2	B	1	Total	C	Cl	N	O	0	0
			37	30	1	2	4		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	111	Total	O	0	0
			111	111		
4	B	65	Total	O	0	0
			65	65		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RNA-directed RNA polymerase



• Molecule 1: RNA-directed RNA polymerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	106.29Å 107.97Å 133.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	83.33 – 2.75 83.33 – 2.75	Depositor EDS
% Data completeness (in resolution range)	100.0 (83.33-2.75) 100.0 (83.33-2.75)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 2.73Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.174 , 0.234 0.198 , 0.248	Depositor DCC
R_{free} test set	2037 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	32.0	Xtriage
Anisotropy	0.549	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 33.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.028 for k,h,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8957	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: XNZ, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	0/4480	0.85	0/6080
1	B	0.65	0/4404	0.86	0/5974
All	All	0.67	0/8884	0.86	0/12054

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4384	0	4397	38	0
1	B	4311	0	4322	35	0
2	A	37	0	28	0	0
2	B	37	0	28	0	0
3	A	6	0	8	0	0
3	B	6	0	8	0	0
4	A	111	0	0	3	0
4	B	65	0	0	4	0
All	All	8957	0	8791	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:198:LYS:HD2	4:B:631:HOH:O	1.85	0.74
1:B:30:LEU:HD12	1:B:31:LEU:HB2	1.76	0.67
1:A:32:ARG:HD3	1:A:493:GLY:O	1.95	0.67
1:B:530:VAL:O	1:B:533:LYS:NZ	2.28	0.67
1:A:236:GLU:OE1	1:A:280:ARG:NH2	2.27	0.66
1:A:546:ASP:OD1	1:A:548:SER:HB2	1.97	0.65
1:A:260:LEU:O	1:A:277:ARG:NH2	2.32	0.63
1:A:180:SER:HB3	1:A:555:TYR:OH	2.00	0.62
1:A:499:THR:HG23	1:B:128:GLU:HG2	1.82	0.60
1:B:46:SER:HA	1:B:49:GLN:HE21	1.67	0.60
1:B:541:ALA:HA	4:B:626:HOH:O	2.04	0.58
1:B:368:SER:HB2	4:B:620:HOH:O	2.04	0.56
1:A:438:GLN:HB3	1:A:441:LYS:HZ2	1.70	0.56
1:A:277:ARG:NH1	4:A:663:HOH:O	2.39	0.55
1:A:219:TYR:HB3	1:A:320:LEU:HD23	1.89	0.54
1:A:327:ALA:O	1:A:331:GLU:HG3	2.08	0.54
1:B:219:TYR:HB3	1:B:320:LEU:HD23	1.89	0.53
1:A:364:THR:HA	1:A:368:SER:O	2.08	0.53
1:A:187:MET:HE3	1:A:296:TYR:CD2	2.44	0.53
1:A:172:LYS:NZ	4:A:643:HOH:O	2.33	0.53
1:A:280:ARG:HD2	1:A:291:ASN:OD1	2.08	0.52
1:B:189:SER:HA	1:B:194:GLN:HE22	1.75	0.52
1:B:280:ARG:HD2	1:B:291:ASN:OD1	2.10	0.52
1:A:48:ARG:HG2	1:A:159:LEU:HG	1.93	0.50
1:B:187:MET:HE1	1:B:292:THR:HG22	1.94	0.49
1:B:46:SER:HA	1:B:49:GLN:NE2	2.27	0.49
1:A:144:VAL:HG21	1:A:397:TRP:CG	2.48	0.49
1:B:419:LEU:HD21	1:B:497:LEU:HD13	1.94	0.49
1:B:183:PRO:HG3	1:B:289:CYS:SG	2.53	0.49
1:B:30:LEU:HD12	1:B:30:LEU:C	2.38	0.48
1:B:197:PRO:O	1:B:201:VAL:HG23	2.13	0.48
1:A:223:CYS:O	1:A:224:PHE:C	2.55	0.48
1:A:24:ASN:HB3	1:A:27:SER:OG	2.15	0.47
1:B:82:LEU:HD13	1:B:249:ALA:HB2	1.97	0.47
1:B:359:ASP:HB3	1:B:362:LEU:HD22	1.96	0.47
1:A:187:MET:HE1	1:A:292:THR:HG22	1.97	0.46
1:A:485:VAL:O	1:A:489:LEU:HG	2.16	0.46
1:A:438:GLN:HB3	1:A:441:LYS:NZ	2.31	0.46
1:B:52:VAL:HG22	1:B:226:SER:OG	2.16	0.45
1:B:237:GLU:OE2	1:B:254:ARG:HD2	2.17	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:461:GLN:HB2	1:A:542:ALA:HA	1.98	0.45
1:B:5:THR:O	1:B:275:GLY:HA3	2.17	0.44
1:B:465:ARG:NH1	1:B:545:LEU:O	2.51	0.44
1:A:446:GLN:HA	1:A:450:ALA:O	2.17	0.44
1:A:234:ARG:CZ	1:A:262:ILE:HD11	2.48	0.44
1:B:120:ARG:O	1:B:124:GLU:HG2	2.16	0.44
1:A:130:THR:HG22	1:A:131:GLU:HG3	1.99	0.44
1:A:416:ALA:N	1:A:417:PRO:CD	2.81	0.43
1:B:211:LYS:HE2	1:B:324:CYS:HA	2.00	0.43
1:A:57:LEU:HD13	1:A:57:LEU:HA	1.84	0.43
1:A:197:PRO:O	1:A:201:VAL:HG23	2.18	0.43
1:B:18:GLU:HG2	1:B:401:ARG:CZ	2.48	0.43
1:B:544:GLN:HG3	4:B:626:HOH:O	2.19	0.43
1:A:201:VAL:HG22	1:A:384:LEU:HG	2.00	0.43
1:A:346:TYR:O	1:A:347:SER:HB3	2.18	0.43
1:B:546:ASP:OD1	1:B:548:SER:HB2	2.19	0.43
1:A:82:LEU:HD13	1:A:249:ALA:HB2	2.01	0.42
1:B:80:ALA:HB3	1:B:245:LEU:HD23	2.00	0.42
1:B:83:LEU:HB2	1:B:173:MET:HA	2.01	0.42
1:B:215:MET:HB2	1:B:326:SER:HB2	2.01	0.42
1:A:284:VAL:O	1:A:287:THR:HG22	2.19	0.42
1:A:314:LEU:HB3	1:A:321:VAL:HG13	2.02	0.42
1:B:416:ALA:HB3	1:B:417:PRO:HD3	2.01	0.42
1:B:421:ALA:HA	1:B:425:LEU:HB2	2.01	0.41
1:A:52:VAL:HG22	1:A:226:SER:OG	2.21	0.41
1:A:459:LEU:N	1:A:460:PRO:CD	2.83	0.41
1:B:30:LEU:HD23	1:B:428:HIS:CD2	2.56	0.41
1:A:403:THR:HB	1:A:404:PRO:HD2	2.03	0.40
1:A:331:GLU:CD	1:A:331:GLU:H	2.28	0.40
1:B:141:LYS:HZ2	1:B:158:ARG:NH1	2.19	0.40
1:A:173:MET:CE	4:A:621:HOH:O	2.69	0.40
1:B:387:ASP:HA	1:B:388:PRO:HD3	1.99	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	561/581 (97%)	538 (96%)	21 (4%)	2 (0%)	30	50
1	B	547/581 (94%)	528 (96%)	17 (3%)	2 (0%)	30	50
All	All	1108/1162 (95%)	1066 (96%)	38 (3%)	4 (0%)	30	50

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	149	PRO
1	B	223	CYS
1	A	152	GLY
1	B	532	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	479/495 (97%)	449 (94%)	30 (6%)	16	30
1	B	470/495 (95%)	435 (93%)	35 (7%)	13	24
All	All	949/990 (96%)	884 (93%)	65 (7%)	14	27

All (65) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	LEU
1	A	52	VAL
1	A	56	ARG
1	A	57	LEU
1	A	68	LEU
1	A	83	LEU
1	A	86	GLU
1	A	96	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	106	LYS
1	A	124	GLU
1	A	126	LEU
1	A	130	THR
1	A	148	GLN
1	A	201	VAL
1	A	211	LYS
1	A	220	ASP
1	A	222	ARG
1	A	319	ASP
1	A	321	VAL
1	A	330	GLN
1	A	331	GLU
1	A	352	ASP
1	A	355	GLN
1	A	365	SER
1	A	425	LEU
1	A	432	ILE
1	A	440	GLU
1	A	469	LEU
1	A	517	ARG
1	A	556	SER
1	B	14	CYS
1	B	18	GLU
1	B	31	LEU
1	B	32	ARG
1	B	52	VAL
1	B	56	ARG
1	B	57	LEU
1	B	68	LEU
1	B	77	THR
1	B	86	GLU
1	B	100	LYS
1	B	126	LEU
1	B	130	THR
1	B	148	GLN
1	B	201	VAL
1	B	220	ASP
1	B	223	CYS
1	B	316	ASN
1	B	319	ASP
1	B	331	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	362	LEU
1	B	364	THR
1	B	425	LEU
1	B	434	LEU
1	B	441	LYS
1	B	459	LEU
1	B	465	ARG
1	B	469	LEU
1	B	483	ASN
1	B	487	SER
1	B	499	THR
1	B	508	ARG
1	B	517	ARG
1	B	547	LEU
1	B	556	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (23) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	ASN
1	A	28	ASN
1	A	35	ASN
1	A	49	GLN
1	A	206	ASN
1	A	273	ASN
1	A	316	ASN
1	A	330	GLN
1	A	355	GLN
1	A	438	GLN
1	A	446	GLN
1	A	483	ASN
1	A	514	GLN
1	A	562	HIS
1	B	34	HIS
1	B	35	ASN
1	B	49	GLN
1	B	95	HIS
1	B	206	ASN
1	B	316	ASN
1	B	446	GLN
1	B	461	GLN
1	B	527	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	XNZ	A	579	-	40,41,41	1.10	4 (10%)	58,61,61	1.78	6 (10%)
3	GOL	A	580	-	5,5,5	0.42	0	5,5,5	0.23	0
3	GOL	B	580	-	5,5,5	0.45	0	5,5,5	0.14	0
2	XNZ	B	579	-	40,41,41	1.03	4 (10%)	58,61,61	2.06	12 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XNZ	A	579	-	-	2/13/47/47	0/4/5/5
3	GOL	A	580	-	-	4/4/4/4	-
3	GOL	B	580	-	-	2/4/4/4	-
2	XNZ	B	579	-	-	1/13/47/47	0/4/5/5

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	579	XNZ	C15-C10	3.59	1.54	1.50
2	A	579	XNZ	C8-C9	3.44	1.54	1.51
2	A	579	XNZ	C15-C10	3.28	1.54	1.50
2	B	579	XNZ	C8-C9	2.63	1.54	1.51
2	B	579	XNZ	C30-N7	2.56	1.41	1.37
2	A	579	XNZ	C30-N7	2.27	1.41	1.37
2	A	579	XNZ	C12-C9	2.16	1.49	1.45
2	B	579	XNZ	C12-C9	2.09	1.49	1.45

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	579	XNZ	C31-C30-N7	7.08	127.64	117.87
2	B	579	XNZ	C4-N7-C30	-6.62	115.80	122.78
2	A	579	XNZ	C31-C30-N7	6.32	126.59	117.87
2	A	579	XNZ	C5-C4-N7	5.83	123.15	119.37
2	B	579	XNZ	C5-C4-N7	5.34	122.83	119.37
2	A	579	XNZ	O32-C30-N7	-4.78	115.50	120.71
2	A	579	XNZ	C4-N7-C30	-4.67	117.85	122.78
2	B	579	XNZ	O32-C30-N7	-4.34	115.98	120.71
2	B	579	XNZ	C14-C15-C10	3.53	117.26	113.40
2	B	579	XNZ	C9-C10-N11	-2.79	124.59	126.58
2	B	579	XNZ	C23-O22-C19	2.77	124.22	117.62
2	A	579	XNZ	C14-C15-C10	2.67	116.31	113.40
2	B	579	XNZ	C5-N11-C10	2.46	138.63	130.93
2	B	579	XNZ	O36-C12-C9	-2.46	118.79	121.32
2	A	579	XNZ	C14-C13-C12	-2.36	110.73	114.44
2	B	579	XNZ	C15-C10-N11	2.24	113.70	112.00
2	B	579	XNZ	C2-C3-C4	2.20	122.73	118.28
2	B	579	XNZ	C17-C16-C8	-2.19	117.54	121.72

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	579	XNZ	O32-C30-N7-C8
2	A	579	XNZ	C31-C30-N7-C8
3	A	580	GOL	O1-C1-C2-C3
3	A	580	GOL	C1-C2-C3-O3
3	B	580	GOL	C1-C2-C3-O3
3	A	580	GOL	O1-C1-C2-O2

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	580	GOL	O2-C2-C3-O3
3	B	580	GOL	O2-C2-C3-O3
2	B	579	XNZ	C24-C23-O22-C19

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	563/581 (96%)	-0.15	9 (1%) 70 69	7, 18, 34, 58	5 (0%)
1	B	553/581 (95%)	0.05	6 (1%) 78 77	6, 19, 33, 48	1 (0%)
All	All	1116/1162 (96%)	-0.05	15 (1%) 75 75	6, 19, 33, 58	6 (0%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	149	PRO	4.5
1	B	149	PRO	3.3
1	A	151	LYS	3.2
1	A	152	GLY	3.1
1	A	531	ARG	2.6
1	A	563	SER	2.6
1	B	563	SER	2.5
1	A	18	GLU	2.5
1	A	57	LEU	2.4
1	B	531	ARG	2.3
1	B	435	ALA	2.1
1	A	402	HIS	2.1
1	A	153	GLY	2.0
1	B	153	GLY	2.0
1	B	535	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	B	580	6/6	0.76	0.22	43,51,54,56	0
3	GOL	A	580	6/6	0.91	0.10	14,22,22,29	0
2	XNZ	B	579	37/37	0.96	0.07	14,22,32,34	0
2	XNZ	A	579	37/37	0.97	0.07	4,17,23,24	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around XNZ A 579:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.