



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 07:01 AM UTC

PDB ID : 6GN8 / pdb_00006gn8
Title : Exoenzyme S from *Pseudomonas aeruginosa* in complex with human 14-3-3 protein beta, trimeric crystal form
Authors : Karlberg, T.; Pinto, A.F.; Hornyak, P.; Thorsell, A.G.; Nareoja, K.; Schuler, H.
Deposited on : 2018-05-30
Resolution : 2.34 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3031 (2.36-2.32)
Clashscore	190562	3127 (2.36-2.32)
Ramachandran outliers	187476	3095 (2.36-2.32)
Sidechain outliers	187428	3095 (2.36-2.32)
RSRZ outliers	180081	3033 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	243	5% 83% 11% • 5%
1	B	243	19% 83% 8% • 7%
2	C	244	14% 74% 10% • 13%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5411 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 14-3-3 protein beta/alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	232	Total	C	N	O	S	0	0	0
			1871	1170	314	378	9			
1	B	225	Total	C	N	O	S	0	0	0
			1814	1137	306	362	9			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	235	GLU	-	expression tag	UNP P31946
A	236	ASN	-	expression tag	UNP P31946
A	237	LEU	-	expression tag	UNP P31946
A	238	TYR	-	expression tag	UNP P31946
A	239	PHE	-	expression tag	UNP P31946
A	240	GLN	-	expression tag	UNP P31946
A	241	SER	-	expression tag	UNP P31946
A	242	LEU	-	expression tag	UNP P31946
A	243	GLU	-	expression tag	UNP P31946
B	235	GLU	-	expression tag	UNP P31946
B	236	ASN	-	expression tag	UNP P31946
B	237	LEU	-	expression tag	UNP P31946
B	238	TYR	-	expression tag	UNP P31946
B	239	PHE	-	expression tag	UNP P31946
B	240	GLN	-	expression tag	UNP P31946
B	241	SER	-	expression tag	UNP P31946
B	242	LEU	-	expression tag	UNP P31946
B	243	GLU	-	expression tag	UNP P31946


- Molecule 2 is a protein called Exoenzyme S.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	212	Total	C	N	O	S	0	1	0
			1587	978	292	313	4			

There are 25 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	210	MET	-	initiating methionine	UNP Q93SQ1
C	211	GLY	-	expression tag	UNP Q93SQ1
C	212	SER	-	expression tag	UNP Q93SQ1
C	213	SER	-	expression tag	UNP Q93SQ1
C	214	HIS	-	expression tag	UNP Q93SQ1
C	215	HIS	-	expression tag	UNP Q93SQ1
C	216	HIS	-	expression tag	UNP Q93SQ1
C	217	HIS	-	expression tag	UNP Q93SQ1
C	218	HIS	-	expression tag	UNP Q93SQ1
C	219	HIS	-	expression tag	UNP Q93SQ1
C	220	SER	-	expression tag	UNP Q93SQ1
C	221	GLN	-	expression tag	UNP Q93SQ1
C	222	ASP	-	expression tag	UNP Q93SQ1
C	223	PRO	-	expression tag	UNP Q93SQ1
C	224	ASN	-	expression tag	UNP Q93SQ1
C	225	SER	-	expression tag	UNP Q93SQ1
C	226	GLU	-	expression tag	UNP Q93SQ1
C	227	ASN	-	expression tag	UNP Q93SQ1
C	228	LEU	-	expression tag	UNP Q93SQ1
C	229	TYR	-	expression tag	UNP Q93SQ1
C	230	PHE	-	expression tag	UNP Q93SQ1
C	231	GLN	-	expression tag	UNP Q93SQ1
C	232	GLY	-	expression tag	UNP Q93SQ1
C	379	ALA	GLU	engineered mutation	UNP Q93SQ1
C	381	ALA	GLU	engineered mutation	UNP Q93SQ1

- Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			8	6	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	59	Total	O	0	0
			59	59		
4	B	18	Total	O	0	0
			18	18		
4	C	54	Total	O	0	0
			54	54		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 14-3-3 protein beta/alpha



- Molecule 1: 14-3-3 protein beta/alpha



- Molecule 2: Exoenzyme S



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	159.91Å 59.40Å 120.33Å 90.00° 125.79° 90.00°	Depositor
Resolution (Å)	97.61 – 2.34 97.61 – 2.34	Depositor EDS
% Data completeness (in resolution range)	99.7 (97.61-2.34) 99.8 (97.61-2.34)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.76 (at 2.34Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.200 , 0.248 0.202 , 0.252	Depositor DCC
R_{free} test set	1958 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	47.3	Xtriage
Anisotropy	0.401	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 53.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5411	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.87	0/1898	1.36	8/2555 (0.3%)
1	B	0.82	0/1840	1.40	11/2474 (0.4%)
2	C	0.92	2/1607 (0.1%)	1.38	8/2155 (0.4%)
All	All	0.87	2/5345 (0.0%)	1.38	27/7184 (0.4%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	351	ALA	CA-C	8.06	1.57	1.52
2	C	301	MET	SD-CE	-7.25	1.61	1.79

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	427	ASP	CA-CB-CG	7.40	120.00	112.60
2	C	397	SER	CA-C-N	7.00	132.69	120.87
2	C	397	SER	C-N-CA	7.00	132.69	120.87
2	C	398	ASP	CA-CB-CG	6.96	119.56	112.60
1	B	215	ASP	CA-C-N	6.53	129.33	120.38
1	B	215	ASP	C-N-CA	6.53	129.33	120.38
1	B	114	PRO	CA-C-N	5.89	128.49	120.54
1	B	114	PRO	C-N-CA	5.89	128.49	120.54
1	B	99	VAL	N-CA-CB	5.71	119.14	110.58
1	A	99	VAL	N-CA-CB	5.45	118.76	110.58
1	A	4	ASP	CA-CB-CG	5.32	117.92	112.60
1	A	54	VAL	CA-C-N	5.32	126.00	119.99
1	A	54	VAL	C-N-CA	5.32	126.00	119.99
1	B	115	GLU	CA-C-N	5.25	127.74	120.29
1	B	115	GLU	C-N-CA	5.25	127.74	120.29
2	C	348	PRO	CA-C-N	5.19	125.70	119.94

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	348	PRO	C-N-CA	5.19	125.70	119.94
1	B	54	VAL	CA-C-N	5.17	125.83	119.99
1	B	54	VAL	C-N-CA	5.17	125.83	119.99
1	A	114	PRO	CA-C-N	5.16	127.51	120.54
1	A	114	PRO	C-N-CA	5.16	127.51	120.54
1	B	152	GLN	CA-C-N	5.11	127.12	120.28
1	B	152	GLN	C-N-CA	5.11	127.12	120.28
1	A	80	MET	CA-C-N	5.05	125.54	119.94
1	A	80	MET	C-N-CA	5.05	125.54	119.94
2	C	355	GLY	CA-C-N	5.02	127.84	120.71
2	C	355	GLY	C-N-CA	5.02	127.84	120.71

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1871	0	1848	10	0
1	B	1814	0	1798	11	0
2	C	1587	0	1577	18	0
3	C	8	0	14	0	0
4	A	59	0	0	0	0
4	B	18	0	0	0	0
4	C	54	0	0	0	0
All	All	5411	0	5237	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:447:MET:HA	2:C:448:ARG:HB3	1.34	1.08
2:C:297:ILE:HG22	2:C:301:MET:HE2	1.66	0.77

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:447:MET:HA	2:C:448:ARG:CB	2.15	0.77
2:C:447:MET:CA	2:C:448:ARG:HB3	2.16	0.73
2:C:447:MET:HB3	2:C:450:LEU:HB2	1.71	0.72
1:B:215:ASP:HA	2:C:446:ARG:HG3	1.71	0.70
1:A:115:GLU:HG2	1:A:168:ILE:HD12	1.90	0.53
2:C:448:ARG:HG2	2:C:449:GLY:H	1.75	0.51
2:C:447:MET:CB	2:C:450:LEU:HB2	2.39	0.50
1:B:221:GLN:HG2	1:B:224:ARG:HH12	1.78	0.48
2:C:347:ASN:HB3	2:C:350:VAL:HG22	1.96	0.48
2:C:356:GLN:HB3	2:C:359:ILE:HD11	1.95	0.48
1:A:118:VAL:HG21	1:A:162:MET:HE1	1.95	0.47
1:A:114:PRO:HG3	1:A:161:GLU:HG2	1.96	0.46
1:B:215:ASP:CA	2:C:446:ARG:HG3	2.44	0.46
2:C:245:PHE:O	2:C:248:ASP:HB2	2.16	0.46
1:A:74:ASN:HB3	1:A:77:LYS:HB2	1.98	0.46
1:B:74:ASN:HB3	1:B:77:LYS:HB2	1.99	0.45
1:A:103:LEU:HA	1:A:107:LEU:HB2	1.99	0.45
2:C:371:SER:HB3	2:C:380:LYS:HA	1.98	0.44
2:C:398:ASP:HB3	2:C:400:GLN:H	1.83	0.44
2:C:327:PHE:CE1	2:C:390:MET:HB2	2.52	0.44
1:B:89:GLU:HG2	1:B:134:VAL:HB	1.99	0.44
1:A:205:LEU:HD21	1:A:220:MET:HE1	2.00	0.44
1:B:103:LEU:HA	1:B:107:LEU:HB2	2.00	0.44
1:B:205:LEU:HD21	1:B:220:MET:HE1	1.99	0.43
2:C:280:LEU:CD2	2:C:301:MET:HE1	2.49	0.43
1:A:118:VAL:CG2	1:A:162:MET:HE1	2.48	0.43
1:A:89:GLU:HG2	1:A:134:VAL:HB	2.01	0.42
2:C:448:ARG:HG2	2:C:449:GLY:N	2.35	0.42
1:A:64:ILE:HD11	1:B:18:ALA:HB2	2.02	0.41
1:B:68:GLU:OE1	1:B:85:ARG:HD3	2.19	0.41
1:B:48:VAL:HG21	2:C:448:ARG:HG3	2.03	0.41
1:B:114:PRO:HG3	1:B:161:GLU:HG2	2.02	0.40
1:A:68:GLU:OE1	1:A:85:ARG:HD3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	230/243 (95%)	227 (99%)	3 (1%)	0	100	100
1	B	221/243 (91%)	215 (97%)	6 (3%)	0	100	100
2	C	209/244 (86%)	193 (92%)	12 (6%)	4 (2%)	6	4
All	All	660/730 (90%)	635 (96%)	21 (3%)	4 (1%)	21	22

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	259	GLY
2	C	447	MET
2	C	448	ARG
2	C	275	ILE

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	206/217 (95%)	195 (95%)	11 (5%)	20	25
1	B	199/217 (92%)	195 (98%)	4 (2%)	48	61
2	C	160/189 (85%)	152 (95%)	8 (5%)	22	27
All	All	565/623 (91%)	542 (96%)	23 (4%)	27	35

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	SER
1	A	78	GLN
1	A	92	LEU
1	A	93	GLN
1	A	99	VAL
1	A	138	ASP
1	A	156	GLU
1	A	161	GLU
1	A	193	LEU
1	A	210	GLU
1	A	211	GLU
1	B	99	VAL
1	B	161	GLU
1	B	162	MET
1	B	218	LEU
2	C	335	VAL
2	C	345	SER
2	C	352	ARG
2	C	356	GLN
2	C	405	ARG
2	C	446	ARG
2	C	447	MET
2	C	450	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	GLN
1	A	113	GLN
1	A	152	GLN
1	B	10	GLN
1	B	79	GLN
1	B	113	GLN
1	B	152	GLN
2	C	276	HIS
2	C	385	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MPD	C	501	-	7,7,7	0.57	0	9,10,10	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MPD	C	501	-	-	0/5/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	232/243 (95%)	0.27	12 (5%) 33 39	35, 52, 98, 134	0
1	B	225/243 (92%)	1.15	46 (20%) 3 3	46, 85, 163, 211	0
2	C	212/244 (86%)	0.69	35 (16%) 4 5	32, 58, 119, 162	1 (0%)
All	All	669/730 (91%)	0.70	93 (13%) 6 7	32, 62, 145, 211	1 (0%)

All (93) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	453	ALA	6.2
2	C	352	ARG	5.8
2	C	445	LEU	5.2
2	C	443	VAL	5.1
1	B	205	LEU	4.6
2	C	448	ARG	4.5
2	C	449	GLY	4.4
1	A	76	LYS	4.4
1	B	232	SER	4.4
1	B	229	LEU	4.4
1	B	112	THR	4.3
2	C	354	PHE	4.1
2	C	351	ALA	3.9
1	A	2	THR	3.9
1	B	202	ILE	3.8
2	C	350	VAL	3.8
2	C	446	ARG	3.8
2	C	355	GLY	3.8
2	C	432	PRO	3.6
1	B	230	TRP	3.3
1	B	212	SER	3.3
2	C	353	SER	3.3
2	C	415	LEU	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	259	GLY	3.1
1	B	216	SER	3.0
2	C	450	LEU	3.0
2	C	243	LYS	3.0
2	C	234	ASP	3.0
1	B	213	TYR	2.9
1	A	77	LYS	2.9
2	C	232	GLY	2.8
2	C	236	ALA	2.8
1	A	149	GLN	2.8
1	B	80	MET	2.8
1	B	76	LYS	2.7
1	B	160	LYS	2.7
1	A	3	MET	2.7
1	A	71	THR	2.7
1	B	228	THR	2.7
1	B	50	TYR	2.6
2	C	447	MET	2.6
1	B	117	LYS	2.6
1	B	121	LEU	2.6
1	B	218	LEU	2.6
2	C	452	LEU	2.6
1	A	73	ARG	2.5
1	B	3	MET	2.5
1	B	165	THR	2.5
2	C	237	LEU	2.5
1	B	214	LYS	2.5
1	B	187	PRO	2.5
1	B	113	GLN	2.5
1	B	153	GLU	2.5
1	B	4	ASP	2.5
1	B	217	THR	2.4
1	B	231	THR	2.4
1	B	201	ALA	2.4
1	A	72	GLU	2.4
1	A	233	GLU	2.4
2	C	413	GLY	2.4
2	C	444	ARG	2.4
1	B	161	GLU	2.4
1	B	108	ILE	2.4
2	C	255	ARG	2.4
1	B	203	ALA	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	258	GLY	2.3
1	A	145	SER	2.3
1	B	162	MET	2.3
1	B	206	ASP	2.3
2	C	257	PRO	2.3
1	B	114	PRO	2.2
2	C	348	PRO	2.2
2	C	233	ALA	2.2
2	C	431	LYS	2.2
1	B	168	ILE	2.2
1	B	137	GLY	2.2
1	B	111	ALA	2.2
1	B	181	TYR	2.2
1	B	115	GLU	2.2
1	B	30	ALA	2.2
1	B	222	LEU	2.2
1	B	195	LYS	2.2
2	C	256	GLN	2.1
1	B	104	ASP	2.1
1	B	116	SER	2.1
1	B	154	ALA	2.1
2	C	249	ALA	2.1
1	A	80	MET	2.1
1	B	105	LYS	2.0
1	B	186	SER	2.0
1	A	70	LYS	2.0
2	C	264	ALA	2.0
2	C	260	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MPD	C	501	8/8	0.93	0.14	61,62,67,70	0

6.5 Other polymers [i](#)

There are no such residues in this entry.