



# wwPDB X-ray Structure Validation Summary Report

Mar 5, 2026 – 02:57 PM UTC

PDB ID : 6GNF / pdb\_00006gnf  
Title : Granule Bound Starch Synthase from Cyanobacterium sp. CLg1 bound to acarbose and ADP  
Authors : Cuesta-Seijo, J.A.; Nielsen, M.M.; Palcic, M.M.  
Deposited on : 2018-05-30  
Resolution : 2.20 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	544	 5% 92% 5% • 10%
1	B	544	 10% 85% 5% • 10%
1	C	544	 7% 89% 5% • 6%
2	D	3	 33% 67%
2	E	3	 100%

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 12463 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycogen synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	522	4076	2630	667	752	27	0	2	0
1	B	492	3835	2476	632	700	27	0	1	0
1	C	514	3995	2577	660	731	27	0	0	0

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP V5SNJ5
A	-18	GLY	-	expression tag	UNP V5SNJ5
A	-17	SER	-	expression tag	UNP V5SNJ5
A	-16	SER	-	expression tag	UNP V5SNJ5
A	-15	HIS	-	expression tag	UNP V5SNJ5
A	-14	HIS	-	expression tag	UNP V5SNJ5
A	-13	HIS	-	expression tag	UNP V5SNJ5
A	-12	HIS	-	expression tag	UNP V5SNJ5
A	-11	HIS	-	expression tag	UNP V5SNJ5
A	-10	HIS	-	expression tag	UNP V5SNJ5
A	-9	SER	-	expression tag	UNP V5SNJ5
A	-8	SER	-	expression tag	UNP V5SNJ5
A	-7	GLY	-	expression tag	UNP V5SNJ5
A	-6	LEU	-	expression tag	UNP V5SNJ5
A	-5	VAL	-	expression tag	UNP V5SNJ5
A	-4	PRO	-	expression tag	UNP V5SNJ5
A	-3	ARG	-	expression tag	UNP V5SNJ5
A	-2	GLY	-	expression tag	UNP V5SNJ5
A	-1	SER	-	expression tag	UNP V5SNJ5
A	0	HIS	-	expression tag	UNP V5SNJ5
B	-19	MET	-	initiating methionine	UNP V5SNJ5
B	-18	GLY	-	expression tag	UNP V5SNJ5
B	-17	SER	-	expression tag	UNP V5SNJ5

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	SER	-	expression tag	UNP V5SNJ5
B	-15	HIS	-	expression tag	UNP V5SNJ5
B	-14	HIS	-	expression tag	UNP V5SNJ5
B	-13	HIS	-	expression tag	UNP V5SNJ5
B	-12	HIS	-	expression tag	UNP V5SNJ5
B	-11	HIS	-	expression tag	UNP V5SNJ5
B	-10	HIS	-	expression tag	UNP V5SNJ5
B	-9	SER	-	expression tag	UNP V5SNJ5
B	-8	SER	-	expression tag	UNP V5SNJ5
B	-7	GLY	-	expression tag	UNP V5SNJ5
B	-6	LEU	-	expression tag	UNP V5SNJ5
B	-5	VAL	-	expression tag	UNP V5SNJ5
B	-4	PRO	-	expression tag	UNP V5SNJ5
B	-3	ARG	-	expression tag	UNP V5SNJ5
B	-2	GLY	-	expression tag	UNP V5SNJ5
B	-1	SER	-	expression tag	UNP V5SNJ5
B	0	HIS	-	expression tag	UNP V5SNJ5
C	-19	MET	-	initiating methionine	UNP V5SNJ5
C	-18	GLY	-	expression tag	UNP V5SNJ5
C	-17	SER	-	expression tag	UNP V5SNJ5
C	-16	SER	-	expression tag	UNP V5SNJ5
C	-15	HIS	-	expression tag	UNP V5SNJ5
C	-14	HIS	-	expression tag	UNP V5SNJ5
C	-13	HIS	-	expression tag	UNP V5SNJ5
C	-12	HIS	-	expression tag	UNP V5SNJ5
C	-11	HIS	-	expression tag	UNP V5SNJ5
C	-10	HIS	-	expression tag	UNP V5SNJ5
C	-9	SER	-	expression tag	UNP V5SNJ5
C	-8	SER	-	expression tag	UNP V5SNJ5
C	-7	GLY	-	expression tag	UNP V5SNJ5
C	-6	LEU	-	expression tag	UNP V5SNJ5
C	-5	VAL	-	expression tag	UNP V5SNJ5
C	-4	PRO	-	expression tag	UNP V5SNJ5
C	-3	ARG	-	expression tag	UNP V5SNJ5
C	-2	GLY	-	expression tag	UNP V5SNJ5
C	-1	SER	-	expression tag	UNP V5SNJ5
C	0	HIS	-	expression tag	UNP V5SNJ5

- Molecule 2 is an oligosaccharide called 4,6-dideoxy-4- $\{[(1S,4R,5S,6S)-4,5,6\text{-trihydroxy-3-(hydroxymethyl)cyclohex-2-en-1-yl]amino}\}$ - $\alpha$ -D-glucopyranose-(1-4)- $\alpha$ -D-glucopyranose-(1-4)- $\beta$ -D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	3	Total	C	N	O	0	0	0
			44	25	1	18			
2	E	3	Total	C	N	O	0	0	0
			44	25	1	18			

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	C	1	Total	C	N	O	P	0	1
			54	20	10	20	4		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula:  $O_4S$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	C	1	Total O S 5 4 1	0	0
4	C	1	Total O S 5 4 1	0	0
4	C	1	Total O S 5 4 1	0	0

- Molecule 5 is alpha-D-glucopyranose (CCD ID: GLC) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 12 6 6	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	113	Total O 113 113	0	0
6	B	73	Total O 73 73	0	0
6	C	118	Total O 118 118	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glycogen synthase





- Molecule 1: Glycogen synthase



- Molecule 1: Glycogen synthase





- Molecule 2: 4,6-dideoxy-4- $\{[(1S,4R,5S,6S)-4,5,6\text{-trihydroxy-3-(hydroxymethyl)cyclohex-2-en-1-yl]amino}\}$ - $\alpha$ -D-glucopyranose-(1-4)- $\alpha$ -D-glucopyranose-(1-4)- $\beta$ -D-glucopyranose

Chain D:   33% 67%



- Molecule 2: 4,6-dideoxy-4- $\{[(1S,4R,5S,6S)-4,5,6\text{-trihydroxy-3-(hydroxymethyl)cyclohex-2-en-1-yl]amino}\}$ - $\alpha$ -D-glucopyranose-(1-4)- $\alpha$ -D-glucopyranose-(1-4)- $\beta$ -D-glucopyranose

Chain E:  100%



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	194.26Å 132.65Å 123.69Å 90.00° 126.30° 90.00°	Depositor
Resolution (Å)	48.56 – 2.20 48.56 – 2.20	Depositor EDS
% Data completeness (in resolution range)	95.6 (48.56-2.20) 95.6 (48.56-2.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.01 (at 2.20Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.205 , 0.224 0.209 , 0.224	Depositor DCC
$R_{free}$ test set	3647 reflections (3.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	60.8	Xtrriage
Anisotropy	0.051	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 49.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	12463	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	75.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.20% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, GLC, ADP, SO4, AC1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.49	0/4183	0.72	1/5682 (0.0%)
1	B	0.48	0/3929	0.73	1/5328 (0.0%)
1	C	0.51	0/4091	0.71	1/5548 (0.0%)
All	All	0.49	0/12203	0.72	3/16558 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	262	SER	N-CA-C	5.57	113.23	108.22
1	C	262	SER	N-CA-C	5.48	113.15	108.22
1	B	262	SER	N-CA-C	5.41	113.09	108.22

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4076	0	4101	9	0
1	B	3835	0	3855	15	0
1	C	3995	0	4022	12	0
2	D	44	0	30	0	0
2	E	44	0	30	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	27	0	12	0	0
3	B	27	0	12	0	0
3	C	54	0	24	3	0
4	A	20	0	0	0	0
4	B	10	0	0	1	0
4	C	15	0	0	1	0
5	B	12	0	12	0	0
6	A	113	0	0	2	0
6	B	73	0	0	1	0
6	C	118	0	0	0	0
All	All	12463	0	12098	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 37 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:SER:HB3	6:A:701:HOH:O	1.81	0.81
3:C:601[B]:ADP:O3B	3:C:601[B]:ADP:O2A	2.11	0.69
1:C:443:PRO:HB2	1:C:445:GLN:OE1	1.98	0.63
1:C:338:ARG:HG3	3:C:601[B]:ADP:O1A	1.98	0.63
1:B:17:THR:OG1	6:B:701:HOH:O	2.16	0.60

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	522/544 (96%)	506 (97%)	16 (3%)	0	100   100
1	B	487/544 (90%)	472 (97%)	15 (3%)	0	100   100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	510/544 (94%)	493 (97%)	17 (3%)	0	100	100
All	All	1519/1632 (93%)	1471 (97%)	48 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	449/466 (96%)	438 (98%)	11 (2%)	43	58
1	B	418/466 (90%)	408 (98%)	10 (2%)	43	58
1	C	437/466 (94%)	426 (98%)	11 (2%)	42	56
All	All	1304/1398 (93%)	1272 (98%)	32 (2%)	42	56

5 of 32 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	457	LYS
1	C	461	VAL
1	B	66	LYS
1	B	17	THR
1	C	492	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	293	GLN
1	C	291	ASN
1	B	166	GLN
1	C	166	GLN
1	B	153	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	GLC	B	602	-	12,12,12	0.76	0	17,17,17	1.76	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GLC	B	602	-	-	0/2/22/22	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	602	GLC	O5-C1-C2	3.97	117.28	110.30
5	B	602	GLC	C1-C2-C3	3.44	117.37	110.36
5	B	602	GLC	C1-O5-C5	2.97	119.39	113.65

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	BGC	D	1	2	12,12,12	0.54	0	17,17,17	1.02	1 (5%)
2	GLC	D	2	2	11,11,12	0.44	0	15,15,17	0.90	0
2	AC1	D	3	2	21,22,23	0.80	1 (4%)	22,32,34	1.25	3 (13%)
2	BGC	E	1	2	12,12,12	0.51	0	17,17,17	0.85	1 (5%)
2	GLC	E	2	2	11,11,12	0.52	0	15,15,17	1.08	1 (6%)
2	AC1	E	3	2	21,22,23	0.94	1 (4%)	22,32,34	1.63	5 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	D	1	2	-	2/2/22/22	0/1/1/1
2	GLC	D	2	2	-	0/2/19/22	0/1/1/1
2	AC1	D	3	2	-	1/6/43/46	0/2/2/2
2	BGC	E	1	2	-	0/2/22/22	0/1/1/1
2	GLC	E	2	2	-	2/2/19/22	0/1/1/1
2	AC1	E	3	2	-	1/6/43/46	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	3	AC1	C2B-C1B	2.48	1.56	1.53
2	E	3	AC1	C4A-C5B	-2.30	1.49	1.51

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	2	GLC	C1-O5-C5	3.66	117.09	112.19
2	D	3	AC1	O2B-C2B-C3B	-3.03	103.22	110.38
2	E	3	AC1	C2B-C3B-C4A	3.02	114.99	110.22
2	E	3	AC1	C7B-C1B-N4A	2.86	114.89	110.68
2	E	3	AC1	O3B-C3B-C4A	-2.77	104.15	109.64

There are no chirality outliers.


5 of 6 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	E	2	GLC	O5-C5-C6-O6
2	E	2	GLC	C4-C5-C6-O6
2	D	1	BGC	C4-C5-C6-O6
2	D	1	BGC	O5-C5-C6-O6
2	E	3	AC1	C7B-C1B-N4A-C4

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	SO4	A	605	-	4,4,4	0.45	0	6,6,6	0.27	0
4	SO4	C	604	-	4,4,4	0.51	0	6,6,6	0.22	0
3	ADP	C	601[A]	-	28,29,29	1.44	6 (21%)	43,45,45	1.84	10 (23%)
5	GLC	B	602	-	12,12,12	0.76	0	17,17,17	1.76	3 (17%)
3	ADP	B	601	-	28,29,29	1.50	6 (21%)	43,45,45	1.69	9 (20%)
4	SO4	C	603	-	4,4,4	0.54	0	6,6,6	0.31	0
4	SO4	A	606	-	4,4,4	0.45	0	6,6,6	0.19	0
4	SO4	C	605	-	4,4,4	0.41	0	6,6,6	0.15	0
3	ADP	A	601	-	28,29,29	1.48	5 (17%)	43,45,45	1.86	10 (23%)
3	ADP	C	601[B]	-	28,29,29	1.35	5 (17%)	43,45,45	1.89	10 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	SO4	A	603	-	4,4,4	0.49	0	6,6,6	0.24	0
4	SO4	A	604	-	4,4,4	0.45	0	6,6,6	0.17	0
4	SO4	B	603	-	4,4,4	0.49	0	6,6,6	0.36	0
4	SO4	B	604	-	4,4,4	0.45	0	6,6,6	0.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	C	601[A]	-	-	3/16/32/32	0/3/3/3
5	GLC	B	602	-	-	0/2/22/22	0/1/1/1
3	ADP	B	601	-	-	4/16/32/32	0/3/3/3
3	ADP	A	601	-	-	4/16/32/32	0/3/3/3
3	ADP	C	601[B]	-	-	5/16/32/32	0/3/3/3

The worst 5 of 22 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	601	ADP	C5-C4	4.93	1.47	1.39
3	A	601	ADP	C5-C4	4.72	1.47	1.39
3	C	601[A]	ADP	C5-C4	4.55	1.47	1.39
3	C	601[B]	ADP	C5-C4	4.50	1.47	1.39
3	A	601	ADP	PA-O3A	2.78	1.62	1.59

The worst 5 of 42 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	ADP	C5-C4-N3	-5.81	118.72	126.72
3	C	601[A]	ADP	C5-C4-N3	-5.42	119.25	126.72
3	C	601[B]	ADP	C5-C4-N3	-5.37	119.32	126.72
3	B	601	ADP	C5-C4-N3	-5.10	119.69	126.72
3	C	601[B]	ADP	N3-C4-N9	4.79	135.32	127.17

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	601	ADP	PA-O3A-PB-O2B
3	C	601[A]	ADP	PA-O3A-PB-O3B

*Continued on next page...*

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	C	601[B]	ADP	PA-O3A-PB-O3B
3	B	601	ADP	O4'-C4'-C5'-O5'
3	C	601[B]	ADP	O4'-C4'-C5'-O5'



There are no ring outliers.


3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	604	SO4	1	0
3	C	601[B]	ADP	3	0
4	B	603	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	522/544 (95%)	0.38	29 (5%) 30 27	40, 67, 109, 132	2 (0%)
1	B	492/544 (90%)	0.77	57 (11%) 9 7	43, 80, 143, 188	1 (0%)
1	C	514/544 (94%)	0.29	37 (7%) 21 18	40, 62, 116, 153	0
All	All	1528/1632 (93%)	0.48	123 (8%) 18 16	40, 70, 129, 188	3 (0%)

The worst 5 of 123 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	107	VAL	5.9
1	C	-8	SER	5.8
1	B	-5	VAL	5.4
1	B	521	ASN	5.3
1	C	213	PHE	5.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	GLC	B	602	12/12	0.97	0.06	47,58,59,63	0


### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	BGC	E	1	12/12	0.64	0.12	112,124,126,127	0
2	GLC	E	2	11/12	0.80	0.17	98,109,120,121	0
2	BGC	D	1	12/12	0.91	0.10	64,73,75,81	0
2	AC1	E	3	21/22	0.94	0.12	41,50,94,100	0
2	GLC	D	2	11/12	0.95	0.08	56,61,62,63	0
2	AC1	D	3	21/22	0.96	0.08	44,51,62,64	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	SO4	B	604	5/5	0.63	0.12	129,133,136,144	0
4	SO4	A	604	5/5	0.71	0.10	110,112,117,122	0
4	SO4	C	605	5/5	0.75	0.10	115,119,127,130	0
4	SO4	B	603	5/5	0.77	0.12	88,97,98,104	0
4	SO4	A	605	5/5	0.83	0.10	99,101,103,106	0
4	SO4	A	606	5/5	0.89	0.17	114,117,127,128	0
4	SO4	C	603	5/5	0.90	0.09	74,81,87,91	0
4	SO4	C	604	5/5	0.90	0.08	78,85,88,93	0
4	SO4	A	603	5/5	0.90	0.08	82,84,88,94	0
3	ADP	B	601	27/27	0.93	0.09	55,73,84,87	0
3	ADP	A	601	27/27	0.95	0.09	60,72,84,85	0
3	ADP	C	601[A]	27/27	0.95	0.10	55,74,75,76	27
3	ADP	C	601[B]	27/27	0.95	0.10	52,58,61,64	27
5	GLC	B	602	12/12	0.97	0.06	47,58,59,63	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around ADP A 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around ADP C 601 (A):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.