



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 07:11 PM UTC

PDB ID : 1GO4 / pdb_00001go4
Title : Crystal structure of Mad1-Mad2 reveals a conserved Mad2 binding motif in Mad1 and Cdc20.
Authors : Sironi, L.; Mapelli, M.; Jeang, K.T.; Musacchio, A.
Deposited on : 2001-10-17
Resolution : 2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	205	12% 59% 33% ...
1	B	205	9% 68% 22% 5%
1	C	205	3% 70% 21% 5%
1	D	205	64% 28% 61% 5% 6%
2	E	100	26% 49% 35% 13%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	100	24%57%27%.13%
2	G	100	40%49%43%7%. .
2	H	100	47%40%48%. .7%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9928 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitotic spindle assembly checkpoint protein MAD2A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	196	Total	C	N	O	S	0	0	0
			1578	1014	254	306	4			
1	B	195	Total	C	N	O	S	0	0	0
			1568	1008	253	303	4			
1	C	195	Total	C	N	O	S	0	0	0
			1568	1008	253	303	4			
1	D	193	Total	C	N	O	S	0	0	0
			1556	1002	250	300	4			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	133	ALA	ARG	engineered mutation	UNP Q13257
B	133	ALA	ARG	engineered mutation	UNP Q13257
C	133	ALA	ARG	engineered mutation	UNP Q13257
D	133	ALA	ARG	engineered mutation	UNP Q13257

- Molecule 2 is a protein called Mitotic spindle assembly checkpoint protein MAD1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	87	Total	C	N	O	S	0	0	0
			722	433	143	142	4			
2	F	87	Total	C	N	O	S	0	0	0
			722	433	143	142	4			
2	G	100	Total	C	N	O	S	0	0	0
			816	489	160	163	4			
2	H	93	Total	C	N	O	S	0	0	0
			770	464	150	152	4			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	88	Total 88	O 88	0	0
3	B	124	Total 124	O 124	0	0
3	C	225	Total 225	O 225	0	0
3	D	45	Total 45	O 45	0	0
3	E	56	Total 56	O 56	0	0
3	F	35	Total 35	O 35	0	0
3	G	24	Total 24	O 24	0	0
3	H	31	Total 31	O 31	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Mitotic spindle assembly checkpoint protein MAD2A




- Molecule 1: Mitotic spindle assembly checkpoint protein MAD2A




- Molecule 1: Mitotic spindle assembly checkpoint protein MAD2A




- Molecule 1: Mitotic spindle assembly checkpoint protein MAD2A




- Molecule 2: Mitotic spindle assembly checkpoint protein MAD1




- Molecule 2: Mitotic spindle assembly checkpoint protein MAD1



- Molecule 2: Mitotic spindle assembly checkpoint protein MAD1



- Molecule 2: Mitotic spindle assembly checkpoint protein MAD1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	111.04Å 63.02Å 139.51Å 90.00° 111.65° 90.00°	Depositor
Resolution (Å)	24.50 – 2.05 24.50 – 2.05	Depositor EDS
% Data completeness (in resolution range)	99.6 (24.50-2.05) 99.7 (24.50-2.05)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 2.04Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.240 , 0.268 0.238 , 0.266	Depositor DCC
R_{free} test set	5665 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	41.5	Xtriage
Anisotropy	0.081	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 51.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.011 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9928	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.90	3/1606 (0.2%)	1.17	15/2179 (0.7%)
1	B	0.80	1/1596 (0.1%)	1.14	9/2166 (0.4%)
1	C	1.03	1/1596 (0.1%)	1.27	18/2166 (0.8%)
1	D	0.54	1/1584 (0.1%)	1.38	14/2150 (0.7%)
2	E	0.82	4/726 (0.6%)	1.22	9/966 (0.9%)
2	F	0.56	0/726	0.85	0/966
2	G	0.60	0/821	1.34	9/1092 (0.8%)
2	H	0.48	0/775	1.17	7/1032 (0.7%)
All	All	0.78	10/9430 (0.1%)	1.22	81/12717 (0.6%)

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	10	GLY	N-CA	23.12	1.78	1.45
2	E	578	MET	CA-C	-7.31	1.43	1.52
1	D	194	ASN	N-CA	6.73	1.54	1.46
2	E	545	MET	SD-CE	6.68	1.96	1.79
1	A	9	GLN	CG-CD	-6.29	1.36	1.52

The worst 5 of 81 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	195	SER	N-CA-CB	-28.46	67.16	110.42
2	G	583	THR	N-CA-C	-20.69	88.28	114.56
1	D	59	LEU	N-CA-CB	-16.75	84.80	110.44
1	D	195	SER	N-CA-C	15.47	133.02	110.59
1	C	203	VAL	O-C-N	14.35	140.51	122.57

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1578	0	1596	74	0
1	B	1568	0	1588	48	0
1	C	1568	0	1588	40	0
1	D	1556	0	1579	206	0
2	E	722	0	720	58	0
2	F	722	0	722	42	0
2	G	816	0	811	66	0
2	H	770	0	766	83	0
3	A	88	0	0	13	0
3	B	124	0	0	18	0
3	C	225	0	0	18	0
3	D	45	0	0	71	0
3	E	56	0	0	13	0
3	F	35	0	0	4	0
3	G	24	0	0	8	0
3	H	31	0	0	24	0
All	All	9928	0	9370	572	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 31.

The worst 5 of 572 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:141:PHE:HE2	1:D:199:TYR:CE2	1.30	1.47
1:A:10:GLY:N	1:A:10:GLY:CA	1.78	1.46
2:E:529:ARG:CZ	3:E:2018:HOH:O	1.81	1.26
1:D:141:PHE:CE2	1:D:199:TYR:CE2	2.22	1.25
1:D:152:ASP:OD2	3:D:2030:HOH:O	1.57	1.16

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	194/205 (95%)	185 (95%)	8 (4%)	1 (0%)	24	16
1	B	193/205 (94%)	187 (97%)	5 (3%)	1 (0%)	24	16
1	C	193/205 (94%)	190 (98%)	3 (2%)	0	100	100
1	D	191/205 (93%)	167 (87%)	22 (12%)	2 (1%)	12	6
2	E	85/100 (85%)	82 (96%)	3 (4%)	0	100	100
2	F	85/100 (85%)	80 (94%)	5 (6%)	0	100	100
2	G	98/100 (98%)	95 (97%)	2 (2%)	1 (1%)	12	6
2	H	91/100 (91%)	84 (92%)	6 (7%)	1 (1%)	11	5
All	All	1130/1220 (93%)	1070 (95%)	54 (5%)	6 (0%)	24	16

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	10	GLY
1	D	41	GLU
1	D	114	SER
2	G	582	GLY
2	H	490	SER

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	181/189 (96%)	176 (97%)	5 (3%)	38	34

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	180/189 (95%)	172 (96%)	8 (4%)	25	19
1	C	180/189 (95%)	173 (96%)	7 (4%)	28	23
1	D	179/189 (95%)	176 (98%)	3 (2%)	53	53
2	E	78/88 (89%)	73 (94%)	5 (6%)	16	9
2	F	78/88 (89%)	74 (95%)	4 (5%)	21	14
2	G	88/88 (100%)	83 (94%)	5 (6%)	18	11
2	H	83/88 (94%)	80 (96%)	3 (4%)	31	26
All	All	1047/1108 (94%)	1007 (96%)	40 (4%)	29	24

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	496	GLU
2	G	578	MET
2	F	499	ASP
2	G	535	TYR
2	H	490	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	194	ASN
2	F	565	GLN
2	E	563	GLN
2	G	525	GLN
1	B	204	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	196/205 (95%)	0.83	24 (12%) 8 7	34, 53, 80, 95	0
1	B	195/205 (95%)	0.59	18 (9%) 14 14	29, 45, 69, 86	0
1	C	195/205 (95%)	-0.00	7 (3%) 46 46	21, 33, 61, 87	0
1	D	193/205 (94%)	2.67	131 (67%) 0 0	80, 98, 103, 105	0
2	E	87/100 (87%)	1.37	26 (29%) 1 0	27, 68, 94, 100	0
2	F	87/100 (87%)	1.13	24 (27%) 1 1	30, 69, 100, 104	0
2	G	100/100 (100%)	1.98	40 (40%) 1 0	49, 78, 92, 95	0
2	H	93/100 (93%)	2.08	47 (50%) 0 0	52, 82, 95, 101	0
All	All	1146/1220 (93%)	1.22	317 (27%) 1 1	21, 62, 100, 105	0

The worst 5 of 317 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	493	PHE	8.7
2	G	584	VAL	8.3
1	D	190	ILE	7.2
1	D	181	VAL	6.4
1	D	124	ILE	6.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.