



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 07:08 AM UTC

PDB ID : 4GO1 / pdb_00004go1
Title : Crystal Structure of full length transcription repressor LsrR from E. coli.
Authors : Wu, M.; Tao, Y.; Liu, X.; Zang, J.
Deposited on : 2012-08-17
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	325	 67% 20% 5% 9%
1	B	325	 2% 58% 26% 7% 9%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4452 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional regulator LsrR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	297	Total	C	N	O	S	0	0	0
			2221	1391	400	418	12			
1	B	296	Total	C	N	O	S	0	0	0
			2208	1387	396	413	12			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	318	LEU	-	expression tag	UNP P76141
A	319	GLU	-	expression tag	UNP P76141
A	320	HIS	-	expression tag	UNP P76141
A	321	HIS	-	expression tag	UNP P76141
A	322	HIS	-	expression tag	UNP P76141
A	323	HIS	-	expression tag	UNP P76141
A	324	HIS	-	expression tag	UNP P76141
A	325	HIS	-	expression tag	UNP P76141
B	318	LEU	-	expression tag	UNP P76141
B	319	GLU	-	expression tag	UNP P76141
B	320	HIS	-	expression tag	UNP P76141
B	321	HIS	-	expression tag	UNP P76141
B	322	HIS	-	expression tag	UNP P76141
B	323	HIS	-	expression tag	UNP P76141
B	324	HIS	-	expression tag	UNP P76141
B	325	HIS	-	expression tag	UNP P76141

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	9	Total	O	0	0
			9	9		
3	B	2	Total	O	0	0
			2	2		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Transcriptional regulator LsrR



• Molecule 1: Transcriptional regulator LsrR



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	79.70Å 113.87Å 185.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.00 50.00 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.5 (50.00-3.00) 99.5 (50.00-3.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.02 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.216 , 0.268 0.210 , 0.256	Depositor DCC
R_{free} test set	1245 reflections (7.20%)	wwPDB-VP
Wilson B-factor (Å ²)	53.1	Xtriage
Anisotropy	0.136	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 22.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4452	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/2246	0.96	3/3029 (0.1%)
1	B	0.66	1/2233 (0.0%)	0.98	6/3012 (0.2%)
All	All	0.65	1/4479 (0.0%)	0.97	9/6041 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	163	LEU	N-CA	7.20	1.55	1.46

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	162	GLN	CA-C-N	-9.26	107.24	121.86
1	B	162	GLN	C-N-CA	-9.26	107.24	121.86
1	B	163	LEU	N-CA-C	7.35	120.77	108.13
1	B	96	ALA	N-CA-C	6.58	120.63	112.47
1	A	276	ILE	CA-C-N	5.55	125.64	119.32
1	A	276	ILE	C-N-CA	5.55	125.64	119.32
1	A	92	GLY	N-CA-C	5.44	118.31	112.33
1	B	167	CYS	N-CA-C	5.15	117.12	109.25
1	B	161	GLY	O-C-N	-5.06	116.97	122.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2221	0	2273	40	0
1	B	2208	0	2269	72	0
2	A	6	0	8	0	0
2	B	6	0	8	0	0
3	A	9	0	0	2	0
3	B	2	0	0	0	0
All	All	4452	0	4558	109	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (109) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:162:GLN:O	1:B:163:LEU:HD23	1.32	1.28
1:B:134:ARG:HH11	1:B:134:ARG:HG3	1.19	1.05
1:B:162:GLN:O	1:B:163:LEU:CD2	2.15	0.95
1:B:160:ILE:CD1	1:B:169:VAL:HG21	1.96	0.94
1:B:160:ILE:HD13	1:B:169:VAL:HG21	1.55	0.87
1:A:231:GLU:O	1:A:235:ILE:HD12	1.76	0.85
1:B:109:MET:HE2	1:B:305:VAL:HG23	1.58	0.85
1:B:261:ILE:HD12	1:B:261:ILE:H	1.40	0.84
1:B:127:ALA:O	1:B:131:THR:HG22	1.78	0.83
1:B:129:MET:HE3	1:B:159:GLY:HA3	1.61	0.82
1:B:191:ASN:H	1:B:191:ASN:HD22	1.29	0.80
1:B:122:ILE:HG12	1:B:132:LEU:HD21	1.63	0.79
1:A:72:LEU:O	1:A:76:THR:HG22	1.85	0.76
1:B:109:MET:CE	1:B:305:VAL:HG23	2.15	0.75
1:B:160:ILE:HD11	1:B:169:VAL:HG21	1.68	0.73
1:A:32:THR:HB	1:A:35:GLU:HB2	1.70	0.73
1:B:134:ARG:HH11	1:B:134:ARG:CG	1.99	0.71
1:A:126:GLU:O	1:A:130:ASN:HB2	1.91	0.71
1:B:109:MET:HE2	1:B:305:VAL:CG2	2.20	0.70
1:A:283:ALA:O	1:A:306:THR:CG2	2.39	0.70
1:A:129:MET:HA	1:A:129:MET:HE3	1.75	0.69
1:A:283:ALA:O	1:A:306:THR:HG22	1.93	0.68
1:B:13:MET:HE2	1:B:18:GLN:HG2	1.75	0.68
1:A:11:GLN:O	1:A:11:GLN:HG3	1.92	0.68
1:B:134:ARG:HG3	1:B:134:ARG:NH1	1.99	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:261:ILE:H	1:B:261:ILE:CD1	2.08	0.66
1:B:34:SER:O	1:B:38:ASP:HB2	1.97	0.65
1:B:160:ILE:HD12	1:B:160:ILE:O	1.97	0.64
1:B:272:ALA:O	1:B:275:THR:HG22	1.95	0.64
1:A:135:LEU:HD22	1:A:138:PHE:HB3	1.80	0.64
1:B:156:TYR:O	1:B:160:ILE:HG22	1.98	0.63
1:A:129:MET:CE	1:A:159:GLY:HA3	2.29	0.62
1:A:33:GLN:H	1:A:33:GLN:HE21	1.46	0.62
1:A:137:GLY:O	1:A:141:SER:HB3	2.00	0.62
1:B:32:THR:HG23	1:B:35:GLU:H	1.64	0.62
1:B:191:ASN:HD22	1:B:191:ASN:N	1.95	0.62
1:B:122:ILE:HD11	1:B:132:LEU:HD11	1.82	0.61
1:B:107:ALA:O	1:B:111:MET:HG3	2.01	0.60
1:A:129:MET:HE2	1:A:159:GLY:HA3	1.84	0.60
1:B:142:GLN:HB2	1:B:144:ILE:CD1	2.32	0.59
1:B:292:ILE:HG22	1:B:296:MET:HE2	1.85	0.59
1:B:129:MET:HE2	1:B:156:TYR:O	2.03	0.58
1:A:101:ARG:NH1	1:B:29:ASP:OD2	2.37	0.58
1:A:100:GLY:O	1:A:104:ILE:HD12	2.03	0.58
1:A:34:SER:O	1:A:37:SER:HB3	2.03	0.58
1:B:160:ILE:O	1:B:160:ILE:CG1	2.54	0.56
1:A:21:ARG:NH2	1:B:75:GLU:OE2	2.38	0.56
1:B:191:ASN:H	1:B:191:ASN:ND2	1.98	0.56
1:B:172:ILE:HG13	1:B:196:VAL:HG11	1.88	0.56
1:B:257:THR:O	1:B:258:ASN:HB2	2.06	0.55
1:B:53:LYS:HA	1:B:56:GLN:HB2	1.89	0.55
1:B:52:GLU:O	1:B:56:GLN:HB2	2.06	0.55
1:A:64:ILE:HB	1:A:69:GLU:HG2	1.89	0.55
1:A:283:ALA:O	1:A:306:THR:HG23	2.06	0.54
1:A:91:PRO:O	1:A:101:ARG:NH2	2.40	0.54
1:A:11:GLN:N	1:A:12:GLY:HA2	2.23	0.54
1:A:205:VAL:HA	1:A:278:VAL:O	2.08	0.53
1:B:160:ILE:CD1	1:B:160:ILE:O	2.57	0.52
1:B:162:GLN:O	1:B:163:LEU:CG	2.57	0.52
1:B:160:ILE:HG12	1:B:169:VAL:HG11	1.92	0.52
1:A:33:GLN:H	1:A:33:GLN:NE2	2.08	0.51
1:B:261:ILE:HD12	1:B:261:ILE:N	2.19	0.51
1:A:66:SER:HB2	1:B:16:GLU:OE2	2.10	0.51
1:A:67:ARG:HG3	1:A:72:LEU:HD22	1.93	0.50
1:B:53:LYS:HA	1:B:56:GLN:CB	2.42	0.50
1:A:43:THR:HG23	1:A:46:LYS:HB2	1.93	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:13:MET:HE1	1:A:40:LEU:HD22	1.94	0.49
1:A:254:ASP:N	1:A:254:ASP:OD1	2.45	0.49
1:B:172:ILE:HG13	1:B:196:VAL:CG1	2.43	0.48
1:B:120:LEU:HG	1:B:122:ILE:HD12	1.96	0.48
1:B:117:GLN:O	1:B:145:ARG:NH2	2.42	0.48
1:B:134:ARG:CG	1:B:134:ARG:NH1	2.64	0.48
1:B:160:ILE:O	1:B:160:ILE:HG13	2.12	0.48
1:B:129:MET:CE	1:B:159:GLY:HA3	2.38	0.48
1:B:152:GLY:O	1:B:154:GLY:N	2.44	0.47
1:B:160:ILE:HD13	1:B:169:VAL:CG2	2.38	0.47
1:A:193:VAL:HG12	1:A:197:LEU:HD22	1.96	0.47
1:B:207:ILE:HD11	1:B:282:VAL:HG22	1.97	0.47
1:A:162:GLN:HE21	1:A:162:GLN:HA	1.81	0.45
1:A:288:LYS:NZ	3:A:509:HOH:O	2.49	0.45
1:B:91:PRO:O	1:B:101:ARG:NH1	2.32	0.45
1:A:76:THR:HA	1:A:79:ARG:HG3	1.98	0.45
1:B:56:GLN:O	1:B:57:SER:HB2	2.17	0.45
1:A:286:GLU:H	1:A:286:GLU:HG3	1.67	0.44
1:B:27:TYR:OH	1:B:55:HIS:HE1	2.00	0.44
1:B:90:ILE:HD11	1:B:102:LEU:HD13	2.00	0.43
1:B:128:THR:HA	1:B:131:THR:CG2	2.48	0.43
1:A:17:GLU:OE1	1:A:21:ARG:NH1	2.52	0.43
1:B:235:ILE:HG22	1:B:240:ALA:HB3	2.00	0.43
1:B:95:ASP:N	1:B:95:ASP:OD1	2.52	0.43
1:B:207:ILE:HD11	1:B:282:VAL:CG2	2.49	0.43
1:B:164:ASN:HD22	1:B:164:ASN:HA	1.65	0.43
1:B:201:GLN:HB3	1:B:268:LEU:HD11	2.00	0.43
1:B:191:ASN:N	1:B:191:ASN:ND2	2.62	0.43
1:B:279:ARG:NH1	1:B:299:GLY:O	2.51	0.42
1:A:135:LEU:CD2	1:A:138:PHE:HB3	2.49	0.42
1:B:98:VAL:O	1:B:102:LEU:HB2	2.19	0.42
1:B:243:ASP:HA	1:B:247:TYR:O	2.19	0.42
1:B:15:GLU:O	1:B:19:VAL:HG23	2.20	0.42
1:A:115:GLN:HB3	1:A:116:PRO:HD2	2.01	0.41
1:B:205:VAL:HG22	1:B:278:VAL:HG13	2.01	0.41
1:A:276:ILE:HA	1:A:277:PRO:HD3	1.92	0.41
1:A:292:ILE:O	1:A:295:ALA:HB3	2.21	0.41
1:B:32:THR:HG22	1:B:35:GLU:HG2	2.03	0.41
1:A:119:MET:HE3	3:A:502:HOH:O	2.20	0.41
1:A:129:MET:HE3	1:A:159:GLY:HA3	1.99	0.41
1:B:32:THR:O	1:B:35:GLU:HB2	2.19	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:119:MET:SD	1:B:147:VAL:HG22	2.61	0.41
1:B:159:GLY:C	1:B:161:GLY:H	2.30	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	293/325 (90%)	273 (93%)	20 (7%)	0	100	100
1	B	292/325 (90%)	268 (92%)	19 (6%)	5 (2%)	7	32
All	All	585/650 (90%)	541 (92%)	39 (7%)	5 (1%)	14	48

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	96	ALA
1	B	287	ASN
1	B	153	VAL
1	B	160	ILE
1	B	285	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/256 (90%)	190 (82%)	41 (18%)	2	10
1	B	229/256 (90%)	181 (79%)	48 (21%)	1	6
All	All	460/512 (90%)	371 (81%)	89 (19%)	1	8

All (89) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	GLU
1	A	31	LEU
1	A	33	GLN
1	A	43	THR
1	A	49	ARG
1	A	51	LEU
1	A	62	VAL
1	A	67	ARG
1	A	73	GLU
1	A	76	THR
1	A	78	LEU
1	A	87	VAL
1	A	88	ARG
1	A	95	ASP
1	A	102	LEU
1	A	128	THR
1	A	129	MET
1	A	130	ASN
1	A	132	LEU
1	A	135	LEU
1	A	136	SER
1	A	141	SER
1	A	158	THR
1	A	176	LEU
1	A	188	LYS
1	A	197	LEU
1	A	207	ILE
1	A	235	ILE
1	A	237	ARG
1	A	243	ASP
1	A	254	ASP
1	A	255	VAL
1	A	265	LEU
1	A	278	VAL
1	A	279	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	280	VAL
1	A	286	GLU
1	A	290	GLU
1	A	296	MET
1	A	306	THR
1	A	320	HIS
1	B	43	THR
1	B	51	LEU
1	B	61	ARG
1	B	63	GLN
1	B	67	ARG
1	B	72	LEU
1	B	73	GLU
1	B	78	LEU
1	B	87	VAL
1	B	88	ARG
1	B	95	ASP
1	B	113	LEU
1	B	122	ILE
1	B	126	GLU
1	B	128	THR
1	B	129	MET
1	B	131	THR
1	B	132	LEU
1	B	134	ARG
1	B	135	LEU
1	B	136	SER
1	B	140	SER
1	B	144	ILE
1	B	147	VAL
1	B	160	ILE
1	B	164	ASN
1	B	176	LEU
1	B	191	ASN
1	B	192	CYS
1	B	194	LYS
1	B	197	LEU
1	B	213	VAL
1	B	227	ILE
1	B	233	LEU
1	B	241	VAL
1	B	255	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	257	THR
1	B	261	ILE
1	B	265	LEU
1	B	275	THR
1	B	278	VAL
1	B	279	ARG
1	B	280	VAL
1	B	282	VAL
1	B	288	LYS
1	B	290	GLU
1	B	296	MET
1	B	318	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	GLN
1	A	33	GLN
1	A	85	GLN
1	A	86	HIS
1	A	162	GLN
1	A	287	ASN
1	B	55	HIS
1	B	130	ASN
1	B	162	GLN
1	B	164	ASN
1	B	191	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	B	401	-	5,5,5	0.36	0	5,5,5	0.29	0
2	GOL	A	401	-	5,5,5	0.41	0	5,5,5	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	401	-	-	2/4/4/4	-
2	GOL	A	401	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	401	GOL	C1-C2-C3-O3
2	B	401	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	297/325 (91%)	-0.21	1 (0%)	90 79	27, 43, 70, 81	4 (1%)
1	B	296/325 (91%)	-0.09	8 (2%)	56 33	26, 44, 79, 87	3 (1%)
All	All	593/650 (91%)	-0.15	9 (1%)	72 49	26, 44, 75, 87	7 (1%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	163	LEU	3.3
1	B	162	GLN	3.2
1	B	166	ALA	3.0
1	B	165	ALA	2.8
1	B	167	CYS	2.6
1	B	161	GLY	2.4
1	B	56	GLN	2.3
1	B	45	LEU	2.3
1	A	95	ASP	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	B	401	6/6	0.95	0.09	53,54,54,55	0
2	GOL	A	401	6/6	0.96	0.09	45,45,45,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.