



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 12:50 PM UTC

PDB ID : 5GON / pdb_00005gon
Title : Structures of a beta-lactam bridged analogue in complex with tubulin
Authors : Zhou, L.; Liu, Y.; Cheng, L.; Wang, Y.
Deposited on : 2016-07-28
Resolution : 2.48 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7589 (2.50-2.46)
Clashscore	190562	8295 (2.50-2.46)
Ramachandran outliers	187476	8164 (2.50-2.46)
Sidechain outliers	187428	8166 (2.50-2.46)
RSRZ outliers	180081	7593 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	440	81% 17% .
1	C	440	85% 14% .
2	B	431	3% 75% 21% ..
2	D	431	4% 67% 28% ..
3	E	136	4% 69% 18% .. 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	378	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	GOL	B	505	-	-	X	-

2 Entry composition [i](#)

There are 13 unique types of molecules in this entry. The entry contains 17847 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha-1B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	439	Total	C	N	O	S	0	13	0
			3488	2217	585	661	25			
1	C	440	Total	C	N	O	S	0	11	0
			3489	2209	588	668	24			

- Molecule 2 is a protein called Tubulin beta-2B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	422	Total	C	N	O	S	1	11	0
			3374	2126	568	653	27			
2	D	421	Total	C	N	O	S	0	4	0
			3326	2094	562	642	28			


- Molecule 3 is a protein called Stathmin-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	121	Total	C	N	O	S	0	3	0
			1016	628	183	199	6			

- Molecule 4 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	319	Total	C	N	O	S	0	5	0
			2633	1706	436	477	14			

- Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (CCD ID: GTP) (formula: C₁₀H₁₆N₅O₁₄P₃).



- | Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|-----------------|---------|---------|
| 6 | A | 1 | Total Mg
1 1 | 0 | 0 |
| 6 | B | 1 | Total Mg
1 1 | 0 | 0 |
| 6 | C | 1 | Total Mg
1 1 | 0 | 0 |
| 6 | D | 1 | Total Mg
1 1 | 0 | 0 |
| 6 | F | 1 | Total Mg
1 1 | 0 | 0 |

- 




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		
7	C	1	Total	C	O	0	0
			6	3	3		
7	C	1	Total	C	O	0	0
			6	3	3		
7	C	1	Total	C	O	0	0
			6	3	3		
7	C	1	Total	C	O	0	0
			6	3	3		
7	D	1	Total	C	O	0	0
			6	3	3		
7	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total Ca 1 1	0	0
8	B	1	Total Ca 1 1	0	0
8	C	1	Total Ca 1 1	0	0

- Molecule 9 is GUANOSINE-5'-DIPHOSPHATE (CCD ID: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	B	1	Total C N O P 28 10 5 11 2	0	0
9	D	1	Total C N O P 28 10 5 11 2	0	0

- Molecule 10 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
10	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 11 is (3R,4R)-4-(4-methoxy-3-oxidanyl-phenyl)-3-methyl-1-(3,4,5-trimethoxyphenyl)azetidin-2-one (CCD ID: 6ZR) (formula: $C_{20}H_{23}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
11	B	1	Total	C	N	O	0	0
			27	20	1	6		

- Molecule 12 is IMIDAZOLE (CCD ID: IMD) (formula: $C_3H_5N_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
12	C	1	Total	C	N	0	0
			5	3	2		
12	C	1	Total	C	N	0	0
			5	3	2		
12	C	1	Total	C	N	0	0
			5	3	2		


- Molecule 13 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	A	44	Total	O	0	0
			44	44		
13	B	52	Total	O	0	0
			52	52		
13	C	110	Total	O	0	0
			110	110		
13	D	25	Total	O	0	0
			25	25		
13	E	16	Total	O	0	0
			16	16		
13	F	20	Total	O	0	0
			20	20		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Tubulin alpha-1B chain




• Molecule 1: Tubulin alpha-1B chain




• Molecule 2: Tubulin beta-2B chain







• Molecule 2: Tubulin beta-2B chain




• Molecule 3: Stathmin-4



• Molecule 4: Uncharacterized protein





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	104.51Å 156.46Å 181.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	118.60 – 2.48 118.60 – 2.48	Depositor EDS
% Data completeness (in resolution range)	92.9 (118.60-2.48) 93.2 (118.60-2.48)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.21 (at 2.48Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.205 , 0.257 0.205 , 0.256	Depositor DCC
R_{free} test set	4920 reflections (4.62%)	wwPDB-VP
Wilson B-factor (Å ²)	36.4	Xtriage
Anisotropy	0.161	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	17847	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MES, GOL, CA, MG, IMD, 6ZR, GTP, GDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.01	2/3605 (0.1%)	1.05	10/4895 (0.2%)
1	C	1.17	3/3597 (0.1%)	1.08	11/4885 (0.2%)
2	B	1.09	2/3478 (0.1%)	1.04	7/4710 (0.1%)
2	D	1.02	3/3411 (0.1%)	1.07	10/4622 (0.2%)
3	E	1.05	0/1033	1.03	2/1370 (0.1%)
4	F	0.99	4/2704 (0.1%)	1.01	5/3650 (0.1%)
All	All	1.06	14/17828 (0.1%)	1.05	45/24132 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	1

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	360	PRO	C-N	15.73	1.54	1.33
2	D	42	LEU	C-N	15.57	1.56	1.33
2	B	172	MET	SD-CE	-8.82	1.57	1.79
4	F	196	HIS	CE1-NE2	7.02	1.39	1.32
4	F	196	HIS	CG-CD2	6.94	1.43	1.35

The worst 5 of 45 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	176	LYS	N-CA-C	-8.08	98.53	110.48
2	B	306	ASP	CA-C-N	8.04	127.76	119.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	306	ASP	C-N-CA	8.04	127.76	119.56
2	D	248	LEU	N-CA-C	-7.49	100.43	110.55
1	A	174	ALA	CA-C-N	6.93	127.22	119.32

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	249	ASN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3488	0	3441	45	0
1	C	3489	0	3413	45	0
2	B	3374	0	3270	78	0
2	D	3326	0	3221	81	0
3	E	1016	0	1041	15	0
4	F	2633	0	2631	84	0
5	A	32	0	12	0	0
5	C	32	0	12	1	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
6	F	1	0	0	0	0
7	A	18	0	24	1	0
7	B	18	0	24	12	0
7	C	24	0	32	6	0
7	D	12	0	16	3	0
8	A	1	0	0	0	0
8	B	1	0	0	0	0
8	C	1	0	0	0	0
9	B	28	0	12	0	0
9	D	28	0	12	0	0
10	B	12	0	13	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
11	B	27	0	0	3	0
12	C	15	0	15	4	0
13	A	44	0	0	2	0
13	B	52	0	0	0	0
13	C	110	0	0	3	0
13	D	25	0	0	1	0
13	E	16	0	0	1	0
13	F	20	0	0	0	0
All	All	17847	0	17189	344	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 344 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:F:240:LEU:HD23	4:F:245:ILE:HD12	1.30	1.13
2:B:320:ARG:HH11	7:B:505:GOL:H11	1.05	1.10
4:F:1:MET:HG2	4:F:28:LYS:HD2	1.19	1.09
2:B:320:ARG:NH1	7:B:505:GOL:H11	1.68	1.06
4:F:320:MET:HG2	4:F:330:ILE:HD11	1.34	1.05

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	450/440 (102%)	431 (96%)	19 (4%)	0	100	100
1	C	449/440 (102%)	437 (97%)	12 (3%)	0	100	100
2	B	429/431 (100%)	409 (95%)	18 (4%)	2 (0%)	24	40

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	D	421/431 (98%)	395 (94%)	18 (4%)	8 (2%)	6	10
3	E	120/136 (88%)	111 (92%)	8 (7%)	1 (1%)	16	28
4	F	307/378 (81%)	264 (86%)	34 (11%)	9 (3%)	3	5
All	All	2176/2256 (96%)	2047 (94%)	109 (5%)	20 (1%)	14	25

5 of 20 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	177	VAL
2	D	399	PHE
4	F	343	TYR
2	B	337	ASN
2	D	181	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	383/371 (103%)	365 (95%)	18 (5%)	23	44
1	C	382/371 (103%)	369 (97%)	13 (3%)	32	57
2	B	375/372 (101%)	351 (94%)	24 (6%)	16	31
2	D	368/372 (99%)	344 (94%)	24 (6%)	15	30
3	E	112/122 (92%)	98 (88%)	14 (12%)	4	8
4	F	293/336 (87%)	261 (89%)	32 (11%)	6	11
All	All	1913/1944 (98%)	1788 (94%)	125 (6%)	16	30

5 of 125 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	15	GLN
4	F	191	LEU
2	D	248	LEU
4	F	165	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	F	258	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	167	ASN
2	D	406	HIS
4	F	243	HIS
2	D	385	GLN
3	E	12	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 29 ligands modelled in this entry, 8 are monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	GOL	B	504	-	5,5,5	0.62	0	5,5,5	0.29	0
7	GOL	D	503	-	5,5,5	0.55	0	5,5,5	0.33	0
7	GOL	A	504	-	5,5,5	0.89	0	5,5,5	0.70	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	GOL	D	504	-	5,5,5	0.30	0	5,5,5	0.53	0
7	GOL	A	503	-	5,5,5	0.53	0	5,5,5	0.42	0
10	MES	B	507	-	12,12,12	2.24	1 (8%)	15,16,16	2.56	5 (33%)
5	GTP	C	503	6	33,34,34	1.42	5 (15%)	50,54,54	1.90	11 (22%)
11	6ZR	B	508	-	28,29,29	2.28	7 (25%)	38,42,42	2.52	14 (36%)
7	GOL	C	506	-	5,5,5	0.57	0	5,5,5	0.60	0
12	IMD	C	509	-	5,5,5	0.75	0	5,5,5	0.87	0
7	GOL	C	507	-	5,5,5	0.94	0	5,5,5	0.81	0
9	GDP	B	501	6	29,30,30	1.71	5 (17%)	45,47,47	1.88	9 (20%)
5	GTP	A	501	6	33,34,34	1.32	4 (12%)	50,54,54	1.79	11 (22%)
7	GOL	C	505	-	5,5,5	0.62	0	5,5,5	0.63	0
9	GDP	D	501	6	29,30,30	1.38	4 (13%)	45,47,47	1.89	10 (22%)
7	GOL	C	502	-	5,5,5	0.96	0	5,5,5	0.65	0
7	GOL	A	506	-	5,5,5	0.58	0	5,5,5	0.85	0
12	IMD	C	510	-	5,5,5	0.74	0	5,5,5	0.75	0
12	IMD	C	508	-	5,5,5	0.68	0	5,5,5	0.78	0
7	GOL	B	503	-	5,5,5	0.47	0	5,5,5	1.19	0
7	GOL	B	505	-	5,5,5	0.66	0	5,5,5	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	B	504	-	-	0/4/4/4	-
7	GOL	D	503	-	-	0/4/4/4	-
7	GOL	A	504	-	-	0/4/4/4	-
7	GOL	D	504	-	-	2/4/4/4	-
7	GOL	A	503	-	-	3/4/4/4	-
10	MES	B	507	-	-	0/6/14/14	0/1/1/1
5	GTP	C	503	6	-	8/22/38/38	0/3/3/3
11	6ZR	B	508	-	-	4/16/32/32	0/3/3/3
7	GOL	C	506	-	-	4/4/4/4	-
12	IMD	C	509	-	-	-	0/1/1/1
7	GOL	C	507	-	-	4/4/4/4	-
9	GDP	B	501	6	-	5/16/32/32	0/3/3/3
5	GTP	A	501	6	-	6/22/38/38	0/3/3/3
7	GOL	C	505	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	GDP	D	501	6	-	6/16/32/32	0/3/3/3
7	GOL	C	502	-	-	0/4/4/4	-
7	GOL	A	506	-	-	2/4/4/4	-
12	IMD	C	510	-	-	-	0/1/1/1
12	IMD	C	508	-	-	-	0/1/1/1
7	GOL	B	503	-	-	2/4/4/4	-
7	GOL	B	505	-	-	4/4/4/4	-

The worst 5 of 26 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	B	507	MES	C8-S	-7.37	1.67	1.77
11	B	508	6ZR	CAB-NAC	-6.18	1.32	1.43
11	B	508	6ZR	CAF-CAH	-5.46	1.50	1.57
9	B	501	GDP	PA-O3A	4.70	1.64	1.59
11	B	508	6ZR	CAI-CAH	-4.56	1.45	1.51

The worst 5 of 60 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	B	507	MES	O1S-S-C8	6.74	116.91	106.73
11	B	508	6ZR	OAN-CAM-CAK	6.65	124.52	114.55
5	C	503	GTP	C5-C4-N3	-6.31	118.34	128.39
9	B	501	GDP	C5-C4-N3	-6.30	118.36	128.39
5	C	503	GTP	C2-N3-C4	6.08	122.78	112.30

There are no chirality outliers.

5 of 52 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	501	GTP	PB-O3B-PG-O2G
5	A	501	GTP	C5'-O5'-PA-O3A
5	A	501	GTP	C5'-O5'-PA-O1A
5	C	503	GTP	C5'-O5'-PA-O3A
5	C	503	GTP	C5'-O5'-PA-O1A

There are no ring outliers.


13 monomers are involved in 29 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	504	GOL	1	0
7	D	504	GOL	3	0
10	B	507	MES	1	0
5	C	503	GTP	1	0
11	B	508	6ZR	3	0
7	C	506	GOL	1	0
12	C	509	IMD	2	0
7	C	507	GOL	2	0
7	C	505	GOL	3	0
12	C	510	IMD	1	0
12	C	508	IMD	1	0
7	B	503	GOL	2	0
7	B	505	GOL	10	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand 6ZR B 508



Ligand GDP B 501





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	439/440 (99%)	0.04	6 (1%) 73 71	17, 37, 62, 99	13 (2%)
1	C	440/440 (100%)	-0.50	3 (0%) 84 82	10, 22, 43, 59	11 (2%)
2	B	422/431 (97%)	-0.03	12 (2%) 55 51	10, 33, 65, 99	12 (2%)
2	D	421/431 (97%)	0.34	17 (4%) 42 38	21, 45, 73, 93	6 (1%)
3	E	121/136 (88%)	0.44	6 (4%) 34 30	12, 44, 76, 96	3 (2%)
4	F	319/378 (84%)	1.07	66 (20%) 2 2	24, 62, 143, 176	5 (1%)
All	All	2162/2256 (95%)	0.15	110 (5%) 33 30	10, 38, 79, 176	50 (2%)

The worst 5 of 110 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	F	149	ALA	6.9
4	F	173	ILE	5.9
4	F	102	PRO	5.7
4	F	245	ILE	5.6
4	F	231	ALA	5.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	GOL	D	503	6/6	0.67	0.18	74,81,86,87	0
7	GOL	A	506	6/6	0.75	0.16	52,63,69,73	0
6	MG	F	401	1/1	0.76	0.19	86,86,86,86	0
7	GOL	A	504	6/6	0.77	0.24	56,71,73,79	0
12	IMD	C	509	5/5	0.77	0.23	64,66,73,76	0
7	GOL	B	505	6/6	0.83	0.29	54,55,62,68	0
7	GOL	C	502	6/6	0.84	0.19	46,52,64,67	0
12	IMD	C	510	5/5	0.84	0.19	56,64,66,67	0
6	MG	A	502	1/1	0.86	0.16	37,37,37,37	0
7	GOL	B	503	6/6	0.86	0.20	42,49,52,72	0
7	GOL	B	504	6/6	0.86	0.19	53,62,69,71	0
6	MG	C	504	1/1	0.86	0.20	38,38,38,38	0
7	GOL	D	504	6/6	0.87	0.17	51,53,65,77	0
7	GOL	C	505	6/6	0.87	0.23	61,65,73,81	0
7	GOL	A	503	6/6	0.87	0.13	47,56,58,59	0
7	GOL	C	507	6/6	0.89	0.13	34,47,48,57	0
7	GOL	C	506	6/6	0.90	0.17	47,52,65,65	0
11	6ZR	B	508	27/27	0.90	0.12	35,47,53,58	0
6	MG	D	502	1/1	0.92	0.13	40,40,40,40	0
6	MG	B	502	1/1	0.93	0.13	16,16,16,16	0
8	CA	C	501	1/1	0.93	0.08	82,82,82,82	0
9	GDP	D	501	28/28	0.93	0.09	33,42,57,61	0
12	IMD	C	508	5/5	0.94	0.12	32,35,36,42	0
8	CA	B	506	1/1	0.95	0.08	81,81,81,81	0
10	MES	B	507	12/12	0.95	0.09	26,38,47,52	0
8	CA	A	505	1/1	0.97	0.04	59,59,59,59	0
5	GTP	A	501	32/32	0.98	0.05	18,23,30,33	0
5	GTP	C	503	32/32	0.98	0.05	12,16,20,22	0
9	GDP	B	501	28/28	0.99	0.05	13,18,21,24	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 6ZR B 508:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around GDP D 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GTP A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GTP C 503:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.