



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:37 PM UTC

PDB ID : 8GP6 / pdb_00008gp6
Title : Structure of the vaccinia virus A16/G9 sub-complex from the orthopoxvirus entry-fusion complex
Authors : Lu, G.W.; Yang, F.L.; Lin, S.
Deposited on : 2022-08-25
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	327	
2	A	348	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4462 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Myristoylated protein G9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	261	Total	C	N	O	S	0	0	0
			2112	1319	374	402	17			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	320	TRP	-	expression tag	UNP P07611
B	321	SER	-	expression tag	UNP P07611
B	322	HIS	-	expression tag	UNP P07611
B	323	PRO	-	expression tag	UNP P07611
B	324	GLN	-	expression tag	UNP P07611
B	325	PHE	-	expression tag	UNP P07611
B	326	GLU	-	expression tag	UNP P07611
B	327	LYS	-	expression tag	UNP P07611

- Molecule 2 is a protein called Virion membrane protein A16.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	285	Total	C	N	O	S	0	0	0
			2344	1485	403	433	23			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	343	HIS	-	expression tag	UNP P16710
A	344	HIS	-	expression tag	UNP P16710
A	345	HIS	-	expression tag	UNP P16710
A	346	HIS	-	expression tag	UNP P16710
A	347	HIS	-	expression tag	UNP P16710
A	348	HIS	-	expression tag	UNP P16710

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Myristoylated protein G9



• Molecule 2: Virion membrane protein A16



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α , β , γ	41.27Å 122.01Å 123.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.70 30.00 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.6 (30.00-2.70) 99.6 (30.00-2.70)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.16 (at 2.68Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.243 , 0.273 0.245 , 0.275	Depositor DCC
R_{free} test set	859 reflections (4.79%)	wwPDB-VP
Wilson B-factor (Å ²)	63.6	Xtriage
Anisotropy	0.061	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 34.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.016 for -h,l,k	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4462	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	B	0.83	0/2163	1.01	6/2927 (0.2%)
2	A	0.84	0/2412	1.05	9/3258 (0.3%)
All	All	0.84	0/4575	1.03	15/6185 (0.2%)

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	252	LYS	N-CA-C	-9.43	93.53	108.90
2	A	35	ARG	N-CA-C	7.87	121.03	109.07
1	B	13	ASP	CA-C-N	-7.10	114.47	119.66
1	B	13	ASP	C-N-CA	-7.10	114.47	119.66
2	A	255	SER	N-CA-C	6.95	120.40	108.02
1	B	138	ASP	N-CA-C	-6.84	104.67	113.16
2	A	273	GLU	N-CA-C	-6.84	96.23	110.80
2	A	271	THR	N-CA-C	-6.51	100.78	110.23
1	B	111	GLY	N-CA-C	-6.07	107.47	114.69
2	A	37	VAL	N-CA-C	5.93	116.72	108.89
2	A	120	ASP	N-CA-C	-5.87	104.51	112.26
2	A	136	LEU	N-CA-C	5.77	118.79	111.69
1	B	239	GLU	CA-C-N	-5.52	114.10	119.78
1	B	239	GLU	C-N-CA	-5.52	114.10	119.78
2	A	137	LEU	N-CA-C	5.27	117.07	110.91

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2112	0	2024	22	0
2	A	2344	0	2209	37	0
3	A	6	0	8	0	0
All	All	4462	0	4241	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:26:LEU:CB	2:A:35:ARG:HD2	2.14	0.77
1:B:94:ARG:NH2	1:B:106:ASP:OD1	2.18	0.74
2:A:26:LEU:HB2	2:A:35:ARG:HD2	1.70	0.73
1:B:18:VAL:O	2:A:71:SER:OG	2.04	0.72
2:A:92:SER:OG	2:A:99:SER:O	2.08	0.72
2:A:94:LYS:HG3	2:A:123:ALA:HB2	1.75	0.68
2:A:7:LEU:N	2:A:28:PHE:HA	2.11	0.65
2:A:169:ASP:OD2	2:A:193:ARG:NH2	2.35	0.59
2:A:267:LEU:HD22	2:A:269:GLU:HG2	1.85	0.58
2:A:119:THR:HG23	2:A:121:TYR:H	1.70	0.57
2:A:10:ILE:HG23	2:A:26:LEU:CD1	2.37	0.55
1:B:25:LEU:HD21	2:A:136:LEU:HB3	1.89	0.54
2:A:62:ILE:H	2:A:65:MET:HE3	1.74	0.53
1:B:67:ASN:OD1	1:B:67:ASN:C	2.52	0.52
1:B:241:ARG:HG2	1:B:245:ASP:HB2	1.92	0.52
1:B:10:PRO:HD3	2:A:10:ILE:HG21	1.92	0.51
1:B:40:LEU:HA	1:B:77:LEU:HD11	1.90	0.51
2:A:281:TYR:H	2:A:284:GLN:HE21	1.58	0.51
2:A:271:THR:O	2:A:272:ASP:C	2.51	0.51
2:A:35:ARG:HB3	2:A:111:TYR:OH	2.11	0.51
2:A:166:GLU:N	2:A:166:GLU:OE2	2.44	0.50
1:B:22:GLU:HA	1:B:25:LEU:HD22	1.93	0.50
1:B:267:CYS:HB2	2:A:266:TRP:CZ2	2.47	0.50
2:A:26:LEU:HB2	2:A:35:ARG:CD	2.40	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:147:GLN:O	1:B:151:SER:OG	2.31	0.49
1:B:85:ALA:HB3	1:B:86:PRO:HD3	1.95	0.48
2:A:10:ILE:HG23	2:A:26:LEU:HD12	1.95	0.47
2:A:62:ILE:HD11	2:A:92:SER:HB3	1.96	0.47
2:A:26:LEU:CB	2:A:35:ARG:CD	2.89	0.47
2:A:55:ILE:HD11	2:A:74:ILE:HG12	1.96	0.47
2:A:11:LYS:HD2	2:A:11:LYS:HA	1.64	0.47
1:B:150:GLY:HA2	1:B:192:HIS:CD2	2.50	0.47
1:B:12:ARG:HE	1:B:12:ARG:HB3	1.35	0.46
2:A:177:ARG:HA	2:A:203:ILE:HD11	1.97	0.46
1:B:99:LYS:HG2	1:B:104:ILE:HG13	1.98	0.46
1:B:267:CYS:CB	2:A:266:TRP:CZ2	2.99	0.46
2:A:194:LYS:HG2	2:A:195:PRO:HD3	1.98	0.46
1:B:159:THR:HG23	1:B:162:GLY:H	1.81	0.45
1:B:23:MET:SD	2:A:67:ILE:HG22	2.56	0.45
1:B:245:ASP:OD1	1:B:245:ASP:C	2.59	0.44
2:A:93:PHE:O	2:A:94:LYS:C	2.60	0.44
1:B:207:TYR:CD1	1:B:207:TYR:C	2.97	0.43
2:A:255:SER:HB2	2:A:258:LYS:HG3	2.02	0.42
2:A:115:LEU:O	2:A:119:THR:HG22	2.20	0.42
1:B:118:TRP:O	1:B:121:VAL:HB	2.20	0.42
2:A:24:MET:HG2	2:A:35:ARG:HH12	1.85	0.41
2:A:105:ASN:O	2:A:106:GLU:C	2.63	0.41
1:B:26:ASN:O	1:B:30:MET:HG3	2.20	0.41
2:A:38:LYS:HE3	2:A:104:GLN:HB2	2.02	0.41
1:B:164:GLN:O	1:B:165:SER:C	2.64	0.41
2:A:232:LEU:HD23	2:A:281:TYR:CD1	2.56	0.41
2:A:94:LYS:HD3	2:A:120:ASP:O	2.19	0.40
2:A:39:PHE:HA	2:A:100:LEU:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	259/327 (79%)	246 (95%)	13 (5%)	0	100	100
2	A	283/348 (81%)	266 (94%)	17 (6%)	0	100	100
All	All	542/675 (80%)	512 (94%)	30 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	236/297 (80%)	207 (88%)	29 (12%)	4	12
2	A	258/312 (83%)	211 (82%)	47 (18%)	2	5
All	All	494/609 (81%)	418 (85%)	76 (15%)	2	7

All (76) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	9	LEU
1	B	12	ARG
1	B	24	LEU
1	B	25	LEU
1	B	29	LYS
1	B	39	LEU
1	B	48	LEU
1	B	50	LYS
1	B	55	LYS
1	B	99	LYS
1	B	104	ILE
1	B	109	GLU
1	B	110	GLU
1	B	115	GLU
1	B	120	ASP
1	B	139	THR
1	B	151	SER
1	B	156	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	157	ASN
1	B	158	ARG
1	B	159	THR
1	B	160	VAL
1	B	174	GLN
1	B	199	GLU
1	B	210	ARG
1	B	218	GLN
1	B	228	ARG
1	B	230	LYS
1	B	236	LYS
2	A	7	LEU
2	A	9	ARG
2	A	10	ILE
2	A	11	LYS
2	A	12	ILE
2	A	16	ILE
2	A	20	ARG
2	A	24	MET
2	A	25	GLU
2	A	26	LEU
2	A	32	GLU
2	A	34	ASN
2	A	35	ARG
2	A	38	LYS
2	A	44	TYR
2	A	45	THR
2	A	51	SER
2	A	55	ILE
2	A	68	ASP
2	A	75	VAL
2	A	81	GLN
2	A	85	ILE
2	A	92	SER
2	A	108	THR
2	A	113	LYS
2	A	116	LYS
2	A	131	ILE
2	A	132	LYS
2	A	136	LEU
2	A	137	LEU
2	A	139	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	A	142	SER
2	A	152	THR
2	A	153	LYS
2	A	173	THR
2	A	194	LYS
2	A	226	THR
2	A	230	THR
2	A	237	ASN
2	A	252	LYS
2	A	253	SER
2	A	258	LYS
2	A	267	LEU
2	A	269	GLU
2	A	287	GLN
2	A	289	THR
2	A	290	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	44	HIS
1	B	79	GLN
1	B	93	ASN
1	B	146	HIS
1	B	157	ASN
1	B	171	ASN
1	B	250	ASN
1	B	254	ASN
2	A	69	HIS
2	A	126	GLN
2	A	151	ASN
2	A	167	HIS
2	A	254	ASN
2	A	283	ASN
2	A	284	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	401	-	5,5,5	0.32	0	5,5,5	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	401	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	401	GOL	C1-C2-C3-O3
3	A	401	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	261/327 (79%)	0.07	0 100 100	43, 61, 92, 116	0
2	A	285/348 (81%)	0.31	15 (5%) 32 28	43, 65, 109, 145	0
All	All	546/675 (80%)	0.20	15 (2%) 56 53	43, 63, 102, 145	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	35	ARG	4.3
2	A	26	LEU	3.4
2	A	10	ILE	2.9
2	A	194	LYS	2.7
2	A	144	ALA	2.6
2	A	44	TYR	2.5
2	A	43	SER	2.4
2	A	45	THR	2.4
2	A	7	LEU	2.3
2	A	280	LEU	2.3
2	A	31	PRO	2.3
2	A	42	GLU	2.2
2	A	276	ASP	2.1
2	A	266	TRP	2.1
2	A	127	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	A	401	6/6	0.74	0.23	66,68,71,76	0

6.5 Other polymers

There are no such residues in this entry.