



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2026 – 06:34 PM UTC

PDB ID : 2GQD / pdb_00002gqd
Title : The crystal structure of B-ketoacyl-ACP synthase II (FabF) from Staphylococcus aureus
Authors : Miller, D.J.; White, S.W.; Zhang, Y.M.; Rock, C.O.
Deposited on : 2006-04-20
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	437	 76% 17% • 6%
1	B	437	 76% 17% • 6%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6358 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-oxoacyl-[acyl-carrier-protein] synthase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	411	Total	C	N	O	S	0	0	0
			3044	1894	528	606	16			
1	B	411	Total	C	N	O	S	0	0	0
			3041	1893	527	605	16			

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-22	MET	-	expression tag	UNP Q8NXE1
A	-21	GLY	-	expression tag	UNP Q8NXE1
A	-20	SER	-	expression tag	UNP Q8NXE1
A	-19	SER	-	expression tag	UNP Q8NXE1
A	-18	HIS	-	expression tag	UNP Q8NXE1
A	-17	HIS	-	expression tag	UNP Q8NXE1
A	-16	HIS	-	expression tag	UNP Q8NXE1
A	-15	HIS	-	expression tag	UNP Q8NXE1
A	-14	HIS	-	expression tag	UNP Q8NXE1
A	-13	HIS	-	expression tag	UNP Q8NXE1
A	-12	SER	-	expression tag	UNP Q8NXE1
A	-11	SER	-	expression tag	UNP Q8NXE1
A	-10	GLY	-	expression tag	UNP Q8NXE1
A	-9	LEU	-	expression tag	UNP Q8NXE1
A	-8	VAL	-	expression tag	UNP Q8NXE1
A	-7	PRO	-	expression tag	UNP Q8NXE1
A	-6	ARG	-	expression tag	UNP Q8NXE1
A	-5	GLY	-	expression tag	UNP Q8NXE1
A	-4	SER	-	expression tag	UNP Q8NXE1
A	-3	HIS	-	expression tag	UNP Q8NXE1
A	-2	MET	-	expression tag	UNP Q8NXE1
A	-1	LEU	-	expression tag	UNP Q8NXE1
A	0	GLU	-	expression tag	UNP Q8NXE1
B	-22	MET	-	expression tag	UNP Q8NXE1
B	-21	GLY	-	expression tag	UNP Q8NXE1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-20	SER	-	expression tag	UNP Q8NXE1
B	-19	SER	-	expression tag	UNP Q8NXE1
B	-18	HIS	-	expression tag	UNP Q8NXE1
B	-17	HIS	-	expression tag	UNP Q8NXE1
B	-16	HIS	-	expression tag	UNP Q8NXE1
B	-15	HIS	-	expression tag	UNP Q8NXE1
B	-14	HIS	-	expression tag	UNP Q8NXE1
B	-13	HIS	-	expression tag	UNP Q8NXE1
B	-12	SER	-	expression tag	UNP Q8NXE1
B	-11	SER	-	expression tag	UNP Q8NXE1
B	-10	GLY	-	expression tag	UNP Q8NXE1
B	-9	LEU	-	expression tag	UNP Q8NXE1
B	-8	VAL	-	expression tag	UNP Q8NXE1
B	-7	PRO	-	expression tag	UNP Q8NXE1
B	-6	ARG	-	expression tag	UNP Q8NXE1
B	-5	GLY	-	expression tag	UNP Q8NXE1
B	-4	SER	-	expression tag	UNP Q8NXE1
B	-3	HIS	-	expression tag	UNP Q8NXE1
B	-2	MET	-	expression tag	UNP Q8NXE1
B	-1	LEU	-	expression tag	UNP Q8NXE1
B	0	GLU	-	expression tag	UNP Q8NXE1

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	129	Total	O	0	0
			129	129		
2	B	144	Total	O	0	0
			144	144		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3-oxoacyl-[acyl-carrier-protein] synthase 2



- Molecule 1: 3-oxoacyl-[acyl-carrier-protein] synthase 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	78.40Å 78.40Å 231.27Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.91 – 2.30 29.91 – 2.30	Depositor EDS
% Data completeness (in resolution range)	96.7 (29.91-2.30) 96.7 (29.91-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.54 (at 2.31Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.221 , 0.273 0.219 , 0.272	Depositor DCC
R_{free} test set	1723 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å ²)	50.5	Xtriage
Anisotropy	0.045	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.479 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	6358	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/3095	0.84	5/4189 (0.1%)
1	B	0.42	1/3092 (0.0%)	0.81	2/4185 (0.0%)
All	All	0.42	1/6187 (0.0%)	0.83	7/8374 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	414	ALA	C-O	5.30	1.34	1.23

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	140	ILE	CA-C-N	6.55	127.07	119.47
1	A	140	ILE	C-N-CA	6.55	127.07	119.47
1	B	140	ILE	CA-C-N	6.15	126.04	119.28
1	B	140	ILE	C-N-CA	6.15	126.04	119.28
1	A	381	VAL	CA-C-N	6.12	126.06	119.76
1	A	381	VAL	C-N-CA	6.12	126.06	119.76
1	A	140	ILE	N-CA-C	5.43	114.07	109.02

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3044	0	2996	56	0
1	B	3041	0	2992	51	0
2	A	129	0	0	3	0
2	B	144	0	0	3	0
All	All	6358	0	5988	98	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (98) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:41:ARG:HH21	1:B:74:ARG:HD3	1.33	0.92
1:B:41:ARG:NH2	1:B:74:ARG:HD3	1.87	0.90
1:A:129:ARG:HH11	1:A:129:ARG:HG3	1.39	0.87
1:A:129:ARG:HH11	1:A:129:ARG:CG	1.94	0.80
1:A:41:ARG:HH11	1:A:41:ARG:HG3	1.47	0.78
1:A:140:ILE:HG13	1:A:143:MET:HG2	1.67	0.75
1:B:140:ILE:HG13	1:B:143:MET:HG2	1.67	0.75
1:B:66:LYS:O	1:B:70:ARG:HG2	1.88	0.74
1:B:110:ILE:HG22	1:B:113:MET:HG2	1.72	0.72
1:A:42:ILE:HD11	1:A:201:ILE:HD12	1.72	0.70
1:B:413:GLU:HG2	2:B:542:HOH:O	1.93	0.69
1:B:44:THR:HG23	1:B:50:HIS:HD2	1.58	0.68
1:B:370:VAL:HG12	1:B:371:THR:HG23	1.76	0.68
1:B:41:ARG:NH2	1:B:234:GLU:OE1	2.28	0.66
1:A:199:MET:HG2	1:B:116:PHE:CE2	2.29	0.66
1:B:42:ILE:HD11	1:B:201:ILE:HD12	1.76	0.66
1:A:229:GLY:HA2	1:A:308:THR:HG22	1.79	0.63
1:B:181:ARG:HD2	1:B:183:ASP:OD2	1.99	0.63
1:B:229:GLY:HA2	1:B:308:THR:HG22	1.81	0.62
1:A:199:MET:HG2	1:B:116:PHE:HE2	1.63	0.62
1:A:120:HIS:HD2	1:B:199:MET:HE3	1.65	0.62
1:B:6:ARG:HH22	1:B:413:GLU:HG3	1.65	0.61
1:A:373:ASP:HB2	2:A:514:HOH:O	2.00	0.61
1:A:370:VAL:HG12	1:A:371:THR:HG23	1.83	0.60
1:B:110:ILE:CG2	1:B:113:MET:HG2	2.32	0.59
1:A:137:PRO:O	1:A:143:MET:HG3	2.02	0.59
1:A:282:ARG:NH2	1:B:155:LYS:O	2.37	0.57
1:A:330:LEU:HD13	1:A:332:VAL:HG23	1.87	0.57
1:A:282:ARG:NH2	1:B:155:LYS:HB3	2.21	0.56
1:A:30:LEU:HD13	1:A:362:LYS:HD3	1.87	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:54:GLU:HG2	1:A:196:ILE:HD13	1.89	0.55
1:A:120:HIS:CD2	1:B:199:MET:HE3	2.41	0.55
1:B:61:GLU:OE2	1:B:66:LYS:HE3	2.07	0.55
1:A:129:ARG:CG	1:A:129:ARG:NH1	2.63	0.54
1:A:30:LEU:CD1	1:A:362:LYS:HD3	2.39	0.53
1:B:41:ARG:HH21	1:B:74:ARG:CD	2.15	0.53
1:B:161:THR:O	1:B:167:THR:HG23	2.09	0.53
1:A:413:GLU:O	1:A:414:ALA:CB	2.58	0.52
1:A:41:ARG:HG3	1:A:41:ARG:NH1	2.22	0.52
1:B:161:THR:HG21	1:B:170:ASN:HB3	1.92	0.51
1:A:129:ARG:HG3	1:A:129:ARG:NH1	2.18	0.51
1:B:382:PRO:O	1:B:384:GLU:N	2.38	0.51
1:A:161:THR:O	1:A:167:THR:HG23	2.10	0.51
1:A:413:GLU:O	1:A:414:ALA:HB2	2.11	0.51
1:B:137:PRO:O	1:B:143:MET:HG3	2.12	0.49
1:B:6:ARG:NH2	1:B:413:GLU:HG3	2.27	0.49
1:A:288:MET:HG3	1:A:394:MET:HE3	1.94	0.49
1:B:204:PHE:HB3	1:B:210:LEU:HG	1.94	0.49
1:A:48:SER:HB2	1:A:212:THR:HG23	1.95	0.49
1:B:41:ARG:NH2	1:B:78:TYR:OH	2.42	0.49
1:B:60:ILE:HD11	1:B:64:ILE:HD12	1.95	0.48
1:A:278:GLU:OE2	1:A:282:ARG:HD3	2.13	0.48
1:A:140:ILE:HD11	1:A:143:MET:HE3	1.96	0.48
1:B:17:PRO:O	1:B:56:LYS:HE2	2.14	0.47
1:B:101:ARG:HD3	1:B:185:ASP:OD2	2.15	0.47
1:A:274:ALA:O	1:A:314:ASN:ND2	2.48	0.47
1:B:338:MET:HE2	1:B:365:PRO:O	2.14	0.47
1:B:74:ARG:NH1	1:B:114:GLU:OE1	2.48	0.47
1:B:288:MET:HG3	1:B:394:MET:HE3	1.96	0.46
1:A:204:PHE:HB3	1:A:210:LEU:HG	1.97	0.46
1:A:17:PRO:HG2	1:A:339:THR:O	2.14	0.46
1:A:140:ILE:CG1	1:A:143:MET:HG2	2.43	0.46
1:B:288:MET:HG3	1:B:394:MET:CE	2.45	0.46
1:A:381:VAL:N	1:A:382:PRO:HD3	2.31	0.45
1:A:302:ASN:ND2	1:A:350:GLU:HG2	2.30	0.45
1:A:306:THR:O	1:A:307:SER:HB3	2.17	0.45
1:B:44:THR:HG21	2:B:477:HOH:O	2.17	0.45
1:A:225:GLU:HG3	2:A:514:HOH:O	2.17	0.44
1:A:162:VAL:O	1:B:142:ASP:HB3	2.18	0.44
1:A:302:ASN:HD21	1:A:350:GLU:HG2	1.82	0.44
1:B:381:VAL:N	1:B:382:PRO:HD3	2.33	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:142:ASP:HB3	1:B:162:VAL:O	2.18	0.44
1:A:41:ARG:HH11	1:A:41:ARG:CG	2.23	0.43
1:B:165:CYS:HB2	1:B:399:GLY:HA3	1.99	0.43
1:A:382:PRO:O	1:A:384:GLU:N	2.37	0.43
1:A:161:THR:HG21	1:A:170:ASN:HB3	1.99	0.43
1:B:54:GLU:HG2	1:B:196:ILE:HD13	2.00	0.43
1:A:166:ALA:HA	1:A:397:SER:HB3	2.00	0.43
1:B:306:THR:O	1:B:307:SER:HB3	2.19	0.43
1:A:225:GLU:N	2:A:514:HOH:O	2.53	0.42
1:B:110:ILE:CG2	1:B:113:MET:CG	2.97	0.42
1:B:166:ALA:HA	1:B:397:SER:HB3	2.01	0.42
1:A:220:CYS:O	1:A:368:HIS:HD2	2.03	0.42
1:A:41:ARG:NH2	1:A:234:GLU:OE2	2.49	0.42
1:A:195:PRO:O	1:A:200:ALA:HB3	2.21	0.41
1:A:17:PRO:O	1:A:56:LYS:HE2	2.21	0.41
1:A:47:TYR:HE2	1:A:202:ALA:HB2	1.86	0.41
1:A:272:ALA:HA	1:A:273:PRO:HD3	1.91	0.41
1:B:342:LEU:HB3	1:B:345:ALA:HB3	2.02	0.41
1:A:123:LEU:HB3	1:B:199:MET:HE1	2.02	0.41
1:A:373:ASP:HA	1:A:374:PRO:HD3	1.95	0.41
1:A:222:PRO:O	1:A:224:GLN:HG2	2.20	0.40
1:A:110:ILE:CG2	1:A:113:MET:HG2	2.51	0.40
1:B:44:THR:CG2	2:B:477:HOH:O	2.69	0.40
1:A:41:ARG:NH1	1:A:41:ARG:CG	2.83	0.40
1:B:33:VAL:O	1:B:338:MET:HB3	2.21	0.40
1:B:242:GLU:HG2	1:B:246:SER:HB3	2.04	0.40
1:B:367:ILE:O	1:B:368:HIS:HB2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	409/437 (94%)	387 (95%)	20 (5%)	2 (0%)	24	31
1	B	409/437 (94%)	383 (94%)	23 (6%)	3 (1%)	18	23
All	All	818/874 (94%)	770 (94%)	43 (5%)	5 (1%)	21	27

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	383	ASN
1	B	383	ASN
1	A	305	GLY
1	B	305	GLY
1	B	216	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	316/339 (93%)	299 (95%)	17 (5%)	20	29
1	B	315/339 (93%)	302 (96%)	13 (4%)	27	41
All	All	631/678 (93%)	601 (95%)	30 (5%)	23	35

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	ASN
1	A	5	LYS
1	A	37	ASP
1	A	125	ASP
1	A	129	ARG
1	A	145	THR
1	A	174	GLU
1	A	215	ASP
1	A	239	LEU
1	A	245	GLU
1	A	310	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	328	LYS
1	A	330	LEU
1	A	342	LEU
1	A	349	ILE
1	A	380	ILE
1	A	384	GLU
1	B	37	ASP
1	B	100	ASP
1	B	139	LEU
1	B	147	GLN
1	B	174	GLU
1	B	208	ARG
1	B	216	ILE
1	B	263	THR
1	B	322	THR
1	B	330	LEU
1	B	346	THR
1	B	361	SER
1	B	413	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	4	ASN
1	A	224	GLN
1	A	248	GLN
1	A	253	ASN
1	A	302	ASN
1	A	314	ASN
1	B	50	HIS
1	B	59	ASN
1	B	122	GLN
1	B	147	GLN
1	B	224	GLN
1	B	248	GLN
1	B	253	ASN
1	B	302	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	411/437 (94%)	-1.32	0 100 100	33, 53, 74, 85	0
1	B	411/437 (94%)	-1.32	0 100 100	33, 53, 74, 81	0
All	All	822/874 (94%)	-1.32	0 100 100	33, 53, 74, 85	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.