



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 12:36 AM UTC

PDB ID : 3GQU / pdb_00003gqu
Title : Pyrococcus Horikoshii NOP5 RNA Binding Domain
Authors : Reyes, F.E.; Hardin, J.W.; Batey, R.T.
Deposited on : 2009-03-24
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	169	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	IOD	A	1	-	-	X	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	IOD	A	2	-	-	X	-
2	IOD	A	4	-	-	X	-
2	IOD	A	9	-	-	X	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 881 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NOP5P PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	117	Total	C	N	O	S	0	0	0
			811	520	138	151	2			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	222	GLY	-	expression tag	UNP O57810
A	239	GLY	-	linker	UNP O57810
A	240	ALA	-	linker	UNP O57810
A	241	GLY	-	linker	UNP O57810
A	242	GLY	-	linker	UNP O57810

- Molecule 2 is IODIDE ION (CCD ID: IOD) (formula: I).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	12	Total	I	0	0
			12	12		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	58	Total	O	0	0
			58	58		

i

- Molecule 1: NOP5P PROTEIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	73.60Å 73.60Å 114.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	23.68 – 2.50 23.68 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (23.68-2.50) 99.8 (23.68-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.15	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.54 (at 1.96Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.299 , 0.334 0.276 , 0.315	Depositor DCC
R_{free} test set	1971 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	43.6	Xtriage
Anisotropy	0.088	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 69.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	881	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IOD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	0/825	1.24	7/1126 (0.6%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	352	ILE	CB-CA-C	-10.83	98.30	111.94
1	A	322	SER	CA-C-N	6.92	126.96	119.90
1	A	322	SER	C-N-CA	6.92	126.96	119.90
1	A	356	LEU	N-CA-C	-6.28	104.44	111.28
1	A	355	GLU	N-CA-C	-5.76	105.84	112.92
1	A	324	TRP	N-CA-C	-5.68	98.70	110.80
1	A	353	ALA	N-CA-C	5.32	122.14	110.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	811	0	751	62	0
2	A	12	0	0	11	0
3	A	58	0	0	8	0
All	All	881	0	751	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 43.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:1:IOD:I	2:A:2:IOD:I	2.97	1.21
1:A:293:ALA:H	1:A:311:GLY:HA2	1.18	1.08
1:A:370:LYS:HA	3:A:393:HOH:O	1.62	0.99
1:A:292:GLY:HA3	1:A:312:VAL:H	1.32	0.94
2:A:1:IOD:I	3:A:36:HOH:O	2.53	0.94
1:A:292:GLY:HA3	1:A:312:VAL:HG13	1.64	0.80
2:A:6:IOD:I	2:A:7:IOD:I	3.45	0.75
1:A:293:ALA:N	1:A:311:GLY:HA2	2.02	0.71
1:A:262:VAL:HB	1:A:266:LEU:HD23	1.71	0.71
1:A:269:ARG:HG3	1:A:273:LEU:HD22	1.74	0.70
1:A:292:GLY:CA	1:A:312:VAL:HG13	2.22	0.70
2:A:4:IOD:I	3:A:26:HOH:O	2.81	0.68
1:A:292:GLY:HA3	1:A:312:VAL:N	2.06	0.68
1:A:284:PRO:O	1:A:285:SER:HB3	1.94	0.67
1:A:370:LYS:O	1:A:371:TYR:CB	2.42	0.67
2:A:3:IOD:I	3:A:27:HOH:O	2.83	0.67
1:A:323:PRO:C	1:A:324:TRP:O	2.33	0.67
1:A:261:LEU:HD21	1:A:337:LYS:HB3	1.77	0.66
1:A:318:ALA:HA	2:A:11:IOD:I	2.67	0.65
1:A:324:TRP:NE1	1:A:325:TRP:HE3	1.94	0.65
1:A:284:PRO:O	1:A:285:SER:CB	2.46	0.64
1:A:249:LYS:O	1:A:252:ASP:HB2	1.98	0.63
1:A:269:ARG:HD3	3:A:36:HOH:O	1.98	0.62
1:A:325:TRP:HE1	1:A:369:GLU:CB	2.14	0.61
1:A:354:GLU:CB	3:A:45:HOH:O	2.49	0.60
1:A:366:GLU:C	1:A:368:LYS:H	2.09	0.60
1:A:312:VAL:O	1:A:315:GLN:HG3	2.03	0.58
1:A:269:ARG:HG3	1:A:273:LEU:CD2	2.33	0.58
1:A:244:GLU:HB2	2:A:4:IOD:I	2.75	0.57
1:A:257:ASN:ND2	1:A:345:ASP:OD1	2.38	0.56
1:A:292:GLY:CA	1:A:312:VAL:H	2.13	0.56
1:A:353:ALA:O	1:A:355:GLU:N	2.40	0.55
1:A:261:LEU:HD23	1:A:338:LEU:HD23	1.88	0.54
1:A:354:GLU:N	3:A:45:HOH:O	2.40	0.54
1:A:293:ALA:HB2	1:A:310:HIS:O	2.08	0.53
1:A:360:LEU:CB	3:A:56:HOH:O	2.57	0.53
1:A:337:LYS:NZ	2:A:9:IOD:I	3.11	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:310:HIS:HA	1:A:314:TYR:CE1	2.44	0.51
1:A:339:ALA:O	1:A:343:ARG:HG3	2.10	0.51
1:A:344:VAL:HG21	1:A:352:ILE:CD1	2.40	0.51
1:A:337:LYS:HE2	2:A:9:IOD:I	2.81	0.51
1:A:323:PRO:O	1:A:327:ARG:HG3	2.11	0.50
1:A:276:GLY:N	1:A:279:GLU:HG2	2.27	0.50
1:A:310:HIS:HB3	1:A:314:TYR:CG	2.47	0.50
1:A:355:GLU:HA	1:A:358:LYS:HG3	1.94	0.49
1:A:313:ILE:HB	1:A:334:LEU:HD23	1.95	0.49
1:A:337:LYS:CE	2:A:9:IOD:I	3.32	0.48
1:A:352:ILE:O	1:A:354:GLU:N	2.47	0.47
1:A:366:GLU:C	1:A:368:LYS:N	2.73	0.47
1:A:353:ALA:C	1:A:355:GLU:H	2.22	0.47
1:A:324:TRP:CD2	1:A:325:TRP:HB3	2.50	0.47
1:A:266:LEU:HB2	2:A:2:IOD:I	2.84	0.46
1:A:279:GLU:CD	1:A:279:GLU:H	2.22	0.46
1:A:256:PRO:HD2	1:A:345:ASP:OD1	2.16	0.46
1:A:324:TRP:CD1	1:A:325:TRP:HE3	2.35	0.44
1:A:316:TYR:CD1	1:A:317:PRO:HD2	2.53	0.43
1:A:250:ALA:O	1:A:254:VAL:HG23	2.18	0.43
1:A:269:ARG:O	1:A:273:LEU:HD22	2.18	0.43
1:A:344:VAL:HG21	1:A:352:ILE:HD12	1.99	0.43
1:A:324:TRP:NE1	1:A:325:TRP:CE3	2.82	0.43
1:A:261:LEU:HD12	1:A:261:LEU:HA	1.81	0.42
1:A:326:GLN:HE21	1:A:369:GLU:CB	2.32	0.42
1:A:344:VAL:HG13	1:A:348:SER:HG	1.83	0.42
1:A:344:VAL:HG21	1:A:352:ILE:HD11	2.02	0.40
1:A:366:GLU:O	1:A:368:LYS:N	2.53	0.40
1:A:344:VAL:HG13	1:A:348:SER:OG	2.21	0.40
1:A:359:GLU:O	1:A:360:LEU:C	2.64	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	113/169 (67%)	97 (86%)	11 (10%)	5 (4%)	2 2

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	353	ALA
1	A	354	GLU
1	A	371	TYR
1	A	285	SER
1	A	367	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	67/132 (51%)	57 (85%)	10 (15%)	3 6

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	243	LEU
1	A	272	SER
1	A	273	LEU
1	A	279	GLU
1	A	280	LEU
1	A	322	SER
1	A	325	TRP
1	A	348	SER
1	A	350	GLU
1	A	359	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	315	GLN
1	A	326	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 12 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	117/169 (69%)	0.98	11 (9%) 14 12	34, 48, 61, 74	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	293	ALA	4.4
1	A	368	LYS	4.3
1	A	366	GLU	3.0
1	A	354	GLU	2.9
1	A	246	TYR	2.6
1	A	292	GLY	2.5
1	A	352	ILE	2.4
1	A	333	ALA	2.2
1	A	353	ALA	2.2
1	A	241	GLY	2.1
1	A	371	TYR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	IOD	A	9	1/1	0.86	0.38	169,169,169,169	0
2	IOD	A	10	1/1	0.86	0.17	155,155,155,155	0
2	IOD	A	2	1/1	0.89	0.33	154,154,154,154	0
2	IOD	A	8	1/1	0.92	0.23	137,137,137,137	0
2	IOD	A	11	1/1	0.92	0.16	165,165,165,165	0
2	IOD	A	12	1/1	0.92	0.30	123,123,123,123	0
2	IOD	A	5	1/1	0.93	0.23	124,124,124,124	0
2	IOD	A	4	1/1	0.93	0.33	143,143,143,143	0
2	IOD	A	7	1/1	0.96	0.15	120,120,120,120	0
2	IOD	A	1	1/1	0.97	0.08	72,72,72,72	0
2	IOD	A	3	1/1	0.97	0.12	103,103,103,103	0
2	IOD	A	6	1/1	0.99	0.09	63,63,63,63	0

6.5 Other polymers [i](#)

There are no such residues in this entry.