



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:23 PM UTC

PDB ID : 5GQU / pdb_00005gqu
Title : Crystal structure of branching enzyme from *Cyanothece* sp. ATCC 51142
Authors : Suzuki, R.; Suzuki, E.
Deposited on : 2016-08-08
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	793	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	801	-	X	-	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	807	-	X	-	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7177 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1,4-alpha-glucan branching enzyme GlgB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	755	6279	4061	1043	1151	24	0	1	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP B1WPM8
A	-18	GLY	-	expression tag	UNP B1WPM8
A	-17	SER	-	expression tag	UNP B1WPM8
A	-16	SER	-	expression tag	UNP B1WPM8
A	-15	HIS	-	expression tag	UNP B1WPM8
A	-14	HIS	-	expression tag	UNP B1WPM8
A	-13	HIS	-	expression tag	UNP B1WPM8
A	-12	HIS	-	expression tag	UNP B1WPM8
A	-11	HIS	-	expression tag	UNP B1WPM8
A	-10	HIS	-	expression tag	UNP B1WPM8
A	-9	SER	-	expression tag	UNP B1WPM8
A	-8	SER	-	expression tag	UNP B1WPM8
A	-7	GLY	-	expression tag	UNP B1WPM8
A	-6	LEU	-	expression tag	UNP B1WPM8
A	-5	VAL	-	expression tag	UNP B1WPM8
A	-4	PRO	-	expression tag	UNP B1WPM8
A	-3	ARG	-	expression tag	UNP B1WPM8
A	-2	GLY	-	expression tag	UNP B1WPM8
A	-1	SER	-	expression tag	UNP B1WPM8
A	0	HIS	-	expression tag	UNP B1WPM8

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Mg	0	0
			4	4		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	852	Total	O	0	0
			852	852		

- Molecule 1: 1,4-alpha-glucan branching enzyme GlgB



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	133.75Å 133.75Å 185.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.29 – 1.85 47.29 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.29-1.85) 99.9 (47.29-1.85)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	9.72 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.147 , 0.170 0.160 , 0.180	Depositor DCC
R_{free} test set	7187 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	21.5	Xtriage
Anisotropy	0.020	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 39.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7177	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.73	71/6502 (1.1%)	1.38	33/8848 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (71) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	276	GLU	CA-C	-9.54	1.41	1.52
1	A	434	ASP	N-CA	9.29	1.58	1.45
1	A	242	ARG	CZ-NH1	9.27	1.45	1.32
1	A	412	ARG	CD-NE	-9.04	1.33	1.46
1	A	564	MET	SD-CE	-8.93	1.57	1.79
1	A	274	ASN	N-CA	8.74	1.57	1.46
1	A	343	ARG	CZ-NH2	-8.14	1.22	1.33
1	A	672	SER	CA-C	7.96	1.60	1.53
1	A	104	PHE	C-O	7.82	1.32	1.24
1	A	112	GLU	CA-C	7.50	1.62	1.52
1	A	481	GLY	CA-C	7.45	1.62	1.51
1	A	412	ARG	CZ-NH2	7.36	1.43	1.33
1	A	480	PRO	C-O	7.23	1.32	1.24
1	A	148	ALA	N-CA	7.00	1.56	1.46
1	A	179	GLU	CA-C	-6.99	1.44	1.52
1	A	481	GLY	N-CA	6.96	1.55	1.45
1	A	216	ARG	CA-C	6.87	1.61	1.53
1	A	369	HIS	CA-C	-6.83	1.44	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	662	THR	CB-CG2	-6.83	1.30	1.52
1	A	673	ASP	N-CA	6.82	1.54	1.46
1	A	36	THR	C-O	6.79	1.31	1.23
1	A	117	PHE	N-CA	6.77	1.54	1.46
1	A	649	GLU	N-CA	6.72	1.54	1.45
1	A	663	HIS	CA-C	6.67	1.61	1.52
1	A	662	THR	N-CA	-6.66	1.37	1.46
1	A	753	LEU	N-CA	6.63	1.54	1.46
1	A	439	MET	SD-CE	-6.54	1.63	1.79
1	A	558	VAL	CA-C	6.50	1.60	1.53
1	A	686	THR	N-CA	-6.46	1.41	1.46
1	A	527	TYR	N-CA	6.32	1.54	1.46
1	A	451	TRP	CA-C	-6.24	1.44	1.52
1	A	658	CYS	CA-CB	6.19	1.63	1.53
1	A	524	ILE	N-CA	6.15	1.53	1.46
1	A	465	VAL	N-CA	-6.09	1.39	1.46
1	A	648	PHE	N-CA	-6.08	1.39	1.46
1	A	171	ARG	CA-C	-5.98	1.45	1.52
1	A	218	ASN	CA-CB	-5.96	1.43	1.53
1	A	104	PHE	CA-C	-5.92	1.45	1.52
1	A	139	VAL	C-O	5.88	1.29	1.24
1	A	745	PRO	CA-CB	-5.82	1.46	1.53
1	A	302	ILE	CA-CB	-5.77	1.51	1.54
1	A	69	HIS	N-CA	-5.77	1.38	1.46
1	A	534	ASN	CA-C	5.75	1.60	1.52
1	A	752	LYS	CA-C	5.64	1.59	1.52
1	A	401	THR	N-CA	5.62	1.53	1.45
1	A	58	GLU	CD-OE2	5.60	1.35	1.25
1	A	643	LEU	N-CA	5.53	1.53	1.46
1	A	724	GLY	N-CA	-5.52	1.40	1.45
1	A	173	ARG	CA-C	-5.45	1.46	1.52
1	A	521	TYR	CA-C	-5.40	1.46	1.52
1	A	726	TRP	CA-CB	5.39	1.61	1.53
1	A	331	VAL	CA-CB	5.37	1.61	1.54
1	A	242	ARG	CD-NE	5.34	1.53	1.46
1	A	154	VAL	CA-C	-5.33	1.46	1.52
1	A	676	ASN	CA-C	5.33	1.59	1.52
1	A	659	ASN	N-CA	5.31	1.53	1.46
1	A	169	GLN	N-CA	-5.28	1.39	1.46
1	A	51	ALA	CA-C	-5.26	1.46	1.52
1	A	46	PRO	C-N	-5.26	1.27	1.33
1	A	518	MET	N-CA	-5.21	1.40	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	273	LEU	CA-C	5.17	1.59	1.52
1	A	285	TRP	CB-CG	-5.17	1.34	1.50
1	A	299	GLU	CD-OE2	5.12	1.35	1.25
1	A	61	GLU	CA-CB	-5.06	1.45	1.53
1	A	218	ASN	CG-OD1	-5.06	1.14	1.23
1	A	59	ARG	CA-C	5.06	1.59	1.53
1	A	177	ILE	C-O	5.06	1.29	1.24
1	A	218	ASN	CG-ND2	-5.05	1.22	1.33
1	A	747	ALA	CA-C	-5.04	1.46	1.52
1	A	752	LYS	C-O	-5.02	1.18	1.23
1	A	248	LYS	CA-C	-5.01	1.45	1.52

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	412	ARG	NE-CZ-NH2	-12.38	108.05	119.20
1	A	412	ARG	CD-NE-CZ	12.10	141.33	124.40
1	A	658	CYS	CB-CA-C	9.33	126.12	110.44
1	A	242	ARG	NE-CZ-NH2	-8.03	111.98	119.20
1	A	480	PRO	CA-C-N	-7.59	106.53	121.41
1	A	480	PRO	C-N-CA	-7.59	106.53	121.41
1	A	412	ARG	NE-CZ-NH1	7.42	128.92	121.50
1	A	118	SER	CA-CB-OG	-7.33	96.45	111.10
1	A	68	HIS	N-CA-C	7.07	125.85	110.80
1	A	662	THR	N-CA-CB	-7.03	98.37	110.32
1	A	24	ILE	CB-CA-C	-6.52	104.52	110.91
1	A	184	GLU	CB-CA-C	-6.38	100.96	111.36
1	A	659	ASN	N-CA-C	6.32	120.73	112.89
1	A	321	HIS	CA-C-N	-5.98	113.60	119.64
1	A	321	HIS	C-N-CA	-5.98	113.60	119.64
1	A	58	GLU	CA-CB-CG	5.92	125.93	114.10
1	A	104	PHE	CA-C-O	5.74	126.52	120.33
1	A	528	PHE	N-CA-C	5.63	119.99	113.18
1	A	173	ARG	CA-CB-CG	-5.49	103.11	114.10
1	A	390	ASP	CA-C-N	5.43	125.10	119.56
1	A	390	ASP	C-N-CA	5.43	125.10	119.56
1	A	619	GLN	CB-CG-CD	-5.33	103.54	112.60
1	A	282	VAL	N-CA-C	5.29	116.01	110.62
1	A	46	PRO	CA-C-N	-5.17	115.92	125.02
1	A	46	PRO	C-N-CA	-5.17	115.92	125.02
1	A	462	ILE	N-CA-C	5.16	115.88	110.62
1	A	116	LEU	O-C-N	5.12	127.35	122.07

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	340	ARG	NE-CZ-NH2	5.12	123.80	119.20
1	A	179	GLU	N-CA-C	5.11	117.44	109.52
1	A	55	ARG	CA-C-O	5.10	127.63	120.64
1	A	16	ASN	N-CA-CB	-5.07	104.29	111.84
1	A	106	SER	O-C-N	5.03	124.93	121.71
1	A	33	GLY	N-CA-C	-5.02	105.70	112.57

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	184	GLU	Mainchain
1	A	412	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6279	0	5865	5	0
2	A	42	0	56	2	0
3	A	4	0	0	0	0
4	A	852	0	0	1	0
All	All	7177	0	5921	6	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (6) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:46:PRO:O	1:A:47:LYS:HB2	2.05	0.56
1:A:518:MET:HE2	1:A:518:MET:HA	1.98	0.45
2:A:801:GOL:H2	4:A:967:HOH:O	2.16	0.45
1:A:182:VAL:HA	1:A:183:PRO:HD2	1.94	0.42
1:A:244:SER:HB3	1:A:249:GLN:NE2	2.36	0.40
1:A:528:PHE:CD2	2:A:802:GOL:H2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	754/793 (95%)	734 (97%)	19 (2%)	1 (0%)	48 35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	68	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	669/700 (96%)	662 (99%)	7 (1%)	68 60

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	184	GLU
1	A	434	ASP
1	A	447	LYS
1	A	491	GLU
1	A	662	THR
1	A	686	THR
1	A	725	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	16	ASN
1	A	35	ASN
1	A	286	ASN
1	A	634	ASN
1	A	688	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	804	-	5,5,5	1.12	0	5,5,5	1.41	1 (20%)
2	GOL	A	801	-	5,5,5	1.78	2 (40%)	5,5,5	2.92	3 (60%)
2	GOL	A	802	-	5,5,5	2.75	3 (60%)	5,5,5	2.22	1 (20%)
2	GOL	A	806	-	5,5,5	0.83	0	5,5,5	1.99	1 (20%)
2	GOL	A	803	-	5,5,5	0.98	0	5,5,5	0.55	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	805	-	5,5,5	0.35	0	5,5,5	1.31	1 (20%)
2	GOL	A	807	-	5,5,5	2.07	2 (40%)	5,5,5	2.25	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	804	-	-	2/4/4/4	-
2	GOL	A	801	-	-	2/4/4/4	-
2	GOL	A	802	-	-	0/4/4/4	-
2	GOL	A	806	-	-	0/4/4/4	-
2	GOL	A	803	-	-	0/4/4/4	-
2	GOL	A	805	-	-	4/4/4/4	-
2	GOL	A	807	-	-	3/4/4/4	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	802	GOL	O3-C3	4.20	1.60	1.42
2	A	807	GOL	O3-C3	3.80	1.58	1.42
2	A	802	GOL	C3-C2	2.98	1.63	1.51
2	A	801	GOL	O2-C2	2.84	1.51	1.43
2	A	802	GOL	O2-C2	2.72	1.51	1.43
2	A	801	GOL	C3-C2	2.30	1.60	1.51
2	A	807	GOL	O2-C2	-2.24	1.36	1.43

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	GOL	O2-C2-C3	4.77	128.91	109.18
2	A	802	GOL	O3-C3-C2	4.05	128.63	110.38
2	A	806	GOL	C3-C2-C1	3.52	124.71	111.80
2	A	801	GOL	O1-C1-C2	3.46	125.93	110.38
2	A	807	GOL	O1-C1-C2	3.32	125.31	110.38
2	A	807	GOL	O3-C3-C2	2.90	123.42	110.38
2	A	804	GOL	O3-C3-C2	2.66	122.34	110.38
2	A	801	GOL	C3-C2-C1	-2.47	102.75	111.80
2	A	805	GOL	O1-C1-C2	2.19	120.23	110.38

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	801	GOL	C1-C2-C3-O3
2	A	804	GOL	C1-C2-C3-O3
2	A	804	GOL	O2-C2-C3-O3
2	A	805	GOL	O1-C1-C2-C3
2	A	805	GOL	C1-C2-C3-O3
2	A	807	GOL	O1-C1-C2-C3
2	A	805	GOL	O2-C2-C3-O3
2	A	801	GOL	O2-C2-C3-O3
2	A	805	GOL	O1-C1-C2-O2
2	A	807	GOL	O1-C1-C2-O2
2	A	807	GOL	C1-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	GOL	1	0
2	A	802	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	755/793 (95%)	-0.22	10 (1%) 75 79	14, 21, 42, 77	1 (0%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	759	PRO	7.0
1	A	758	VAL	6.1
1	A	285	TRP	3.8
1	A	500	TYR	3.1
1	A	106	SER	3.0
1	A	35	ASN	3.0
1	A	456	TYR	2.4
1	A	105	SER	2.3
1	A	81	GLU	2.2
1	A	307	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	A	801	6/6	0.74	0.20	33,41,43,51	0
2	GOL	A	804	6/6	0.86	0.16	37,54,58,58	0
2	GOL	A	806	6/6	0.86	0.19	33,53,59,60	0
2	GOL	A	802	6/6	0.87	0.18	27,43,55,55	0
2	GOL	A	805	6/6	0.88	0.16	45,53,58,58	0
3	MG	A	811	1/1	0.88	0.20	44,44,44,44	0
2	GOL	A	807	6/6	0.92	0.15	27,41,44,45	0
2	GOL	A	803	6/6	0.94	0.09	33,34,35,35	0
3	MG	A	808	1/1	0.97	0.12	40,40,40,40	0
3	MG	A	809	1/1	0.98	0.03	22,22,22,22	0
3	MG	A	810	1/1	0.99	0.05	23,23,23,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.