



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 03:04 PM UTC

PDB ID : 3GR0 / pdb\_00003gr0  
Title : Periplasmic domain of the T3SS inner membrane protein PrgH from *S.typhimurium* (fragment 170-362)  
Authors : Yip, C.K.; Vockovic, M.; Yu, A.C.; Strynadka, N.C.J.  
Deposited on : 2009-03-24  
Resolution : 2.30 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	197	 74% 19% • 6%
1	B	197	 82% 14% • •
1	C	197	 71% 16% • 11%
1	D	197	 69% 21% • 9%

## 2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 6311 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein prgH.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	186	Total 1541	C 976	N 280	O 281	S 4	0	0	0
1	B	192	Total 1585	C 1003	N 286	O 292	S 4	0	0	0
1	C	176	Total 1461	C 928	N 268	O 261	S 4	0	0	0
1	D	180	Total 1493	C 947	N 272	O 270	S 4	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	166	GLY	-	expression tag	UNP P41783
A	167	SER	-	expression tag	UNP P41783
A	168	HIS	-	expression tag	UNP P41783
A	169	MET	-	expression tag	UNP P41783
B	166	GLY	-	expression tag	UNP P41783
B	167	SER	-	expression tag	UNP P41783
B	168	HIS	-	expression tag	UNP P41783
B	169	MET	-	expression tag	UNP P41783
C	166	GLY	-	expression tag	UNP P41783
C	167	SER	-	expression tag	UNP P41783
C	168	HIS	-	expression tag	UNP P41783
C	169	MET	-	expression tag	UNP P41783
D	166	GLY	-	expression tag	UNP P41783
D	167	SER	-	expression tag	UNP P41783
D	168	HIS	-	expression tag	UNP P41783
D	169	MET	-	expression tag	UNP P41783

- Molecule 2 is water.


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
2	A	66	Total 66	O 66	0	0
2	B	69	Total 69	O 69	0	0
2	C	41	Total 41	O 41	0	0
2	D	55	Total 55	O 55	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Protein prgH

Chain A:  74% 19% 6%




- Molecule 1: Protein prgH

Chain B:  82% 14%




- Molecule 1: Protein prgH


Chain C:  71% 16% 11%



- Molecule 1: Protein prgH

Chain D:  69% 21% 9%





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	53.26Å 53.26Å 282.53Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	35.00 – 2.30 35.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.5 (35.00-2.30) 99.5 (35.00-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.92 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.206 , 0.255 0.200 , 0.246	Depositor DCC
$R_{free}$ test set	2521 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.8	Xtrriage
Anisotropy	0.343	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 25.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.024 for -h,-k,l 0.487 for h,-h-k,-l 0.027 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6311	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.93% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.78	1/1573 (0.1%)	0.94	2/2122 (0.1%)
1	B	0.79	1/1617 (0.1%)	0.92	0/2182
1	C	0.68	0/1491	0.87	0/2011
1	D	0.68	0/1525	0.90	0/2059
All	All	0.74	2/6206 (0.0%)	0.91	2/8374 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	218	LYS	CE-NZ	8.75	1.75	1.49
1	A	196	VAL	CA-CB	5.75	1.60	1.53

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	312	LEU	CA-C-N	-5.46	114.33	119.85
1	A	312	LEU	C-N-CA	-5.46	114.33	119.85

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1541	0	1530	31	0
1	B	1585	0	1572	19	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1461	0	1462	27	0
1	D	1493	0	1483	28	0
2	A	66	0	0	0	0
2	B	69	0	0	1	0
2	C	41	0	0	0	0
2	D	55	0	0	3	0
All	All	6311	0	6047	96	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 96 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:218:LYS:CE	1:B:218:LYS:NZ	1.75	1.48
1:D:263:GLN:HE22	1:D:296:ASP:H	1.25	0.81
1:C:263:GLN:HE22	1:C:296:ASP:H	1.27	0.79
1:B:263:GLN:HE21	1:B:263:GLN:H	1.30	0.78
1:A:263:GLN:HE21	1:A:263:GLN:H	1.30	0.77

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	184/197 (93%)	180 (98%)	4 (2%)	0	100	100
1	B	190/197 (96%)	185 (97%)	4 (2%)	1 (0%)	24	31
1	C	172/197 (87%)	170 (99%)	2 (1%)	0	100	100
1	D	178/197 (90%)	173 (97%)	4 (2%)	1 (1%)	21	27
All	All	724/788 (92%)	708 (98%)	14 (2%)	2 (0%)	36	46

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	215	ASP
1	B	320	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	160/168 (95%)	150 (94%)	10 (6%)	16	23
1	B	165/168 (98%)	152 (92%)	13 (8%)	11	15
1	C	152/168 (90%)	142 (93%)	10 (7%)	15	21
1	D	155/168 (92%)	142 (92%)	13 (8%)	10	14
All	All	632/672 (94%)	586 (93%)	46 (7%)	13	18

5 of 46 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	269	LYS
1	D	201	GLU
1	C	307	LEU
1	C	360	GLU
1	D	218	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	341	GLN
1	D	263	GLN
1	D	290	ASN
1	D	226	ASN
1	B	242	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	186/197 (94%)	-1.44	0 100 100	10, 20, 35, 44	0
1	B	192/197 (97%)	-1.40	0 100 100	8, 22, 35, 51	0
1	C	176/197 (89%)	-1.39	0 100 100	14, 23, 34, 38	0
1	D	180/197 (91%)	-1.39	0 100 100	13, 24, 35, 40	0
All	All	734/788 (93%)	-1.41	0 100 100	8, 22, 35, 51	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.