



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 08:01 PM UTC

PDB ID : 5GRK / pdb_00005grk
Title : Crystal structure of Uracil DNA glycosylase -Xanthine complex from Bradyrhizobium diazoefficiens
Authors : Patil, V.V.; Ullas, V.C.; Ahn, W.; Varshney, U.; Woo, E.
Deposited on : 2016-08-11
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	272	2% 96% .
1	B	272	4% 94% 6%
1	C	272	31% 90% 10%
1	D	272	5% 95% 5%

2 Entry composition [i](#)


There are 2 unique types of molecules in this entry. The entry contains 16682 atoms, of which 8242 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Blr0248 protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	272	Total	C	H	N	O	S	0	0	0
			4158	1344	2059	368	381	6			
1	B	272	Total	C	H	N	O	S	0	0	0
			4143	1344	2044	368	381	6			
1	C	272	Total	C	H	N	O	S	0	0	0
			4152	1344	2053	368	381	6			
1	D	272	Total	C	H	N	O	S	0	0	0
			4169	1344	2070	368	381	6			

- Molecule 2 is XANTHINE (CCD ID: XAN) (formula: C₅H₄N₄O₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	0
			15	5	4	4	2		
2	B	1	Total	C	H	N	O	0	0
			15	5	4	4	2		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	H	N	O	0	0
			15	5	4	4	2		
2	D	1	Total	C	H	N	O	0	0
			15	5	4	4	2		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Blr0248 protein




- Molecule 1: Blr0248 protein



- Molecule 1: Blr0248 protein



- Molecule 1: Blr0248 protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	70.02Å 89.89Å 255.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.17 – 2.80 47.17 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.7 (47.17-2.80) 99.7 (47.17-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.25 (at 2.81Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.220 , 0.244 0.231 , 0.246	Depositor DCC
R_{free} test set	2000 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	37.6	Xtriage
Anisotropy	1.020	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 32.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	16682	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: XAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	1/2159 (0.0%)	0.67	0/2944
1	B	0.54	0/2159	0.67	0/2944
1	C	0.40	0/2159	0.69	0/2944
1	D	0.44	0/2159	0.68	1/2944 (0.0%)
All	All	0.47	1/8636 (0.0%)	0.68	1/11776 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	42	ARG	C-O	-5.36	1.17	1.23

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	104	SER	N-CA-C	5.11	118.97	112.12

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2099	2059	2072	10	0
1	B	2099	2044	2072	29	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2099	2053	2072	34	0
1	D	2099	2070	2072	9	0
2	A	11	4	4	0	0
2	B	11	4	4	1	0
2	C	11	4	4	2	0
2	D	11	4	4	0	0
All	All	8440	8242	8304	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:102:GLY:HA3	1:B:103:GLN:CG	1.42	1.40
1:B:102:GLY:CA	1:B:103:GLN:HG2	1.30	1.29
1:B:102:GLY:HA3	1:B:103:GLN:CD	1.56	1.28
1:B:102:GLY:CA	1:B:103:GLN:CG	2.04	1.14
1:B:102:GLY:HA3	1:B:103:GLN:OE1	1.56	1.05
1:B:103:GLN:HA	1:B:104:SER:HB2	1.40	1.03
1:B:103:GLN:NE2	1:B:104:SER:O	1.93	1.01
1:A:44:ARG:NH2	1:A:232:ASP:OD1	2.02	0.93
1:B:102:GLY:CA	1:B:103:GLN:OE1	2.16	0.93
1:C:33:ARG:NH1	1:C:259:ARG:HG3	1.84	0.91
1:C:107:SER:HB2	1:C:110:LYS:HB3	1.54	0.88
1:C:33:ARG:HH12	1:C:259:ARG:HG3	1.39	0.86
1:C:104:SER:N	1:C:105:GLY:HA3	1.92	0.85
1:A:102:GLY:HA2	1:A:103:GLN:HB2	1.58	0.84
1:B:103:GLN:HA	1:B:104:SER:CB	2.06	0.81
1:A:102:GLY:CA	1:A:103:GLN:HB2	2.15	0.76
1:C:107:SER:OG	1:C:141:LEU:CD2	2.35	0.74
1:B:58:PRO:O	1:B:99:SER:OG	2.04	0.74
1:B:44:ARG:NH1	1:B:88:GLN:O	2.21	0.74
1:B:44:ARG:NH2	1:B:232:ASP:OD1	2.21	0.73
1:B:103:GLN:HB3	1:B:104:SER:C	2.17	0.69
1:B:102:GLY:CA	1:B:103:GLN:CD	2.40	0.69
1:C:33:ARG:NH2	1:C:101:TYR:CD1	2.63	0.67
1:C:103:GLN:C	1:C:105:GLY:HA3	2.21	0.66
1:C:69:LEU:N	2:C:301:XAN:O2	2.31	0.62
1:C:33:ARG:HD2	1:C:99:SER:OG	2.00	0.62
1:B:103:GLN:CD	1:B:104:SER:O	2.42	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:68:ILE:HD12	1:C:94:ASN:OD1	2.01	0.61
1:C:33:ARG:NH2	1:C:101:TYR:CE1	2.70	0.60
1:C:33:ARG:CZ	1:C:101:TYR:HD1	2.14	0.59
1:D:12:GLN:HB3	1:D:13:PRO:HA	1.85	0.57
1:B:102:GLY:C	1:B:103:GLN:HG2	2.21	0.57
1:C:57:ASP:OD2	1:C:100:VAL:HG23	2.05	0.57
1:C:33:ARG:NH2	1:C:101:TYR:HD1	2.03	0.56
1:B:102:GLY:HA2	1:B:103:GLN:OE1	2.02	0.56
1:D:104:SER:H	1:D:105:GLY:HA3	1.71	0.55
1:D:90:TYR:OH	1:D:232:ASP:OD2	2.25	0.54
1:C:33:ARG:CZ	1:C:101:TYR:CD1	2.91	0.53
1:C:106:GLY:N	1:C:107:SER:HA	2.24	0.52
1:C:57:ASP:CG	1:C:100:VAL:HG23	2.34	0.52
1:C:89:SER:OG	1:C:208:ASP:OD1	2.23	0.52
1:C:107:SER:OG	1:C:141:LEU:HD21	2.08	0.52
1:B:68:ILE:HD12	1:B:94:ASN:OD1	2.10	0.52
1:B:104:SER:N	1:B:105:GLY:O	2.44	0.51
1:B:33:ARG:HG3	1:B:99:SER:O	2.11	0.51
1:B:168:HIS:CE1	2:B:301:XAN:N7	2.79	0.51
1:C:107:SER:OG	1:C:141:LEU:HD23	2.11	0.50
1:C:107:SER:N	1:C:141:LEU:HD21	2.26	0.50
1:D:89:SER:OG	1:D:208:ASP:OD1	2.22	0.50
1:C:68:ILE:HB	1:C:94:ASN:HD21	1.78	0.48
1:B:34:ILE:HA	1:B:98:TYR:CD1	2.48	0.48
1:C:96:PHE:CZ	1:C:107:SER:OG	2.66	0.47
1:A:103:GLN:O	1:A:104:SER:C	2.56	0.47
1:B:44:ARG:NH1	1:B:44:ARG:HG2	2.29	0.47
1:B:104:SER:N	1:B:105:GLY:C	2.73	0.47
1:C:69:LEU:HG	2:C:301:XAN:O2	2.14	0.47
1:C:96:PHE:HZ	1:C:107:SER:OG	1.98	0.46
1:C:107:SER:H	1:C:141:LEU:HD21	1.81	0.46
1:C:108:LYS:HD2	1:C:109:HIS:NE2	2.32	0.45
1:A:102:GLY:HA3	1:A:104:SER:N	2.32	0.45
1:D:6:ASP:OD1	1:D:7:ALA:N	2.50	0.44
1:A:178:ASP:OD2	1:A:181:THR:OG1	2.33	0.44
1:D:12:GLN:CB	1:D:13:PRO:HA	2.45	0.44
1:B:44:ARG:HH11	1:B:44:ARG:CG	2.31	0.44
1:B:104:SER:N	1:B:105:GLY:CA	2.82	0.43
1:B:68:ILE:HB	1:B:94:ASN:HD21	1.82	0.43
1:C:104:SER:HA	1:C:108:LYS:H	1.83	0.43
1:A:102:GLY:HA3	1:A:103:GLN:C	2.44	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:171:TRP:N	1:C:172:PRO:CD	2.81	0.42
1:C:105:GLY:N	1:C:107:SER:HA	2.34	0.42
1:C:261:THR:HG21	1:D:263:MET:HE3	2.01	0.42
1:D:171:TRP:N	1:D:172:PRO:CD	2.82	0.42
1:B:58:PRO:O	1:B:99:SER:CB	2.68	0.41
1:C:103:GLN:HG2	1:C:103:GLN:O	2.21	0.41
1:A:41:HIS:CG	1:A:42:ARG:N	2.88	0.41
1:A:104:SER:O	1:A:104:SER:OG	2.28	0.41
1:C:1:MET:HE1	1:D:25:GLU:OE2	2.21	0.41
1:A:178:ASP:OD1	1:A:178:ASP:N	2.54	0.40
1:C:34:ILE:HA	1:C:98:TYR:CD2	2.57	0.40
1:B:103:GLN:CB	1:B:104:SER:C	2.92	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	270/272 (99%)	263 (97%)	7 (3%)	0	100	100
1	B	270/272 (99%)	260 (96%)	10 (4%)	0	100	100
1	C	270/272 (99%)	260 (96%)	10 (4%)	0	100	100
1	D	270/272 (99%)	261 (97%)	9 (3%)	0	100	100
All	All	1080/1088 (99%)	1044 (97%)	36 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	213/213 (100%)	212 (100%)	1 (0%)	81	93
1	B	213/213 (100%)	213 (100%)	0	100	100
1	C	213/213 (100%)	213 (100%)	0	100	100
1	D	213/213 (100%)	213 (100%)	0	100	100
All	All	852/852 (100%)	851 (100%)	1 (0%)	88	97

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	20	ASN
1	B	88	GLN
1	B	165	HIS
1	B	185	ASN
1	C	30	HIS
1	C	185	ASN
1	D	30	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	XAN	C	301	-	12,12,12	1.97	3 (25%)	14,17,17	2.47	7 (50%)
2	XAN	D	301	-	12,12,12	1.97	3 (25%)	14,17,17	2.47	8 (57%)
2	XAN	A	301	-	12,12,12	1.97	3 (25%)	14,17,17	2.47	7 (50%)
2	XAN	B	301	-	12,12,12	1.96	3 (25%)	14,17,17	2.47	7 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XAN	C	301	-	-	-	0/2/2/2
2	XAN	D	301	-	-	-	0/2/2/2
2	XAN	A	301	-	-	-	0/2/2/2
2	XAN	B	301	-	-	-	0/2/2/2

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	301	XAN	C5-C4	4.81	1.47	1.38
2	A	301	XAN	C5-C4	4.78	1.47	1.38
2	C	301	XAN	C5-C4	4.78	1.47	1.38
2	B	301	XAN	C5-C4	4.75	1.47	1.38
2	C	301	XAN	C6-N1	-2.46	1.34	1.38
2	A	301	XAN	C6-N1	-2.45	1.34	1.38
2	B	301	XAN	C6-N1	-2.42	1.34	1.38
2	D	301	XAN	C6-N1	-2.42	1.34	1.38
2	C	301	XAN	C5-N7	-2.04	1.35	1.39
2	B	301	XAN	C5-N7	-2.02	1.35	1.39
2	D	301	XAN	C5-N7	-2.02	1.35	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	XAN	C5-N7	-2.00	1.35	1.39

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	XAN	N1-C2-N3	4.83	123.34	115.74
2	D	301	XAN	N1-C2-N3	4.83	123.33	115.74
2	A	301	XAN	N1-C2-N3	4.83	123.33	115.74
2	C	301	XAN	N1-C2-N3	4.81	123.30	115.74
2	B	301	XAN	C6-N1-C2	-3.34	121.77	126.37
2	C	301	XAN	C6-N1-C2	-3.34	121.77	126.37
2	A	301	XAN	C6-N1-C2	-3.34	121.77	126.37
2	D	301	XAN	C6-N1-C2	-3.33	121.79	126.37
2	D	301	XAN	C6-C5-N7	3.22	136.15	130.29
2	A	301	XAN	C6-C5-N7	3.21	136.13	130.29
2	C	301	XAN	C6-C5-N7	3.21	136.13	130.29
2	B	301	XAN	C6-C5-N7	3.20	136.11	130.29
2	A	301	XAN	N9-C4-N3	3.10	134.94	128.34
2	D	301	XAN	N9-C4-N3	3.10	134.94	128.34
2	C	301	XAN	N9-C4-N3	3.09	134.94	128.34
2	B	301	XAN	N9-C4-N3	3.09	134.93	128.34
2	C	301	XAN	C8-N7-C5	2.63	107.93	104.38
2	D	301	XAN	C8-N7-C5	2.63	107.93	104.38
2	B	301	XAN	C8-N7-C5	2.62	107.92	104.38
2	A	301	XAN	C8-N7-C5	2.61	107.90	104.38
2	B	301	XAN	N9-C8-N7	-2.21	110.41	112.98
2	C	301	XAN	N9-C8-N7	-2.18	110.45	112.98
2	D	301	XAN	N9-C8-N7	-2.18	110.45	112.98
2	A	301	XAN	N9-C8-N7	-2.18	110.45	112.98
2	A	301	XAN	O6-C6-C5	-2.08	121.04	126.53
2	C	301	XAN	O6-C6-C5	-2.07	121.07	126.53
2	B	301	XAN	O6-C6-C5	-2.06	121.08	126.53
2	D	301	XAN	O6-C6-C5	-2.06	121.10	126.53
2	D	301	XAN	O2-C2-N3	-2.00	118.30	121.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	301	XAN	2	0
2	B	301	XAN	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	272/272 (100%)	0.17	6 (2%) 62 52	26, 35, 55, 115	0
1	B	272/272 (100%)	0.11	10 (3%) 45 36	24, 35, 50, 80	0
1	C	272/272 (100%)	1.42	83 (30%) 1 1	30, 62, 94, 127	0
1	D	272/272 (100%)	0.25	13 (4%) 35 28	30, 39, 61, 110	0
All	All	1088/1088 (100%)	0.49	112 (10%) 12 9	24, 39, 82, 127	0

All (112) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	103	GLN	8.3
1	B	102	GLY	8.2
1	B	103	GLN	7.3
1	C	105	GLY	7.1
1	B	104	SER	7.0
1	A	105	GLY	6.9
1	A	104	SER	6.4
1	C	218	ASP	5.8
1	A	103	GLN	5.7
1	C	217	GLY	5.5
1	B	106	GLY	5.4
1	C	177	HIS	5.2
1	C	106	GLY	5.1
1	D	8	GLY	5.0
1	C	171	TRP	4.9
1	C	195	ALA	4.8
1	D	1	MET	4.5
1	A	1	MET	4.4
1	C	200	LEU	4.4
1	C	219	ALA	4.2
1	C	178	ASP	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	109	HIS	4.0
1	C	154	ASP	3.9
1	C	99	SER	3.9
1	C	220	PHE	3.8
1	D	177	HIS	3.7
1	D	105	GLY	3.7
1	C	194	ASN	3.7
1	B	177	HIS	3.7
1	C	1	MET	3.7
1	C	210	PRO	3.6
1	C	107	SER	3.6
1	C	104	SER	3.6
1	C	153	SER	3.6
1	C	102	GLY	3.5
1	B	105	GLY	3.4
1	C	114	GLY	3.4
1	D	104	SER	3.4
1	C	198	ALA	3.4
1	D	7	ALA	3.4
1	C	176	ALA	3.4
1	D	108	LYS	3.3
1	C	196	ALA	3.3
1	C	190	LEU	3.3
1	C	103	GLN	3.3
1	C	159	LYS	3.2
1	A	177	HIS	3.2
1	C	163	TYR	3.2
1	D	102	GLY	3.2
1	C	193	TRP	3.2
1	C	164	GLN	3.1
1	C	206	HIS	3.1
1	C	181	THR	3.1
1	C	139	GLY	3.0
1	C	166	ILE	3.0
1	C	175	SER	3.0
1	C	15	ARG	2.9
1	C	202	PRO	2.9
1	C	185	ASN	2.9
1	C	191	ALA	2.8
1	A	106	GLY	2.8
1	C	4	GLU	2.7
1	C	47	GLY	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	149	TRP	2.6
1	D	2	LEU	2.6
1	C	192	LYS	2.6
1	C	205	LYS	2.6
1	B	99	SER	2.6
1	C	161	LEU	2.6
1	C	213	LEU	2.6
1	B	100	VAL	2.6
1	C	212	THR	2.6
1	C	215	PRO	2.5
1	C	141	LEU	2.5
1	D	263	MET	2.5
1	C	203	GLU	2.5
1	C	167	THR	2.5
1	C	211	THR	2.5
1	C	142	ALA	2.4
1	C	137	SER	2.4
1	C	170	THR	2.4
1	C	126	VAL	2.3
1	C	151	LYS	2.3
1	C	155	GLY	2.3
1	C	133	GLU	2.3
1	C	204	VAL	2.3
1	C	32	PHE	2.3
1	C	131	ASN	2.3
1	D	178	ASP	2.3
1	C	160	THR	2.3
1	C	172	PRO	2.3
1	C	165	HIS	2.3
1	C	120	ASN	2.2
1	C	140	GLY	2.2
1	C	179	SER	2.2
1	C	243	GLY	2.2
1	C	127	LEU	2.2
1	C	199	ALA	2.2
1	C	33	ARG	2.2
1	C	144	GLU	2.2
1	C	201	ALA	2.2
1	C	121	LYS	2.1
1	D	106	GLY	2.1
1	C	128	GLY	2.1
1	C	162	ALA	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	1	MET	2.1
1	C	146	TRP	2.1
1	C	183	ALA	2.1
1	C	207	PRO	2.1
1	B	107	SER	2.0
1	C	26	ALA	2.0
1	C	87	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	XAN	C	301	11/11	0.84	0.14	51,57,64,73	0
2	XAN	B	301	11/11	0.94	0.10	25,31,36,37	0
2	XAN	A	301	11/11	0.94	0.11	32,36,42,45	0
2	XAN	D	301	11/11	0.94	0.10	35,37,46,46	0

6.5 Other polymers [i](#)

There are no such residues in this entry.