



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 08:15 AM UTC

PDB ID : 1GS5 / pdb_00001gs5
Title : N-acetyl-L-glutamate kinase from Escherichia coli complexed with its substrate
N-acetylglutamate and its substrate analog AMPPNP
Authors : Ramon-Maiques, S.; Marina, A.; Gil-Ortiz, F.; Fita, I.; Rubio, V.
Deposited on : 2001-12-28
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	11% 69% 25% 5%

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 2189 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACETYLGLUTAMATE KINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	258	1946	1231	334	371	10	0	10	0

- Molecule 2 is N-ACETYL-L-GLUTAMATE (CCD ID: NLG) (formula: C₇H₁₁NO₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	13	7	1	5	0	0

- Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (CCD ID: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 31	C 10	N 6	O 12	P 3	0	0

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Mg 1 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	198	Total O 198 198	0	1

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ACETYLGLUTAMATE KINASE



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	59.56Å 72.33Å 107.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.50 50.00 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-1.50) 99.9 (50.00-1.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.90 (at 1.44Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.209 , 0.213 0.198 , 0.222	Depositor DCC
R_{free} test set	2129 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	22.7	Xtriage
Anisotropy	0.202	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 44.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2189	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NLG, ANP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.23	4/2010 (0.2%)	1.95	65/2724 (2.4%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	131	LYS	N-CA	6.20	1.53	1.46
1	A	1[A]	MET	C-N	6.11	1.41	1.33
1	A	1[B]	MET	C-N	6.11	1.41	1.33
1	A	39	VAL	C-O	5.29	1.29	1.24

All (65) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	160	ASN	OD1-CG-ND2	12.06	134.66	122.60
1	A	36	ARG	O-C-N	10.47	130.62	121.30
1	A	139	ASN	CA-CB-CG	9.47	122.07	112.60
1	A	93	LYS	CA-C-O	-9.38	110.61	120.55
1	A	218	VAL	O-C-N	-9.27	112.88	121.87
1	A	233	ASP	O-C-N	8.98	133.69	123.19
1	A	234	ILE	O-C-N	-8.87	112.72	123.10
1	A	192	ARG	NE-CZ-NH1	-7.86	113.64	121.50
1	A	22	ARG	CD-NE-CZ	7.66	135.12	124.40
1	A	137	LEU	O-C-N	7.44	129.73	122.07
1	A	218	VAL	CA-C-O	7.41	128.66	120.95
1	A	210	ILE	O-C-N	7.32	131.87	122.94
1	A	198	ALA	O-C-N	7.32	129.88	122.12
1	A	242	GLN	O-C-N	6.90	132.20	122.41
1	A	233	ASP	CA-C-N	-6.81	114.27	123.12
1	A	233	ASP	C-N-CA	-6.81	114.27	123.12
1	A	211	THR	CA-C-O	6.77	129.09	121.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	249	GLY	CA-C-N	6.73	130.04	120.49
1	A	249	GLY	C-N-CA	6.73	130.04	120.49
1	A	96[A]	GLN	CA-C-O	6.61	128.12	120.32
1	A	96[B]	GLN	CA-C-O	6.61	128.12	120.32
1	A	210	ILE	CA-C-O	-6.55	112.99	120.65
1	A	214	MET	CA-C-N	6.37	129.47	120.42
1	A	214	MET	C-N-CA	6.37	129.47	120.42
1	A	3	ASN	N-CA-C	-6.23	101.73	109.64
1	A	141	TYR	O-C-N	6.16	129.96	123.06
1	A	91	TRP	O-C-N	-6.16	115.59	122.12
1	A	133	ILE	O-C-N	6.12	128.13	121.83
1	A	192	ARG	CD-NE-CZ	6.07	132.90	124.40
1	A	204	LEU	CA-C-O	6.06	126.95	120.10
1	A	214	MET	CA-C-O	6.06	126.94	119.79
1	A	256	ILE	CA-C-O	6.04	127.72	120.65
1	A	230	ARG	CA-CB-CG	6.04	126.18	114.10
1	A	173	GLY	O-C-N	-5.80	116.14	122.41
1	A	254	THR	CA-C-O	-5.76	113.98	120.32
1	A	167	ALA	O-C-N	5.74	128.20	122.12
1	A	93	LYS	O-C-N	5.74	128.20	122.12
1	A	41	VAL	O-C-N	-5.70	117.33	123.14
1	A	204	LEU	N-CA-C	5.66	118.22	111.71
1	A	256	ILE	O-C-N	-5.64	116.05	122.94
1	A	201	ALA	O-C-N	5.64	128.10	122.12
1	A	215	ILE	N-CA-C	-5.58	105.09	110.72
1	A	134	ASN	OD1-CG-ND2	5.57	128.17	122.60
1	A	87	THR	N-CA-C	-5.55	105.31	111.36
1	A	177	ILE	CA-C-N	5.54	129.79	122.42
1	A	177	ILE	C-N-CA	5.54	129.79	122.42
1	A	153	GLU	CA-C-O	5.51	125.95	119.28
1	A	116	ASP	CA-CB-CG	5.51	118.11	112.60
1	A	51	LEU	N-CA-C	-5.50	105.30	112.23
1	A	85	ASN	CA-C-O	-5.46	114.63	120.42
1	A	213	GLY	N-CA-C	-5.41	108.60	115.31
1	A	42	HIS	O-C-N	5.37	129.50	123.27
1	A	103	PHE	CA-C-O	5.34	128.06	121.60
1	A	198	ALA	CA-C-O	-5.33	114.90	120.55
1	A	192	ARG	NE-CZ-NH2	5.29	123.97	119.20
1	A	42	HIS	ND1-CG-CD2	5.29	111.39	106.10
1	A	48	VAL	CA-C-O	-5.26	115.60	121.17
1	A	93	LYS	N-CA-C	-5.24	105.56	111.28
1	A	141	TYR	CA-C-O	-5.19	115.13	121.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	166	THR	CA-CB-OG1	-5.18	101.83	109.60
1	A	202	GLU	CG-CD-OE1	5.18	130.31	118.40
1	A	222	LEU	N-CA-C	5.16	116.90	111.28
1	A	48	VAL	O-C-N	5.08	126.89	121.91
1	A	241	GLU	CG-CD-OE2	5.02	129.95	118.40
1	A	36	ARG	CD-NE-CZ	5.02	131.43	124.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1946	0	2017	62	0
2	A	13	0	9	0	0
3	A	31	0	13	3	0
4	A	1	0	0	0	0
5	A	198	0	0	6	0
All	All	2189	0	2039	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:52:MET:HA	1:A:52:MET:HE2	1.28	1.12
1:A:214:MET:HE1	1:A:217:LYS:HD2	1.44	1.00
1:A:138[B]:GLU:OE1	5:A:2118:HOH:O	1.81	0.96
1:A:51:LEU:HD21	1:A:75:ILE:HD11	1.49	0.94
1:A:51:LEU:HD21	1:A:75:ILE:CD1	2.04	0.86
1:A:205[B]:ILE:HD13	1:A:210:ILE:HB	1.57	0.84
1:A:75:ILE:O	5:A:2064:HOH:O	1.96	0.82
1:A:52:MET:HA	1:A:52:MET:CE	2.12	0.79
1:A:250:MET:HG2	1:A:251:PRO:HD2	1.65	0.79

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:51:LEU:CD2	1:A:75:ILE:HD11	2.13	0.79
1:A:185:ILE:HG12	1:A:214:MET:HE2	1.67	0.77
1:A:205[B]:ILE:HD12	1:A:215:ILE:HG12	1.68	0.75
1:A:1[A]:MET:HE1	1:A:137:LEU:C	2.13	0.74
1:A:1[A]:MET:HE1	1:A:137:LEU:O	1.89	0.73
1:A:138[B]:GLU:OE1	5:A:2117:HOH:O	2.08	0.70
1:A:214:MET:CE	1:A:217:LYS:HD2	2.23	0.68
1:A:52:MET:HE2	1:A:52:MET:CA	2.16	0.67
1:A:52:MET:CE	1:A:55:LEU:HD12	2.29	0.62
1:A:12:VAL:HG12	1:A:12:VAL:O	1.99	0.62
1:A:243:LEU:HB2	1:A:244:PRO:HD3	1.81	0.62
1:A:201:ALA:O	1:A:205[B]:ILE:HG12	2.00	0.61
1:A:181[B]:ASP:OD1	1:A:182:VAL:HG13	2.01	0.60
1:A:250:MET:SD	1:A:252:MET:SD	3.00	0.60
1:A:246:LEU:HD12	5:A:2188:HOH:O	2.02	0.59
1:A:181[A]:ASP:HB3	1:A:238:ARG:HH21	1.68	0.59
1:A:181[B]:ASP:HB2	1:A:238:ARG:HH21	1.65	0.59
1:A:204:LEU:CD2	1:A:209[B]:ILE:HD12	2.33	0.59
1:A:63:ASN:ND2	5:A:2046:HOH:O	2.30	0.57
1:A:205[B]:ILE:CD1	1:A:215:ILE:HG12	2.34	0.57
1:A:196:MET:HB3	1:A:256:ILE:HD13	1.86	0.56
1:A:204:LEU:HD23	1:A:209[B]:ILE:HD12	1.88	0.56
1:A:205[B]:ILE:HD11	1:A:210:ILE:CG2	2.35	0.56
1:A:181[A]:ASP:HB3	1:A:238:ARG:NH2	2.22	0.55
1:A:205[B]:ILE:CD1	1:A:210:ILE:CG2	2.83	0.55
1:A:51:LEU:CG	1:A:75:ILE:HD11	2.37	0.54
1:A:205[B]:ILE:HD11	1:A:210:ILE:HG21	1.90	0.54
1:A:222:LEU:O	1:A:226:ARG:HG2	2.09	0.52
1:A:201:ALA:O	1:A:205[A]:ILE:HG13	2.09	0.51
1:A:51:LEU:CD2	1:A:75:ILE:CD1	2.79	0.51
1:A:188:GLY:CA	1:A:209[A]:ILE:HD12	2.41	0.50
1:A:250:MET:HG2	1:A:251:PRO:CD	2.37	0.50
1:A:188:GLY:HA2	1:A:209[A]:ILE:HD12	1.93	0.49
1:A:62:LYS:NZ	1:A:118:GLU:HB3	2.28	0.49
1:A:181[B]:ASP:HB2	1:A:238:ARG:NH2	2.27	0.49
1:A:205[B]:ILE:HD13	1:A:210:ILE:CB	2.37	0.48
1:A:11:GLY:N	3:A:1260:ANP:HNB1	2.11	0.48
1:A:214:MET:HE3	1:A:217:LYS:HB3	1.96	0.48
1:A:52:MET:HE1	1:A:55:LEU:HD12	1.95	0.47
1:A:210:ILE:HG22	1:A:215:ILE:HG12	1.97	0.47
1:A:52:MET:HE2	1:A:55:LEU:HD12	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:51:LEU:HD21	1:A:75:ILE:HD12	1.93	0.46
3:A:1260:ANP:O1A	3:A:1260:ANP:H4'	2.14	0.46
1:A:196:MET:HE2	1:A:196:MET:HB2	1.81	0.46
1:A:226:ARG:HD2	5:A:2170:HOH:O	2.16	0.45
1:A:12:VAL:O	1:A:12:VAL:CG1	2.64	0.45
1:A:8:LYS:HA	1:A:41:VAL:O	2.17	0.45
1:A:184:GLY:HA2	1:A:234:ILE:HG22	1.98	0.44
1:A:203:GLN:O	1:A:207:GLN:HG3	2.17	0.44
1:A:37:PRO:HB2	1:A:137:LEU:HD22	1.99	0.44
1:A:11:GLY:HA3	1:A:181[A]:ASP:OD1	2.19	0.43
1:A:205[B]:ILE:CD1	1:A:210:ILE:HG22	2.50	0.42
1:A:11:GLY:H	3:A:1260:ANP:HNB1	1.67	0.41
1:A:185:ILE:CG1	1:A:214:MET:HE2	2.46	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/258 (103%)	256 (97%)	9 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	211/201 (105%)	204 (97%)	7 (3%)	33 8

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1[A]	MET
1	A	1[B]	MET
1	A	52	MET
1	A	60	LYS
1	A	103	PHE
1	A	212	ASP
1	A	250	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	ASN
1	A	63	ASN
1	A	121	HIS
1	A	248	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NLG	A	1259	-	12,12,12	4.29	7 (58%)	15,15,15	4.15	8 (53%)
3	ANP	A	1260	4	33,33,33	2.55	14 (42%)	45,52,52	2.09	19 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NLG	A	1259	-	-	0/13/13/13	-
3	ANP	A	1260	4	-	7/18/38/38	0/3/3/3

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1259	NLG	C8-C7	-8.98	1.31	1.50
2	A	1259	NLG	CA-C	-7.23	1.34	1.52
2	A	1259	NLG	C7-N2	-6.92	1.12	1.34
3	A	1260	ANP	C5'-C4'	-6.14	1.33	1.51
3	A	1260	ANP	C1'-N9	-4.88	1.32	1.46
3	A	1260	ANP	O2'-C2'	4.66	1.54	1.43
3	A	1260	ANP	PG-O3G	-4.43	1.45	1.56
3	A	1260	ANP	PB-O2B	-4.03	1.46	1.56
2	A	1259	NLG	CA-N2	3.82	1.53	1.45
3	A	1260	ANP	C4-N3	3.69	1.41	1.34
3	A	1260	ANP	C2-N1	3.44	1.40	1.33
2	A	1259	NLG	OXT-C	2.96	1.40	1.30
3	A	1260	ANP	C2'-C3'	2.63	1.60	1.53
3	A	1260	ANP	PA-O3A	2.61	1.62	1.59
3	A	1260	ANP	PB-N3B	-2.45	1.57	1.63
3	A	1260	ANP	C3'-C4'	2.41	1.59	1.53
3	A	1260	ANP	PG-O2G	-2.34	1.50	1.56
2	A	1259	NLG	O-C	2.33	1.29	1.22
3	A	1260	ANP	O5'-C5'	2.20	1.52	1.44
2	A	1259	NLG	CB-CG	2.18	1.59	1.52
3	A	1260	ANP	C5-N7	-2.15	1.35	1.39

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1259	NLG	OXT-C-O	-9.50	102.54	124.08
2	A	1259	NLG	C8-C7-N2	6.44	126.80	116.12
2	A	1259	NLG	CA-N2-C7	6.38	139.47	121.58
2	A	1259	NLG	O7-C7-N2	-5.78	111.77	121.98
2	A	1259	NLG	OXT-C-CA	4.85	129.94	113.51
3	A	1260	ANP	O4'-C4'-C5'	-4.30	95.55	109.33
3	A	1260	ANP	O1G-PG-N3B	4.24	118.01	111.77
3	A	1260	ANP	O3'-C3'-C4'	3.91	122.30	111.08
3	A	1260	ANP	O2B-PB-O1B	3.54	117.47	109.87
3	A	1260	ANP	O4'-C1'-N9	3.21	114.25	108.09
3	A	1260	ANP	O4'-C4'-C3'	-2.99	99.22	105.15
3	A	1260	ANP	O1B-PB-N3B	-2.94	107.45	111.77
3	A	1260	ANP	C4-N9-C8	-2.92	102.67	105.74
3	A	1260	ANP	O3A-PB-N3B	2.91	114.65	106.59
3	A	1260	ANP	C5-C6-N6	2.86	130.36	123.29
2	A	1259	NLG	OE1-CD-CG	2.73	131.75	123.09
3	A	1260	ANP	N9-C8-N7	2.73	117.80	113.94
2	A	1259	NLG	OE2-CD-CG	-2.73	105.39	114.00
3	A	1260	ANP	O3G-PG-O1G	-2.57	107.00	113.45
3	A	1260	ANP	C3'-C2'-C1'	2.44	106.08	101.46
2	A	1259	NLG	CB-CA-N2	2.41	115.69	110.91
3	A	1260	ANP	O5'-PA-O1A	-2.41	99.39	108.94
3	A	1260	ANP	N6-C6-N1	-2.37	113.11	118.38
3	A	1260	ANP	O2'-C2'-C3'	-2.36	104.26	111.82
3	A	1260	ANP	O3'-C3'-C2'	2.26	119.05	111.82
3	A	1260	ANP	O2A-PA-O3A	2.21	113.24	107.27
3	A	1260	ANP	C2'-C3'-C4'	-2.20	98.37	102.61

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1260	ANP	PB-N3B-PG-O1G
3	A	1260	ANP	C4'-C5'-O5'-PA
3	A	1260	ANP	O4'-C4'-C5'-O5'
3	A	1260	ANP	C3'-C4'-C5'-O5'
3	A	1260	ANP	PB-O3A-PA-O1A
3	A	1260	ANP	PA-O3A-PB-O2B
3	A	1260	ANP	PA-O3A-PB-O1B

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1260	ANP	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	258/258 (100%)	0.60	28 (10%)	10 11	9, 24, 46, 70	10 (3%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	258	ALA	4.5
1	A	14	LEU	3.9
1	A	75	ILE	3.5
1	A	53	LYS	3.2
1	A	1[A]	MET	3.2
1	A	12	VAL	2.9
1	A	56[A]	ASN	2.8
1	A	55	LEU	2.8
1	A	10	GLY	2.8
1	A	249	GLY	2.7
1	A	214	MET	2.6
1	A	13	LEU	2.5
1	A	237	TRP	2.5
1	A	251	PRO	2.5
1	A	57	LEU	2.4
1	A	59	VAL	2.4
1	A	228	LEU	2.4
1	A	52	MET	2.4
1	A	250	MET	2.4
1	A	54	GLY	2.3
1	A	213	GLY	2.3
1	A	182	VAL	2.2
1	A	2	MET	2.2
1	A	257	LEU	2.1
1	A	22	ARG	2.1
1	A	18	GLU	2.1
1	A	15	ASP	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	212	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	ANP	A	1260	31/31	0.84	0.12	23,27,48,55	21
4	MG	A	1261	1/1	0.85	0.29	38,38,38,38	0
2	NLG	A	1259	13/13	0.91	0.08	19,22,28,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ANP A 1260:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.