



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 09:27 PM UTC

PDB ID : 3GSI / pdb_00003gsi
Title : Crystal structure of D552A dimethylglycine oxidase mutant of *Arthrobacter globiformis* in complex with tetrahydrofolate
Authors : Tralau, T.; Lafite, P.; Levy, C.; Combe, J.P.; Scrutton, N.S.; Leys, D.
Deposited on : 2009-03-27
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	827	 92% 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	FAD	A	902	X	-	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7436 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called N,N-dimethylglycine oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	827	6270	3977	1082	1189	22	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	552	ALA	ASP	engineered mutation	UNP Q9AGP8

- Molecule 2 is (6S)-5,6,7,8-TETRAHYDROFOLATE (CCD ID: THG) (formula: C₁₉H₂₃N₇O₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	32	19	7	6	0	0

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		


- Molecule 4 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: N,N-dimethylglycine oxidase



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	71.22Å 224.55Å 119.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.79 – 2.00 39.79 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.8 (39.79-2.00) 96.9 (39.79-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.53 (at 2.00Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.150 , 0.202 0.151 , 0.201	Depositor DCC
R_{free} test set	3195 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	19.8	Xtriage
Anisotropy	0.107	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7436	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, MG, THG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	0/6418	0.91	0/8743

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6270	0	6134	45	2
2	A	32	0	23	0	0
3	A	1	0	0	0	0
4	A	53	0	29	8	0
5	A	1080	0	0	12	3
All	All	7436	0	6186	45	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:48:HIS:NE2	4:A:902:FAD:C8M	2.03	1.20
1:A:30:ASN:HB2	5:A:1510:HOH:O	1.60	0.98
1:A:48:HIS:NE2	4:A:902:FAD:HM81	1.75	0.97
1:A:48:HIS:NE2	4:A:902:FAD:HM83	1.76	0.95
1:A:335:PHE:CE2	5:A:1237:HOH:O	2.31	0.83
1:A:785:LYS:HD3	5:A:1649:HOH:O	1.85	0.75
1:A:755:ARG:HD3	5:A:1737:HOH:O	1.88	0.73
1:A:337:PHE:HE1	5:A:1237:HOH:O	1.75	0.67
1:A:135:ASN:HB2	5:A:1075:HOH:O	1.96	0.64
1:A:7:ILE:HD11	1:A:28:TRP:HE3	1.69	0.58
1:A:337:PHE:CE1	5:A:1237:HOH:O	2.52	0.58
1:A:48:HIS:NE2	4:A:902:FAD:C8	2.66	0.56
1:A:125:GLN:NE2	1:A:131:LEU:O	2.39	0.55
1:A:760:GLY:O	1:A:761:LYS:HB2	2.05	0.55
1:A:48:HIS:CE1	4:A:902:FAD:HM83	2.43	0.53
1:A:419:HIS:CG	1:A:420:PRO:HD2	2.44	0.52
1:A:106:LEU:HD22	1:A:118:LEU:HG	1.91	0.52
1:A:67:LYS:HD3	5:A:1787:HOH:O	2.09	0.51
1:A:613:ARG:HG3	1:A:655:LEU:HD21	1.95	0.49
1:A:409:GLN:HG3	5:A:1238:HOH:O	2.11	0.49
1:A:510:ARG:HD3	5:A:1604:HOH:O	2.13	0.48
1:A:651:TYR:HD2	1:A:690:PHE:CZ	2.30	0.48
1:A:258:LEU:HD21	1:A:311:ALA:HB1	1.95	0.48
1:A:337:PHE:HD2	1:A:410:ASN:HD22	1.60	0.47
1:A:471:GLU:H	1:A:471:GLU:CD	2.20	0.47
1:A:818:ASP:HB2	1:A:819:PRO:HA	1.96	0.47
1:A:223:LEU:HD23	1:A:278:PRO:HA	1.97	0.47
1:A:494:TRP:CZ3	1:A:495:LYS:HE2	2.49	0.47
1:A:545:HIS:HB2	5:A:1496:HOH:O	2.15	0.45
1:A:50:PRO:HD2	4:A:902:FAD:C5X	2.47	0.44
1:A:17:THR:HG21	1:A:154:VAL:HG13	2.00	0.43
1:A:651:TYR:CD2	1:A:690:PHE:CZ	3.07	0.43
1:A:345:LEU:HG	1:A:388:CYS:HB3	2.00	0.43
1:A:649:LEU:HD23	1:A:649:LEU:N	2.33	0.43
1:A:159:LYS:HE3	1:A:159:LYS:HB2	1.83	0.42
1:A:536:ALA:HB3	5:A:1588:HOH:O	2.20	0.42
1:A:48:HIS:CE1	4:A:902:FAD:C8M	2.92	0.42
1:A:208:ALA:HA	1:A:218:VAL:HB	2.02	0.42
1:A:28:TRP:CE2	1:A:377:THR:HG22	2.55	0.42
1:A:497:ARG:HD3	1:A:654:GLU:OE2	2.20	0.41
1:A:259:TYR:CZ	1:A:270:GLY:HA3	2.56	0.41
1:A:431:ARG:HB2	1:A:452:TRP:HA	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:50:PRO:HD2	4:A:902:FAD:N5	2.36	0.41
1:A:604:CYS:HA	1:A:659:LEU:O	2.21	0.40
1:A:69:THR:HA	1:A:366:ALA:HB3	2.03	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:594:GLN:OE1	1:A:594:GLN:OE1[3_656]	1.62	0.58
1:A:755:ARG:NH2	5:A:911:HOH:O[2_665]	1.96	0.24
5:A:1037:HOH:O	5:A:1037:HOH:O[4_566]	1.96	0.24
5:A:1447:HOH:O	5:A:1447:HOH:O[2_665]	1.96	0.24

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	825/827 (100%)	809 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	626/656 (95%)	618 (99%)	8 (1%)	61 68

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	329	ASP
1	A	380	ARG
1	A	390	ILE
1	A	471	GLU
1	A	593	VAL
1	A	619	VAL
1	A	746	LEU
1	A	795	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	ASN
1	A	240	ASN
1	A	410	ASN
1	A	422	GLN
1	A	564	GLN
1	A	568	ASN
1	A	583	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	FAD	A	902	-	58,58,58	3.30	21 (36%)	85,89,89	1.87	20 (23%)
2	THG	A	2887	-	32,34,34	4.88	19 (59%)	39,47,47	3.06	19 (48%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FAD	A	902	-	1/1/9/9	5/34/50/50	0/6/6/6
2	THG	A	2887	-	-	5/22/31/31	0/3/3/3

All (40) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2887	THG	C4A-C8A	15.27	1.64	1.38
4	A	902	FAD	O2'-C2'	-11.45	1.19	1.43
2	A	2887	THG	O4-C4	9.28	1.41	1.23
4	A	902	FAD	C4X-N5	8.53	1.49	1.30
2	A	2887	THG	C2-N3	8.39	1.58	1.37
4	A	902	FAD	O4-C4	7.40	1.37	1.23
4	A	902	FAD	C8A-N7A	7.21	1.45	1.31
2	A	2887	THG	C5'-C4'	6.97	1.51	1.39
4	A	902	FAD	C9-C8	6.39	1.48	1.39
2	A	2887	THG	OX1-C	6.19	1.40	1.22
2	A	2887	THG	C6'-C1'	6.10	1.48	1.39
2	A	2887	THG	C2'-C1'	5.97	1.48	1.39
4	A	902	FAD	C6-C5X	5.89	1.48	1.40
4	A	902	FAD	O2-C2	5.82	1.35	1.24
2	A	2887	THG	OE1-CD	5.78	1.41	1.22

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2887	THG	C11-N	5.67	1.47	1.34
2	A	2887	THG	C2-N2	5.31	1.46	1.34
4	A	902	FAD	C10-N10	4.73	1.47	1.37
2	A	2887	THG	C2'-C3'	4.71	1.46	1.38
4	A	902	FAD	C4A-N9A	4.69	1.47	1.37
4	A	902	FAD	C2A-N3A	4.51	1.41	1.33
2	A	2887	THG	C2-N1	4.26	1.43	1.33
2	A	2887	THG	C3'-C4'	3.82	1.45	1.39
4	A	902	FAD	C6A-N6A	3.71	1.43	1.34
2	A	2887	THG	OX2-C	3.50	1.41	1.30
4	A	902	FAD	P-O1P	3.48	1.62	1.50
4	A	902	FAD	C8A-N9A	-3.23	1.32	1.37
2	A	2887	THG	OE2-CD	3.22	1.41	1.30
4	A	902	FAD	P-O3P	3.15	1.62	1.59
4	A	902	FAD	PA-O1A	3.14	1.61	1.50
4	A	902	FAD	C4X-C10	-3.08	1.35	1.44
2	A	2887	THG	C4A-N5	2.94	1.44	1.37
4	A	902	FAD	C7M-C7	2.92	1.56	1.51
4	A	902	FAD	C10-N1	2.91	1.39	1.33
4	A	902	FAD	C5'-C4'	2.80	1.55	1.51
4	A	902	FAD	PA-O3P	2.70	1.62	1.59
2	A	2887	THG	CG-CD	2.50	1.56	1.50
4	A	902	FAD	C9-C9A	2.41	1.43	1.39
2	A	2887	THG	C4'-N10	2.30	1.45	1.38
2	A	2887	THG	CA-N	2.11	1.50	1.45

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2887	THG	C9-N10-C4'	12.70	154.32	122.00
4	A	902	FAD	N3A-C2A-N1A	-5.69	119.97	128.58
4	A	902	FAD	O2'-C2'-C3'	5.23	121.49	109.25
2	A	2887	THG	C9-C6-N5	4.63	118.76	108.90
4	A	902	FAD	N9A-C8A-N7A	-4.57	107.45	113.94
2	A	2887	THG	C2-N1-C8A	4.29	120.94	113.36
4	A	902	FAD	O4B-C4B-C5B	4.12	122.53	109.33
4	A	902	FAD	C4-N3-C2	-4.08	118.40	125.64
4	A	902	FAD	O3B-C3B-C4B	3.88	122.21	111.08
2	A	2887	THG	C4-C4A-N5	3.76	126.52	116.27
4	A	902	FAD	C4A-N9A-C8A	3.75	109.67	105.74
2	A	2887	THG	C5'-C4'-C3'	3.66	123.90	119.04
2	A	2887	THG	C6'-C1'-C2'	3.63	123.19	118.57

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2887	THG	C4A-C4-N3	3.46	121.63	112.13
4	A	902	FAD	C5B-C4B-C3B	3.16	126.59	115.21
2	A	2887	THG	OX2-C-OX1	-3.12	117.01	124.08
2	A	2887	THG	C2'-C3'-C4'	-3.04	116.80	120.30
2	A	2887	THG	OX1-C-CA	-3.03	112.47	122.26
2	A	2887	THG	OE1-CD-CG	-3.00	113.58	123.09
2	A	2887	THG	C5'-C6'-C1'	-2.82	117.78	120.80
2	A	2887	THG	OE2-CD-OE1	-2.82	116.09	123.33
2	A	2887	THG	O4-C4-C4A	-2.73	120.68	127.26
4	A	902	FAD	C4X-C10-N10	2.71	120.36	116.48
2	A	2887	THG	C2-N3-C4	-2.69	120.23	125.11
4	A	902	FAD	C10-C4X-N5	-2.59	119.51	124.81
4	A	902	FAD	O4-C4-C4X	-2.58	119.72	126.53
4	A	902	FAD	C4'-C3'-C2'	-2.56	109.32	113.57
4	A	902	FAD	C5A-N7A-C8A	2.52	107.41	103.45
4	A	902	FAD	C4-C4X-C10	2.50	121.22	116.93
4	A	902	FAD	C4X-C4-N3	2.49	119.60	113.25
4	A	902	FAD	C2B-C3B-C4B	-2.43	97.91	102.61
4	A	902	FAD	C2A-N3A-C4A	2.41	117.71	111.83
4	A	902	FAD	C5A-C4A-N3A	-2.36	123.47	126.72
2	A	2887	THG	C6-C7-N8	-2.35	106.33	110.60
2	A	2887	THG	C6-C9-N10	2.32	117.07	110.24
2	A	2887	THG	C5'-C4'-N10	-2.31	115.74	120.96
2	A	2887	THG	N3-C2-N1	-2.18	119.34	123.32
4	A	902	FAD	C4A-C5A-N7A	-2.12	108.16	110.58
4	A	902	FAD	O4'-C4'-C3'	2.09	114.15	109.25

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	902	FAD	C4B

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	2887	THG	N5-C6-C9-N10
4	A	902	FAD	N10-C1'-C2'-O2'
4	A	902	FAD	C3B-C4B-C5B-O5B
4	A	902	FAD	O4B-C4B-C5B-O5B
2	A	2887	THG	C3'-C4'-N10-C9
2	A	2887	THG	C7-C6-C9-N10
2	A	2887	THG	C5'-C4'-N10-C9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	902	FAD	C1'-C2'-C3'-O3'
4	A	902	FAD	C1'-C2'-C3'-C4'
2	A	2887	THG	OE1-CD-CG-CB


There are no ring outliers.

1 monomer is involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	902	FAD	8	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand FAD A 902



Ligand THG A 2887



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	827/827 (100%)	-0.69	0 100 100	10, 18, 32, 50	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	THG	A	2887	32/32	0.95	0.06	18,27,35,39	0
3	MG	A	901	1/1	0.97	0.04	19,19,19,19	0
4	FAD	A	902	53/53	0.98	0.04	9,14,17,18	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around THG A 2887:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD A 902:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.