



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:44 PM UTC

PDB ID : 4GS7 / pdb_00004gs7
Title : Structure of the Interleukin-15 quaternary complex
Authors : Ring, A.M.; Ozkan, E.; Feng, D.; Garcia, K.C.
Deposited on : 2012-08-27
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1596 (2.36-2.36)
Clashscore	190562	1663 (2.36-2.36)
Ramachandran outliers	187476	1646 (2.36-2.36)
Sidechain outliers	187428	1646 (2.36-2.36)
RSRZ outliers	180081	1598 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	116	
2	B	217	
3	C	203	
4	D	69	
5	E	2	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	F	2	50% 50%

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 5053 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-15.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	114	Total	C	N	O	S	0	0	0
			887	560	141	179	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	expression tag	UNP P40933
A	0	GLY	-	expression tag	UNP P40933

- Molecule 2 is a protein called Interleukin-2 receptor subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	196	Total	C	N	O	S	0	2	0
			1640	1051	289	290	10			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	ALA	-	expression tag	UNP P14784
B	-1	ASP	-	expression tag	UNP P14784
B	0	PRO	-	expression tag	UNP P14784
B	3	GLN	ASN	engineered mutation	UNP P14784
B	17	GLN	ASN	engineered mutation	UNP P14784
B	45	GLN	ASN	engineered mutation	UNP P14784

- Molecule 3 is a protein called Cytokine receptor common subunit gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	196	Total	C	N	O	S	0	1	0
			1659	1058	294	299	8			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	30	ALA	-	expression tag	UNP P31785
C	31	ASP	-	expression tag	UNP P31785
C	32	PRO	-	expression tag	UNP P31785
C	53	GLN	ASN	engineered mutation	UNP P31785

- Molecule 4 is a protein called Interleukin-15 receptor subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	67	Total	C	N	O	S	0	0	0
			521	328	92	97	4			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-1	MET	-	initiating methionine	UNP Q13261

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	2	Total	C	N	O		0	0	0
			28	16	2	10				
5	F	2	Total	C	N	O		0	0	0
			28	16	2	10				

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2^-$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	C	1	Total	C	N	O	0	0
			14	8	1	5		
8	C	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	50	Total	O	0	0
			50	50		
9	B	103	Total	O	0	0
			103	103		
9	C	81	Total	O	0	0
			81	81		
9	D	20	Total	O	0	0
			20	20		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Interleukin-15




- Molecule 2: Interleukin-2 receptor subunit beta



- Molecule 3: Cytokine receptor common subunit gamma



- Molecule 4: Interleukin-15 receptor subunit alpha



- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  50% 50%

MAG1
MAG2

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	70.95Å 74.61Å 129.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.23 – 2.35 40.23 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.2 (40.23-2.35) 99.4 (40.23-2.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.33 (at 2.34Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, R_{free}	0.182 , 0.227 0.185 , 0.230	Depositor DCC
R_{free} test set	1456 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	34.4	Xtriage
Anisotropy	0.186	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 50.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.018 for k,h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5053	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, ACT, NAG, MLY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/854	0.64	0/1159
2	B	0.27	0/1648	0.66	2/2253 (0.1%)
3	C	0.25	0/1685	0.67	0/2301
4	D	0.26	0/535	0.64	0/730
All	All	0.26	0/4722	0.66	2/6443 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	99	LYS	CA-C-N	5.40	124.85	119.24
2	B	99	LYS	C-N-CA	5.40	124.85	119.24

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	887	0	867	5	0
2	B	1640	0	1593	6	0
3	C	1659	0	1551	11	0
4	D	521	0	507	3	0
5	E	28	0	25	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	F	28	0	25	0	0
6	B	4	0	6	1	0
7	C	4	0	3	0	0
8	C	28	0	26	0	0
9	A	50	0	0	2	0
9	B	103	0	0	2	0
9	C	81	0	0	2	0
9	D	20	0	0	0	0
All	All	5053	0	4603	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:145:THR:O	9:B:482:HOH:O	2.01	0.78
3:C:136:GLU:OE1	3:C:155:ARG:NH1	2.26	0.68
2:B:91:ARG:HH21	2:B:93:MET:HE1	1.61	0.66
3:C:54:SER:HB3	3:C:89:ILE:HD11	1.85	0.57
3:C:144:SER:OG	3:C:145:GLU:N	2.36	0.56
3:C:67:TYR:OH	3:C:99[A]:GLU:OE2	2.26	0.49
4:D:4:PRO:O	4:D:56:TRP:NE1	2.42	0.49
4:D:17:LYS:HG3	4:D:25:GLU:OE2	2.13	0.49
2:B:159:THR:HG23	9:B:417:HOH:O	2.14	0.48
3:C:196:ARG:NH1	9:C:478:HOH:O	2.03	0.46
2:B:40:ASP:HA	6:B:301:EDO:H21	1.98	0.46
2:B:139:LEU:HD23	2:B:186:PRO:HA	1.97	0.45
1:A:19:MET:HE1	1:A:54:SER:HB2	1.99	0.45
1:A:87:GLU:HB2	1:A:90:GLU:HG3	1.99	0.44
3:C:138:LEU:HB3	3:C:222:ILE:HD13	1.98	0.44
2:B:146:LEU:HB3	2:B:179:GLU:HB2	2.00	0.43
3:C:196:ARG:HB2	3:C:226:SER:HA	2.00	0.43
1:A:96:ILE:HG12	9:A:204:HOH:O	2.19	0.42
3:C:33:LEU:O	3:C:118:ARG:NH1	2.49	0.42
3:C:200:ARG:NH1	9:C:470:HOH:O	2.26	0.42
1:A:26:TYR:OH	4:D:35:ARG:O	2.37	0.41
1:A:-1:MET:N	9:A:248:HOH:O	2.45	0.41
3:C:151:ASN:OD1	3:C:185:LYS:HG2	2.22	0.40
3:C:206:ASN:HA	3:C:207:PRO:HA	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	108/116 (93%)	105 (97%)	3 (3%)	0	100	100
2	B	190/217 (88%)	181 (95%)	9 (5%)	0	100	100
3	C	192/203 (95%)	181 (94%)	11 (6%)	0	100	100
4	D	65/69 (94%)	63 (97%)	2 (3%)	0	100	100
All	All	555/605 (92%)	530 (96%)	25 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	98/104 (94%)	98 (100%)	0	100	100
2	B	178/190 (94%)	177 (99%)	1 (1%)	78	87
3	C	183/192 (95%)	183 (100%)	0	100	100
4	D	59/62 (95%)	57 (97%)	2 (3%)	32	43
All	All	518/548 (94%)	515 (99%)	3 (1%)	78	87

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	171	THR
4	D	13	ASP
4	D	17	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	101	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

11 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MLY	C	125	3	9,10,11	0.61	0	6,11,13	0.67	0
1	MLY	A	11	1	9,10,11	0.51	0	6,11,13	0.81	0
1	MLY	A	10	1	9,10,11	0.55	0	6,11,13	0.77	0
2	MLY	B	185	2	9,10,11	0.58	0	6,11,13	0.82	0
3	MLY	C	76	3	9,10,11	0.50	0	6,11,13	0.95	0
2	MLY	B	163	2	9,10,11	0.56	0	6,11,13	0.89	0
1	MLY	A	94	1	9,10,11	0.52	0	6,11,13	0.87	0
2	MLY	B	161	2	9,10,11	0.46	0	6,11,13	0.79	0
3	MLY	C	70	3	9,10,11	0.48	0	6,11,13	0.98	0
2	MLY	B	71	2	9,10,11	0.53	0	6,11,13	0.86	0
1	MLY	A	97	1	9,10,11	0.52	0	6,11,13	0.85	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MLY	C	125	3	-	3/8/9/11	-
1	MLY	A	11	1	-	0/8/9/11	-
1	MLY	A	10	1	-	1/8/9/11	-
2	MLY	B	185	2	-	0/8/9/11	-
3	MLY	C	76	3	-	0/8/9/11	-
2	MLY	B	163	2	-	0/8/9/11	-
1	MLY	A	94	1	-	2/8/9/11	-
2	MLY	B	161	2	-	0/8/9/11	-
3	MLY	C	70	3	-	3/8/9/11	-
2	MLY	B	71	2	-	1/8/9/11	-
1	MLY	A	97	1	-	5/8/9/11	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	94	MLY	C-CA-CB-CG
1	A	97	MLY	C-CA-CB-CG
3	C	70	MLY	O-C-CA-CB
3	C	125	MLY	O-C-CA-CB
1	A	97	MLY	CD-CE-NZ-CH1
1	A	97	MLY	CD-CE-NZ-CH2
3	C	70	MLY	CG-CD-CE-NZ
3	C	125	MLY	CA-CB-CG-CD
1	A	97	MLY	CE-CD-CG-CB
1	A	97	MLY	CA-CB-CG-CD
1	A	94	MLY	CE-CD-CG-CB
3	C	70	MLY	CE-CD-CG-CB
1	A	10	MLY	CD-CE-NZ-CH2
2	B	71	MLY	C-CA-CB-CG
3	C	125	MLY	CD-CE-NZ-CH1

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	NAG	E	1	5,2	14,14,15	0.54	0	17,19,21	0.67	0
5	NAG	E	2	5	14,14,15	0.48	0	17,19,21	0.80	1 (5%)
5	NAG	F	1	5,3	14,14,15	0.53	0	17,19,21	0.78	0
5	NAG	F	2	5	14,14,15	0.45	0	17,19,21	1.26	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	E	1	5,2	-	0/6/23/26	0/1/1/1
5	NAG	E	2	5	-	0/6/23/26	0/1/1/1
5	NAG	F	1	5,3	-	0/6/23/26	0/1/1/1
5	NAG	F	2	5	-	3/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	F	2	NAG	C1-O5-C5	3.67	117.10	112.19
5	E	2	NAG	C2-N2-C7	-2.01	120.21	122.90

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	F	2	NAG	C8-C7-N2-C2
5	F	2	NAG	O7-C7-N2-C2

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	F	2	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	ACT	C	301	-	3,3,3	0.80	0	3,3,3	1.49	0
8	NAG	C	305	3	14,14,15	0.48	0	17,19,21	0.68	1 (5%)
8	NAG	C	302	3	14,14,15	0.45	0	17,19,21	0.92	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	EDO	B	301	-	3,3,3	0.43	0	2,2,2	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	NAG	C	305	3	-	2/6/23/26	0/1/1/1
8	NAG	C	302	3	-	0/6/23/26	0/1/1/1
6	EDO	B	301	-	-	0/1/1/1	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	C	305	NAG	C1-O5-C5	2.01	114.88	112.19

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	C	305	NAG	C8-C7-N2-C2
8	C	305	NAG	O7-C7-N2-C2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	301	EDO	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	110/116 (94%)	-0.05	7 (6%) 25 28	20, 34, 78, 122	0
2	B	192/217 (88%)	-0.21	4 (2%) 63 68	14, 31, 61, 86	2 (1%)
3	C	193/203 (95%)	0.13	13 (6%) 24 27	21, 37, 81, 123	1 (0%)
4	D	67/69 (97%)	0.41	5 (7%) 20 23	32, 53, 87, 108	0
All	All	562/605 (92%)	0.01	29 (5%) 33 38	14, 35, 78, 123	3 (0%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	79	ASN	4.7
3	C	156	PHE	4.4
1	A	-1	MET	3.8
3	C	72	SER	3.7
2	B	31	THR	3.6
3	C	171	TRP	3.6
3	C	58	PRO	3.6
1	A	77	ASN	3.5
4	D	53	VAL	3.3
3	C	32	PRO	3.2
3	C	145	GLU	3.1
2	B	6	SER	3.0
2	B	24	GLN	3.0
3	C	54	SER	3.0
1	A	80	VAL	2.9
4	D	50	ALA	2.9
1	A	0	GLY	2.7
3	C	155	ARG	2.7
3	C	59	GLN	2.6
2	B	88	VAL	2.5
3	C	146	SER	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	71	ASN	2.4
4	D	51	THR	2.3
4	D	1	ILE	2.3
4	D	67	PRO	2.3
3	C	55	SER	2.3
1	A	78	GLY	2.2
3	C	117	ARG	2.1
1	A	20	HIS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MLY	C	70	11/12	0.86	0.16	34,52,63,63	0
3	MLY	C	76	11/12	0.89	0.14	46,50,73,74	0
1	MLY	A	94	11/12	0.91	0.12	30,36,49,49	0
1	MLY	A	97	11/12	0.93	0.10	26,33,42,45	0
2	MLY	B	185	11/12	0.94	0.10	25,31,52,54	0
2	MLY	B	71	11/12	0.95	0.07	17,27,39,41	0
3	MLY	C	125	11/12	0.95	0.09	21,28,49,55	0
2	MLY	B	161	11/12	0.96	0.09	22,32,55,56	0
2	MLY	B	163	11/12	0.96	0.07	22,26,38,41	0
1	MLY	A	10	11/12	0.96	0.10	26,32,57,59	0
1	MLY	A	11	11/12	0.97	0.06	23,29,35,36	0


6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	F	2	14/15	0.68	0.16	63,74,84,87	0
5	NAG	E	2	14/15	0.90	0.08	35,39,51,52	0
5	NAG	F	1	14/15	0.91	0.10	30,41,52,62	0
5	NAG	E	1	14/15	0.97	0.05	18,27,32,34	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
8	NAG	C	305	14/15	0.66	0.16	51,69,77,79	0
7	ACT	C	301	4/4	0.69	0.20	59,62,66,68	0
8	NAG	C	302	14/15	0.70	0.13	44,60,71,74	0
6	EDO	B	301	4/4	0.87	0.21	50,52,53,54	0

6.5 Other polymers [i](#)

There are no such residues in this entry.