



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:24 PM UTC

PDB ID : 5GS7 / pdb_00005gs7
Title : Crystal structure determination of Cys2Ala mutant of Bile Salt Hydrolase from *Enterococcus faecalis*
Authors : Ramasamy, S.; Deepak, C.; Varshney, N.K.; Suresh, C.G.
Deposited on : 2016-08-14
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	323	4% 85% 13% .
1	B	323	4% 87% 10% .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5351 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Choloylglycine hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	323	Total	C	N	O	S	0	0	0
			2595	1657	420	510	8			
1	B	323	Total	C	N	O	S	0	0	0
			2595	1658	420	509	8			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	ALA	CYS	engineered mutation	UNP Q8KUA7
B	2	ALA	CYS	engineered mutation	UNP Q8KUA7


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	82	Total	O	0	0
			82	82		
2	B	79	Total	O	0	0
			79	79		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Choloylglycine hydrolase



• Molecule 1: Choloylglycine hydrolase



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	91.19Å 91.19Å 156.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.29 – 1.50 45.29 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.8 (45.29-1.50) 99.8 (45.29-1.50)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.21 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.209 , 0.237 (Not available) , 0.234	Depositor DCC
R_{free} test set	5262 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	18.1	Xtriage
Anisotropy	0.145	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 22.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5351	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.68	26/2653 (1.0%)	1.35	8/3597 (0.2%)
1	B	1.56	16/2654 (0.6%)	1.32	14/3599 (0.4%)
All	All	1.62	42/5307 (0.8%)	1.34	22/7196 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (42) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	9	SER	CA-C	14.37	1.59	1.52
1	A	170	ASN	CA-C	9.66	1.59	1.53
1	A	102	ILE	CA-CB	7.68	1.58	1.54
1	A	102	ILE	C-O	-7.63	1.18	1.24
1	B	102	ILE	C-O	-7.58	1.18	1.24
1	B	170	ASN	CA-C	7.32	1.57	1.53
1	B	9	SER	CA-C	6.93	1.55	1.52
1	B	165	VAL	C-O	-6.78	1.16	1.24
1	A	80	PHE	C-O	6.62	1.30	1.23
1	B	16	ARG	CZ-NH2	-6.14	1.25	1.33
1	A	141	LEU	CA-CB	6.10	1.62	1.53
1	B	217	ASP	CA-C	6.00	1.61	1.53
1	A	24	TYR	CA-C	5.92	1.60	1.53
1	B	54	ILE	CA-CB	5.86	1.61	1.54
1	B	182	ASN	CA-C	5.79	1.60	1.52
1	A	130	SER	CA-C	-5.77	1.45	1.52
1	B	149	ILE	N-CA	-5.71	1.39	1.46
1	A	4	ALA	CA-C	-5.62	1.46	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	53	GLY	N-CA	5.60	1.51	1.45
1	B	215	PRO	C-O	5.56	1.29	1.23
1	A	275	SER	CA-C	-5.55	1.46	1.52
1	A	54	ILE	N-CA	-5.53	1.39	1.46
1	A	220	SER	C-O	-5.49	1.17	1.24
1	B	8	VAL	C-O	-5.48	1.18	1.24
1	A	180	ASN	N-CA	5.48	1.53	1.46
1	A	14	PHE	C-O	5.46	1.30	1.23
1	B	53	GLY	N-CA	5.42	1.51	1.45
1	A	126	ASN	N-CA	-5.41	1.39	1.46
1	A	20	TYR	CA-C	-5.40	1.46	1.52
1	A	74	GLY	N-CA	-5.32	1.37	1.45
1	A	318	GLN	N-CA	-5.31	1.39	1.46
1	A	95	ASN	N-CA	-5.31	1.39	1.46
1	A	296	ALA	CA-CB	-5.30	1.44	1.53
1	A	106	LEU	CA-C	5.30	1.60	1.52
1	B	177	GLN	C-O	5.28	1.30	1.24
1	A	85	ASP	C-O	-5.27	1.18	1.23
1	A	54	ILE	CA-CB	5.26	1.60	1.54
1	A	306	LYS	N-CA	-5.24	1.39	1.45
1	B	169	THR	C-O	5.12	1.27	1.23
1	A	256	GLN	CA-C	-5.12	1.46	1.52
1	B	175	ASP	CA-CB	5.09	1.61	1.53
1	B	178	LEU	N-CA	-5.02	1.40	1.46

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	9	SER	CA-C-O	7.60	124.10	119.77
1	A	122	LEU	N-CA-C	7.13	120.83	110.28
1	B	102	ILE	O-C-N	6.64	124.67	120.42
1	A	220	SER	N-CA-C	6.37	117.88	111.07
1	B	122	LEU	N-CA-C	6.34	119.67	110.28
1	B	139	HIS	CA-C-O	-6.04	114.58	121.58
1	B	58	ILE	N-CA-C	-5.69	107.74	113.20
1	B	182	ASN	O-C-N	5.67	128.86	122.22
1	B	306	LYS	N-CA-C	5.55	117.50	110.33
1	B	224	PHE	CA-C-O	5.50	126.78	120.90
1	B	24	TYR	N-CA-C	5.49	117.45	109.71
1	A	33	ARG	N-CA-C	5.40	117.17	111.28
1	A	222	SER	N-CA-C	5.39	117.16	111.28
1	A	58	ILE	N-CA-C	-5.33	108.09	113.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	104	TRP	O-C-N	-5.30	116.61	122.07
1	B	108	GLN	N-CA-C	5.25	121.42	114.12
1	A	102	ILE	O-C-N	5.24	123.77	120.42
1	B	145	LYS	CB-CG-CD	5.17	123.20	111.30
1	A	306	LYS	N-CA-C	5.11	117.12	110.43
1	B	320	ILE	N-CA-C	5.08	116.60	108.87
1	B	80	PHE	CA-CB-CG	5.03	118.83	113.80
1	B	212	ILE	O-C-N	5.01	127.65	122.79

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	138	LEU	Peptide
1	A	210	GLY	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2595	0	2503	19	1
1	B	2595	0	2504	17	1
2	A	82	0	0	0	0
2	B	79	0	0	1	0
All	All	5351	0	5007	32	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:24:TYR:O	1:A:58:ILE:HD11	1.89	0.72
1:B:24:TYR:O	1:B:58:ILE:HD11	1.89	0.72
1:B:139:HIS:HD2	1:B:152:GLU:OE2	1.77	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139:HIS:HD2	1:A:152:GLU:OE2	1.79	0.66
1:B:24:TYR:O	1:B:58:ILE:CD1	2.43	0.66
1:A:24:TYR:O	1:A:58:ILE:CD1	2.44	0.65
1:A:298:ASP:H	1:B:324:ASN:HD21	1.45	0.65
1:B:36:LYS:NZ	2:B:401:HOH:O	2.28	0.65
1:A:320:ILE:HD12	1:B:241:TYR:CE1	2.33	0.64
1:A:324:ASN:O	1:A:324:ASN:CA	2.45	0.64
1:A:58:ILE:HG22	1:A:58:ILE:O	1.99	0.62
1:B:58:ILE:HG22	1:B:58:ILE:O	2.00	0.61
1:B:139:HIS:HE1	1:B:169:THR:O	1.84	0.61
1:A:139:HIS:HE1	1:A:169:THR:O	1.87	0.57
1:B:2:ALA:N	1:B:17:ASN:O	2.40	0.55
1:A:58:ILE:O	1:A:58:ILE:CG2	2.56	0.54
1:A:24:TYR:C	1:A:58:ILE:HD11	2.35	0.51
1:B:58:ILE:O	1:B:58:ILE:CG2	2.59	0.50
1:A:85:ASP:HB2	1:A:155:LYS:HD3	1.94	0.49
1:B:104:TRP:CH2	1:B:118:LEU:HG	2.48	0.49
1:A:237:ARG:CG	1:A:237:ARG:HH11	2.28	0.47
1:A:36:LYS:HG3	1:A:46:ASP:OD1	2.15	0.47
1:A:298:ASP:H	1:B:324:ASN:ND2	2.12	0.46
1:B:24:TYR:C	1:B:58:ILE:HD11	2.40	0.45
1:A:241:TYR:CE1	1:B:320:ILE:HD12	2.52	0.45
1:B:139:HIS:CE1	1:B:169:THR:O	2.68	0.43
1:A:56:ALA:HB1	1:A:58:ILE:HD13	2.00	0.43
1:B:2:ALA:HB1	1:B:170:ASN:OD1	2.19	0.43
1:A:104:TRP:CH2	1:A:118:LEU:HG	2.54	0.43
1:B:36:LYS:HG3	1:B:46:ASP:OD1	2.20	0.41
1:A:109:CYS:SG	1:A:118:LEU:HD22	2.61	0.41
1:A:178:LEU:HA	1:A:178:LEU:HD23	1.86	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:176:TYR:CE1	1:B:176:TYR:CE1[5_555]	1.95	0.25
1:A:176:TYR:CE1	1:A:176:TYR:CE1[5_555]	2.19	0.01

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	321/323 (99%)	312 (97%)	9 (3%)	0	100	100
1	B	321/323 (99%)	308 (96%)	13 (4%)	0	100	100
All	All	642/646 (99%)	620 (97%)	22 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	289/289 (100%)	284 (98%)	5 (2%)	53	26
1	B	289/289 (100%)	283 (98%)	6 (2%)	47	19
All	All	578/578 (100%)	567 (98%)	11 (2%)	50	22

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	117	LYS
1	A	139	HIS
1	A	155	LYS
1	A	237	ARG
1	A	239	SER
1	B	88	LYS
1	B	117	LYS
1	B	139	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	258	LYS
1	B	307	ASP
1	B	324	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	ASN
1	A	123	ASN
1	A	139	HIS
1	A	195	ASN
1	A	293	GLN
1	A	300	ASN
1	B	79	ASN
1	B	123	ASN
1	B	139	HIS
1	B	324	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	323/323 (100%)	0.19	13 (4%)	42	46	11, 17, 32, 59	0
1	B	323/323 (100%)	0.20	12 (3%)	45	49	10, 17, 32, 56	0
All	All	646/646 (100%)	0.20	25 (3%)	43	47	10, 17, 32, 59	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	24	TYR	7.0
1	B	58	ILE	5.0
1	A	24	TYR	4.5
1	A	235	VAL	4.0
1	A	236	SER	3.6
1	A	237	ARG	3.3
1	B	155	LYS	3.1
1	A	265	ASP	3.0
1	A	58	ILE	3.0
1	A	238	SER	3.0
1	B	2	ALA	2.9
1	A	266	GLU	2.9
1	B	266	GLU	2.8
1	B	307	ASP	2.6
1	A	176	TYR	2.5
1	B	176	TYR	2.5
1	B	265	ASP	2.4
1	A	23	SER	2.4
1	B	22	ILE	2.4
1	A	22	ILE	2.3
1	B	57	GLY	2.1
1	A	91	GLU	2.0
1	B	60	ASP	2.0
1	A	155	LYS	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	235	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.