



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 10:53 AM UTC

PDB ID : 5GSN / pdb_00005gsn
Title : Tmm in complex with methimazole
Authors : Zhang, Y.Z.; Li, C.Y.
Deposited on : 2016-08-16
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	453	
1	B	453	
1	C	453	
1	D	453	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MMZ	A	503	-	X	-	-
4	MMZ	C	503	-	X	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 15548 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Flavin-containing monooxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	445	Total	C	N	O	S	0	1	0
			3582	2287	607	671	17			
1	B	445	Total	C	N	O	S	0	4	0
			3607	2303	613	674	17			
1	C	445	Total	C	N	O	S	0	0	0
			3573	2282	605	669	17			
1	D	445	Total	C	N	O	S	0	1	0
			3582	2287	606	672	17			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	153	ALA	GLU	engineered mutation	UNP A3SLM3
A	154	ALA	ASP	engineered mutation	UNP A3SLM3
A	207	SER	TYR	engineered mutation	UNP A3SLM3
A	448	HIS	-	expression tag	UNP A3SLM3
A	449	HIS	-	expression tag	UNP A3SLM3
A	450	HIS	-	expression tag	UNP A3SLM3
A	451	HIS	-	expression tag	UNP A3SLM3
A	452	HIS	-	expression tag	UNP A3SLM3
A	453	HIS	-	expression tag	UNP A3SLM3
B	153	ALA	GLU	engineered mutation	UNP A3SLM3
B	154	ALA	ASP	engineered mutation	UNP A3SLM3
B	207	SER	TYR	engineered mutation	UNP A3SLM3
B	448	HIS	-	expression tag	UNP A3SLM3
B	449	HIS	-	expression tag	UNP A3SLM3
B	450	HIS	-	expression tag	UNP A3SLM3
B	451	HIS	-	expression tag	UNP A3SLM3
B	452	HIS	-	expression tag	UNP A3SLM3
B	453	HIS	-	expression tag	UNP A3SLM3
C	153	ALA	GLU	engineered mutation	UNP A3SLM3
C	154	ALA	ASP	engineered mutation	UNP A3SLM3
C	207	SER	TYR	engineered mutation	UNP A3SLM3

Continued on next page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	448	HIS	-	expression tag	UNP A3SLM3
C	449	HIS	-	expression tag	UNP A3SLM3
C	450	HIS	-	expression tag	UNP A3SLM3
C	451	HIS	-	expression tag	UNP A3SLM3
C	452	HIS	-	expression tag	UNP A3SLM3
C	453	HIS	-	expression tag	UNP A3SLM3
D	153	ALA	GLU	engineered mutation	UNP A3SLM3
D	154	ALA	ASP	engineered mutation	UNP A3SLM3
D	207	SER	TYR	engineered mutation	UNP A3SLM3
D	448	HIS	-	expression tag	UNP A3SLM3
D	449	HIS	-	expression tag	UNP A3SLM3
D	450	HIS	-	expression tag	UNP A3SLM3
D	451	HIS	-	expression tag	UNP A3SLM3
D	452	HIS	-	expression tag	UNP A3SLM3
D	453	HIS	-	expression tag	UNP A3SLM3

- # NAP

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 48	C 21	N 7	O 17	P 3	0	0
2	B	1	Total 48	C 21	N 7	O 17	P 3	0	0
2	C	1	Total 48	C 21	N 7	O 17	P 3	0	0



WORLD WIDE
PDB
PROTEIN DATA BANK



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	S	0	0
			7	4	2	1		
4	C	1	Total	C	N	S	0	0
			7	4	2	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	186	Total	O	0	0
			186	186		
5	B	202	Total	O	0	0
			202	202		
5	C	206	Total	O	0	0
			206	206		
5	D	192	Total	O	0	0
			192	192		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Flavin-containing monooxygenase




• Molecule 1: Flavin-containing monooxygenase




• Molecule 1: Flavin-containing monooxygenase





• Molecule 1: Flavin-containing monooxygenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	60.85Å 207.58Å 72.52Å 90.00° 90.31° 90.00°	Depositor
Resolution (Å)	46.74 – 2.20 46.74 – 2.20	Depositor EDS
% Data completeness (in resolution range)	89.1 (46.74-2.20) 91.5 (46.74-2.20)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.73 (at 2.20Å)	Xtriage
Refinement program	PHENIX 1.6.4_486	Depositor
R, R_{free}	0.203 , 0.274 0.203 , 0.269	Depositor DCC
R_{free} test set	4242 reflections (4.69%)	wwPDB-VP
Wilson B-factor (Å ²)	28.9	Xtriage
Anisotropy	0.540	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 14.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.338 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	15548	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 33.25 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.2477e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MMZ, NAP, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/3691	0.80	1/5010 (0.0%)
1	B	0.51	0/3716	0.83	1/5044 (0.0%)
1	C	0.53	2/3682 (0.1%)	0.84	5/4998 (0.1%)
1	D	0.49	0/3691	0.81	1/5010 (0.0%)
All	All	0.51	2/14780 (0.0%)	0.82	8/20062 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	231	ALA	C-N	6.23	1.41	1.33
1	C	232	PRO	N-CD	5.12	1.54	1.47

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	231	ALA	CA-C-N	-7.14	112.96	120.03
1	C	231	ALA	C-N-CA	-7.14	112.96	120.03
1	C	84	TYR	N-CA-C	-5.37	99.36	110.80
1	C	167	THR	CA-C-N	5.33	125.23	119.85
1	C	167	THR	C-N-CA	5.33	125.23	119.85

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3582	0	3363	75	0
1	B	3607	0	3391	83	0
1	C	3573	0	3356	141	0
1	D	3582	0	3361	123	0
2	A	48	0	24	10	0
2	B	48	0	25	6	0
2	C	48	0	24	12	0
2	D	48	0	24	3	0
3	A	53	0	31	4	0
3	B	53	0	31	2	0
3	C	53	0	31	7	0
3	D	53	0	31	3	0
4	A	7	0	6	2	0
4	C	7	0	6	11	0
5	A	186	0	0	6	1
5	B	202	0	0	13	1
5	C	206	0	0	11	2
5	D	192	0	0	4	0
All	All	15548	0	13704	436	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 436 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:323:ASN:HD22	1:D:323:ASN:H	1.01	1.01
1:A:87:GLU:OE2	1:A:93:GLN:NE2	1.98	0.97
1:D:323:ASN:HD22	1:D:323:ASN:N	1.64	0.95
1:C:421:ILE:HG13	1:C:422:THR:HG23	1.52	0.92
1:C:45:GLY:HA2	3:C:502:FAD:O3B	1.70	0.91

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:772:HOH:O	5:C:779:HOH:O[1_556]	1.94	0.26
5:B:794:HOH:O	5:C:803:HOH:O[1_455]	2.19	0.01

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	444/453 (98%)	417 (94%)	25 (6%)	2 (0%)	24	27
1	B	447/453 (99%)	423 (95%)	24 (5%)	0	100	100
1	C	443/453 (98%)	411 (93%)	27 (6%)	5 (1%)	11	10
1	D	444/453 (98%)	405 (91%)	35 (8%)	4 (1%)	14	14
All	All	1778/1812 (98%)	1656 (93%)	111 (6%)	11 (1%)	21	23

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	300	LYS
1	C	27	GLN
1	D	95	ALA
1	A	84	TYR
1	C	235	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	366/373 (98%)	352 (96%)	14 (4%)	29	40
1	B	369/373 (99%)	358 (97%)	11 (3%)	36	49
1	C	365/373 (98%)	353 (97%)	12 (3%)	33	45
1	D	366/373 (98%)	346 (94%)	20 (6%)	19	24
All	All	1466/1492 (98%)	1409 (96%)	57 (4%)	29	39

5 of 57 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	65	SER
1	D	413	ARG
1	C	425	MET
1	D	323	ASN
1	D	230	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	256	HIS
1	C	429	HIS
1	D	415	ASN
1	C	323	ASN
1	C	430	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAP	A	501	-	50,52,52	3.18	17 (34%)	71,80,80	2.23	16 (22%)
2	NAP	B	501	-	50,52,52	3.04	16 (32%)	71,80,80	2.37	19 (26%)
2	NAP	D	501	-	50,52,52	3.12	18 (36%)	71,80,80	2.14	17 (23%)
3	FAD	C	502	-	58,58,58	1.47	10 (17%)	85,89,89	1.51	15 (17%)
2	NAP	C	501	-	50,52,52	3.10	20 (40%)	71,80,80	2.30	19 (26%)
3	FAD	D	502	-	58,58,58	1.47	10 (17%)	85,89,89	1.52	15 (17%)
3	FAD	B	502	-	58,58,58	1.47	10 (17%)	85,89,89	1.53	16 (18%)
4	MMZ	C	503	-	7,7,7	7.56	4 (57%)	9,9,9	11.42	5 (55%)
4	MMZ	A	503	-	7,7,7	7.79	5 (71%)	9,9,9	11.73	5 (55%)
3	FAD	A	502	-	58,58,58	1.49	12 (20%)	85,89,89	1.54	15 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	501	-	-	13/35/67/67	0/5/5/5
2	NAP	B	501	-	-	7/35/67/67	0/5/5/5
2	NAP	D	501	-	-	6/35/67/67	0/5/5/5
3	FAD	C	502	-	-	10/34/50/50	0/6/6/6
2	NAP	C	501	-	-	10/35/67/67	0/5/5/5
3	FAD	D	502	-	-	3/34/50/50	0/6/6/6
3	FAD	B	502	-	-	4/34/50/50	0/6/6/6
4	MMZ	C	503	-	-	-	0/1/1/1
4	MMZ	A	503	-	-	-	0/1/1/1
3	FAD	A	502	-	-	4/34/50/50	0/6/6/6

The worst 5 of 122 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	503	MMZ	C2-N1	16.15	1.45	1.35
4	A	503	MMZ	C2-N1	15.56	1.45	1.35
4	A	503	MMZ	C2-N3	12.22	1.46	1.36
4	C	503	MMZ	C2-N3	10.31	1.44	1.36
2	D	501	NAP	C2N-N1N	9.47	1.45	1.35

The worst 5 of 142 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	503	MMZ	S2-C2-N3	-28.79	109.44	127.14
4	A	503	MMZ	S2-C2-N3	-23.45	112.72	127.14
4	A	503	MMZ	C1A-N1-C2	-17.30	103.43	110.78
4	A	503	MMZ	C3A-N3-C2	-13.78	105.63	109.60
4	A	503	MMZ	S2-C2-N1	-11.81	111.56	127.26

There are no chirality outliers.

5 of 57 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	NAP	O4D-C1D-N1N-C2N
2	A	501	NAP	O4D-C1D-N1N-C6N
2	A	501	NAP	C2D-C1D-N1N-C2N
2	A	501	NAP	C2D-C1D-N1N-C6N
2	A	501	NAP	C2N-C3N-C7N-N7N


There are no ring outliers.


10 monomers are involved in 48 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	NAP	10	0
2	B	501	NAP	6	0
2	D	501	NAP	3	0
3	C	502	FAD	7	0
2	C	501	NAP	12	0
3	D	502	FAD	3	0
3	B	502	FAD	2	0
4	C	503	MMZ	11	0
4	A	503	MMZ	2	0
3	A	502	FAD	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	445/453 (98%)	-1.33	0 100 100	13, 30, 49, 66	1 (0%)
1	B	445/453 (98%)	-1.32	0 100 100	12, 30, 50, 65	4 (0%)
1	C	445/453 (98%)	-1.26	0 100 100	16, 32, 63, 86	0
1	D	445/453 (98%)	-1.27	0 100 100	14, 31, 65, 86	1 (0%)
All	All	1780/1812 (98%)	-1.30	0 100 100	12, 30, 57, 86	6 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAP	A	501	48/48	0.99	0.04	25,34,53,61	0
2	NAP	B	501	48/48	0.99	0.03	21,32,44,49	0
2	NAP	C	501	48/48	0.99	0.03	25,35,53,56	0
2	NAP	D	501	48/48	0.99	0.03	23,33,46,52	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	FAD	A	502	53/53	0.99	0.03	13,22,36,37	0
3	FAD	B	502	53/53	0.99	0.03	14,22,30,37	0
3	FAD	C	502	53/53	0.99	0.03	14,24,40,41	0
3	FAD	D	502	53/53	0.99	0.03	15,26,33,35	0
4	MMZ	A	503	7/7	0.99	0.04	35,41,46,59	0
4	MMZ	C	503	7/7	0.99	0.04	37,41,48,64	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around NAP A 501:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAP B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAP C 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAP D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD A 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around FAD B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD C 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.