



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 09:13 PM UTC

PDB ID : 4GT0 / pdb_00004gt0
Title : Structure of dengue virus serotype 1 sE containing stem to residue 421
Authors : Klein, D.E.; Choi, J.L.; Harrison, S.C.
Deposited on : 2012-08-28
Resolution : 2.57 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4770 (2.60-2.56)
Clashscore	190562	5124 (2.60-2.56)
Ramachandran outliers	187476	5046 (2.60-2.56)
Sidechain outliers	187428	5046 (2.60-2.56)
RSRZ outliers	180081	4770 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	437	
1	B	437	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6226 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Envelope protein E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	390	Total	C	N	O	S	0	4	0
			3007	1902	507	578	20			
1	B	390	Total	C	N	O	S	0	1	0
			2963	1872	498	573	20			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-15	GLY	-	expression tag	UNP P17763
A	-14	HIS	-	expression tag	UNP P17763
A	-13	HIS	-	expression tag	UNP P17763
A	-12	HIS	-	expression tag	UNP P17763
A	-11	HIS	-	expression tag	UNP P17763
A	-10	HIS	-	expression tag	UNP P17763
A	-9	HIS	-	expression tag	UNP P17763
A	-8	HIS	-	expression tag	UNP P17763
A	-7	HIS	-	expression tag	UNP P17763
A	-6	GLY	-	expression tag	UNP P17763
A	-5	SER	-	expression tag	UNP P17763
A	-4	SER	-	expression tag	UNP P17763
A	-3	THR	-	expression tag	UNP P17763
A	-2	SER	-	expression tag	UNP P17763
A	-1	ASN	-	expression tag	UNP P17763
A	0	GLY	-	expression tag	UNP P17763
A	101	HIS	TRP	engineered mutation	UNP P17763
A	161	ILE	THR	conflict	UNP P17763
A	202	LYS	GLU	conflict	UNP P17763
A	203	GLU	LYS	conflict	UNP P17763
B	-15	GLY	-	expression tag	UNP P17763
B	-14	HIS	-	expression tag	UNP P17763
B	-13	HIS	-	expression tag	UNP P17763
B	-12	HIS	-	expression tag	UNP P17763
B	-11	HIS	-	expression tag	UNP P17763

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-10	HIS	-	expression tag	UNP P17763
B	-9	HIS	-	expression tag	UNP P17763
B	-8	HIS	-	expression tag	UNP P17763
B	-7	HIS	-	expression tag	UNP P17763
B	-6	GLY	-	expression tag	UNP P17763
B	-5	SER	-	expression tag	UNP P17763
B	-4	SER	-	expression tag	UNP P17763
B	-3	THR	-	expression tag	UNP P17763
B	-2	SER	-	expression tag	UNP P17763
B	-1	ASN	-	expression tag	UNP P17763
B	0	GLY	-	expression tag	UNP P17763
B	101	HIS	TRP	engineered mutation	UNP P17763
B	161	ILE	THR	conflict	UNP P17763
B	202	LYS	GLU	conflict	UNP P17763
B	203	GLU	LYS	conflict	UNP P17763

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is CADMIUM ION (CCD ID: CD) (formula: Cd).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Cd	0	0
			2	2		
3	B	2	Total	Cd	0	0
			2	2		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	118	Total	O	0	0
			118	118		
5	B	105	Total	O	0	0
			105	105		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Envelope protein E

Chain A: 



• Molecule 1: Envelope protein E

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	77.89Å 77.89Å 292.29Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.48 – 2.57 28.48 – 2.57	Depositor EDS
% Data completeness (in resolution range)	99.6 (28.48-2.57) 99.7 (28.48-2.57)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.57 (at 2.58Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, R_{free}	0.177 , 0.210 0.162 , 0.193	Depositor DCC
R_{free} test set	1563 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	40.3	Xtriage
Anisotropy	0.015	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 21.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.40$, $\langle L^2 \rangle = 0.23$	Xtriage
Estimated twinning fraction	0.499 for h,-h-k,-l	Xtriage
Reported twinning fraction	0.500 for h,-h-k,-l	Depositor
Outliers	0 of 31764 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6226	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CD, CL, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/3071	0.62	0/4156
1	B	0.26	0/3016	0.61	0/4087
All	All	0.26	0/6087	0.61	0/8243

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3007	0	3050	34	0
1	B	2963	0	2965	46	2
2	A	14	0	13	5	0
2	B	14	0	13	3	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	1	0	0	1	0
5	A	118	0	0	6	0
5	B	105	0	0	11	1
All	All	6226	0	6041	81	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 7.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:67:ASN:HD21	2:B:501:NAG:C1	1.25	1.50
1:A:67:ASN:HD21	2:A:501:NAG:C1	1.35	1.36
1:B:399:GLY:N	5:B:653:HOH:O	2.09	0.86
1:B:67:ASN:ND2	2:B:501:NAG:O5	2.10	0.84
1:B:76:THR:OG1	1:B:77:GLN:OE1	1.98	0.80
1:B:399:GLY:O	5:B:621:HOH:O	2.05	0.75
1:A:67:ASN:HD22	2:A:501:NAG:C1	1.99	0.73
1:B:67:ASN:HD22	2:B:501:NAG:C1	1.97	0.71
1:B:399:GLY:O	5:B:653:HOH:O	2.08	0.71
4:A:504:CL:CL	5:A:608:HOH:O	2.45	0.70
1:B:254:GLY:O	1:B:256:GLN:NE2	2.24	0.70
1:B:29:SER:O	5:B:626:HOH:O	2.08	0.69
1:B:102:GLY:N	1:B:105:CYS:O	2.27	0.68
1:B:174:GLN:NE2	5:B:693:HOH:O	2.28	0.67
1:B:229:GLU:OE2	5:B:677:HOH:O	2.11	0.67
1:A:163:THR:O	5:A:662:HOH:O	2.13	0.67
1:A:329:THR:O	5:A:652:HOH:O	2.12	0.66
1:B:368:GLU:OE2	5:B:676:HOH:O	2.15	0.63
1:A:67:ASN:ND2	2:A:501:NAG:C2	2.63	0.62
1:A:67:ASN:HD21	2:A:501:NAG:C2	2.07	0.62
1:A:342:GLU:OE1	1:A:343:LYS:N	2.38	0.57
1:B:65:ILE:HG23	1:B:117:ALA:HB1	1.87	0.56
1:B:27:HIS:CD2	1:B:280:ALA:HB1	2.40	0.56
1:A:326:TYR:OH	1:A:330:ASP:OD1	2.13	0.55
1:A:67:ASN:ND2	2:A:501:NAG:N2	2.56	0.54
1:B:241:LYS:NZ	5:B:703:HOH:O	2.40	0.53
1:A:201:MET:HE1	1:A:260:MET:HB2	1.90	0.53
1:B:400:LYS:HA	5:B:621:HOH:O	2.09	0.52
1:A:36:LYS:NZ	1:A:293:THR:OG1	2.20	0.52
1:A:235:ASP:N	1:A:235:ASP:OD1	2.44	0.51
1:B:75:PRO:C	1:B:76:THR:HG23	2.36	0.51
1:B:330:ASP:OD1	1:B:330:ASP:N	2.45	0.49
1:A:61:ILE:HD13	1:A:201:MET:HE2	1.95	0.49
1:A:305:SER:HB3	1:A:385:LYS:HG3	1.94	0.49
1:B:21:VAL:HB	1:B:287:LEU:HB2	1.95	0.49
1:B:2:ARG:NH2	5:B:689:HOH:O	2.46	0.48
1:B:61:ILE:O	1:B:256:GLN:N	2.46	0.48
1:B:125:LEU:C	1:B:125:LEU:HD12	2.38	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:86:GLN:N	1:B:86:GLN:CD	2.72	0.47
1:B:332:PRO:HA	1:B:358:VAL:O	2.14	0.47
1:A:201:MET:HE1	1:A:260:MET:CB	2.43	0.47
1:A:326:TYR:CZ	1:A:328:GLY:HA3	2.50	0.47
1:B:75:PRO:O	1:B:76:THR:OG1	2.33	0.47
1:A:161:ILE:HD12	1:A:161:ILE:N	2.30	0.47
1:A:124:LYS:O	5:A:613:HOH:O	2.20	0.47
1:A:125:LEU:HD12	1:A:125:LEU:C	2.40	0.47
1:A:181:LEU:C	1:A:181:LEU:HD13	2.40	0.47
1:A:247:LYS:NZ	5:A:680:HOH:O	2.47	0.46
1:B:108:PHE:C	1:B:108:PHE:CD1	2.93	0.46
1:A:305:SER:HB3	1:A:385:LYS:CG	2.46	0.46
1:A:332:PRO:HA	1:A:358:VAL:O	2.17	0.44
1:B:67:ASN:OD1	1:B:118:LYS:HE2	2.18	0.44
1:B:198:LEU:HD13	1:B:270:ILE:HD11	1.99	0.44
1:B:201:MET:HE1	1:B:261:HIS:CD2	2.53	0.44
1:B:52:ASN:N	1:B:53:PRO:CD	2.80	0.44
1:B:108:PHE:O	1:B:108:PHE:CG	2.71	0.44
1:B:331:ALA:N	5:B:625:HOH:O	2.41	0.44
1:A:176:THR:O	1:A:177:ASP:HB2	2.18	0.43
1:B:198:LEU:HD12	1:B:198:LEU:C	2.43	0.43
1:A:32:THR:HA	1:A:41:LEU:O	2.18	0.43
1:A:181:LEU:C	1:A:181:LEU:CD1	2.92	0.43
1:A:8:ASN:HA	1:A:11:PHE:CE2	2.54	0.43
1:A:181:LEU:HD13	1:A:182:THR:N	2.33	0.43
1:B:102:GLY:O	1:B:103:ASN:C	2.61	0.43
1:A:21:VAL:HB	1:A:287:LEU:HB2	2.00	0.43
1:A:35:ALA:O	1:A:37:ASP:N	2.52	0.43
1:B:65:ILE:HG12	1:B:252:VAL:CG1	2.48	0.43
1:A:330:ASP:OD1	1:A:330:ASP:N	2.51	0.43
1:A:321[A]:LEU:HD23	1:A:322:VAL:N	2.35	0.42
1:B:170:THR:HA	1:B:183:LEU:O	2.20	0.42
1:A:12:VAL:HG23	1:A:12:VAL:O	2.19	0.42
1:B:205:SER:OG	1:B:272:THR:CG2	2.68	0.41
1:B:39:PRO:O	1:B:178:TYR:OH	2.25	0.41
1:B:41:LEU:HD13	1:B:143:VAL:HG22	2.01	0.41
1:B:199:LEU:C	1:B:199:LEU:HD23	2.45	0.41
1:A:101:HIS:HB2	1:B:309:GLU:O	2.20	0.41
1:B:83:VAL:O	1:B:86:GLN:HG2	2.21	0.41
1:B:299:TYR:CD2	1:B:357:ILE:HD12	2.56	0.40
1:B:75:PRO:O	1:B:76:THR:CB	2.70	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:361:LYS:C	1:B:363:LYS:H	2.29	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:225:SER:N	5:B:621:HOH:O[3_445]	1.99	0.21
1:B:13:GLU:OE2	1:B:16:SER:OG[2_455]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	390/437 (89%)	375 (96%)	13 (3%)	2 (0%)	24	44
1	B	385/437 (88%)	366 (95%)	18 (5%)	1 (0%)	36	56
All	All	775/874 (89%)	741 (96%)	31 (4%)	3 (0%)	30	49

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	36	LYS
1	B	291	LYS
1	A	202	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	338/370 (91%)	315 (93%)	23 (7%)	14	31
1	B	328/370 (89%)	308 (94%)	20 (6%)	17	35
All	All	666/740 (90%)	623 (94%)	43 (6%)	15	33

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	ARG
1	A	15	LEU
1	A	36	LYS
1	A	51	THR
1	A	69	THR
1	A	81	THR
1	A	84	GLU
1	A	95	THR
1	A	135	LEU
1	A	144	HIS
1	A	177	ASP
1	A	181	LEU
1	A	197	VAL
1	A	204	LYS
1	A	225	SER
1	A	230	THR
1	A	272	THR
1	A	308	LEU
1	A	325	LYS
1	A	342	GLU
1	A	380	VAL
1	A	387	LEU
1	A	389	LEU
1	B	12	VAL
1	B	51	THR
1	B	79	GLU
1	B	81	THR
1	B	86	GLN
1	B	91	VAL
1	B	108	PHE
1	B	124	LYS
1	B	135	LEU
1	B	172	GLU
1	B	177	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	197	VAL
1	B	198	LEU
1	B	207	LEU
1	B	230	THR
1	B	272	THR
1	B	284	LYS
1	B	298	SER
1	B	321	LEU
1	B	363	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	67	ASN
1	A	316	GLN
1	A	355	ASN
1	A	366	ASN
1	B	67	ASN
1	B	86	GLN
1	B	256	GLN
1	B	317	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 5 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	501	1	14,14,15	0.56	0	17,19,21	0.75	0
2	NAG	A	501	1	14,14,15	0.52	0	17,19,21	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	501	1	-	3/6/23/26	0/1/1/1
2	NAG	A	501	1	-	3/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	501	NAG	C1-C2-N2-C7
2	B	501	NAG	C8-C7-N2-C2
2	B	501	NAG	O7-C7-N2-C2
2	A	501	NAG	C8-C7-N2-C2
2	A	501	NAG	O7-C7-N2-C2
2	A	501	NAG	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	NAG	3	0
2	A	501	NAG	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	390/437 (89%)	-1.72	0 100 100	4, 12, 29, 48	4 (1%)
1	B	390/437 (89%)	-1.63	0 100 100	2, 15, 32, 45	1 (0%)
All	All	780/874 (89%)	-1.68	0 100 100	2, 13, 31, 48	5 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	A	501	14/15	0.98	0.04	21,38,46,53	0
2	NAG	B	501	14/15	0.98	0.04	46,53,58,58	0
3	CD	A	502	1/1	1.00	0.03	53,53,53,53	0
3	CD	A	503	1/1	1.00	0.02	52,52,52,52	0
3	CD	B	502	1/1	1.00	0.01	65,65,65,65	0
3	CD	B	503	1/1	1.00	0.03	62,62,62,62	0
4	CL	A	504	1/1	1.00	0.01	9,9,9,9	0

6.5 Other polymers [i](#)

There are no such residues in this entry.