



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:40 AM UTC

PDB ID : 7GTS / pdb_00007gts
Title : PanDDA Analysis group deposition – Crystal structure of PTP1B in complex with FMOPL000604a
Authors : Mehlman, T.; Ginn, H.M.; Keedy, D.A.
Deposited on : 2024-01-03
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	321	38% 76% 11% 12%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	WZY	A	402	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4889 atoms, of which 2329 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein phosphatase non-receptor type 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	284	4621	1471	2304	399	432	15	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	32	SER	CYS	engineered mutation	UNP P18031
A	92	VAL	CYS	engineered mutation	UNP P18031


- Molecule 2 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
2	A	1	20	4	12	1	3	0	0

- Molecule 3 is N-(4-methoxyphenyl)glycinamide (CCD ID: WZY) (formula: $C_9H_{12}N_2O_2$)

(labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	0	0
			26	9	13	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	219	Total	O	0	4
			222	222		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein phosphatase non-receptor type 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	89.85Å 89.85Å 106.38Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	62.81 – 1.80 62.81 – 1.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (62.81-1.80) 100.0 (62.81-1.80)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.38 (at 1.81Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.203 , 0.236 0.276 , 0.295	Depositor DCC
R_{free} test set	1817 reflections (3.94%)	wwPDB-VP
Wilson B-factor (Å ²)	34.8	Xtriage
Anisotropy	0.003	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 50.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.025 for -h,-k,l	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	4889	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, WZY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.21	3/2369 (0.1%)	1.29	3/3192 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	211	VAL	C-O	8.98	1.33	1.24
1	A	212	VAL	C-O	5.44	1.29	1.24
1	A	120	LYS	C-O	5.41	1.31	1.24

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	213	VAL	CA-CB-CG1	5.96	120.53	110.40
1	A	72	ILE	CA-CB-CG1	5.65	120.01	110.40
1	A	106	GLY	CA-C-O	-5.16	117.08	121.57

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2317	2304	2296	22	2
2	A	8	12	12	0	2

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	13	13	0	0	0
4	A	222	0	0	10	0
All	All	2560	2329	2308	22	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:119:LEU:O	4:A:503:HOH:O	1.75	1.05
1:A:73:LYS:HE2	4:A:505:HOH:O	1.65	0.95
1:A:36:LYS:HD2	4:A:520:HOH:O	1.85	0.75
1:A:73:LYS:CE	4:A:505:HOH:O	2.26	0.74
1:A:73:LYS:HE3	4:A:536:HOH:O	1.89	0.73
1:A:279:LYS:O	1:A:284:ASP:HB3	1.97	0.64
1:A:227:LEU:HB2	1:A:253:MET:HE1	1.78	0.64
1:A:73:LYS:NZ	4:A:505:HOH:O	2.28	0.62
1:A:128:LYS:NZ	4:A:504:HOH:O	2.13	0.61
1:A:92:VAL:HG12	1:A:135:PHE:CE1	2.37	0.60
1:A:152:TYR:C	1:A:177:THR:HG1	2.13	0.56
1:A:152:TYR:C	1:A:177:THR:OG1	2.54	0.50
1:A:73:LYS:CE	4:A:536:HOH:O	2.59	0.44
1:A:76:GLU:OE2	1:A:238:ARG:NH1	2.51	0.44
1:A:21:GLN:HG2	4:A:575:HOH:O	2.17	0.44
1:A:148:ASP:OD1	1:A:150:LYS:HE3	2.18	0.43
1:A:196:PHE:O	1:A:200:GLU:HG3	2.18	0.43
1:A:92:VAL:CG1	1:A:135:PHE:CE1	3.02	0.43
1:A:152:TYR:CZ	1:A:190:SER:HB2	2.55	0.42
1:A:39:LYS:HE3	4:A:583:HOH:O	2.21	0.41
1:A:45:ARG:NH2	1:A:121:CYS:HA	2.36	0.41
1:A:111:ASN:HB3	1:A:221:ARG:HH12	1.86	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:130:GLU:OE2	2:A:401:TRS:O2[5_544]	1.73	0.47
1:A:130:GLU:OE2	2:A:401:TRS:HO2[5_544]	1.32	0.28

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	282/321 (88%)	270 (96%)	11 (4%)	1 (0%)	30	19

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	261	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	257/294 (87%)	244 (95%)	13 (5%)	21	9

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	ASP
1	A	15	SER
1	A	72	ILE
1	A	121	CYS
1	A	139	ASN
1	A	146	SER
1	A	150	LYS
1	A	187	SER
1	A	213	VAL
1	A	221	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	236	ASP
1	A	237	LYS
1	A	253	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	139	ASN
1	A	193	ASN
1	A	208	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	WZY	A	402	-	12,13,13	0.93	1 (8%)	14,16,16	1.57	2 (14%)
2	TRS	A	401	-	7,7,7	0.77	0	9,9,9	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	WZY	A	402	-	-	4/8/8/8	0/1/1/1
2	TRS	A	401	-	-	6/9/9/9	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	WZY	C6-N1	2.01	1.45	1.41

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	WZY	C9-C8-N1	4.60	122.17	114.32
3	A	402	WZY	C5-C4-C3	2.22	122.27	119.73

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	TRS	C1-C-C2-O2
2	A	401	TRS	N-C-C2-O2
3	A	402	WZY	C9-C8-N1-C6
3	A	402	WZY	O2-C8-N1-C6
3	A	402	WZY	C4-C3-O1-C7
3	A	402	WZY	C2-C3-O1-C7
2	A	401	TRS	C3-C-C1-O1
2	A	401	TRS	N-C-C1-O1
2	A	401	TRS	C3-C-C2-O2
2	A	401	TRS	C2-C-C1-O1


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	TRS	0	2

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	284/321 (88%)	4.37	123 (43%) 0 0	10, 32, 68, 112	58 (20%)

All (123) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	280	PHE	25.3
1	A	241	PRO	22.6
1	A	179	TRP	20.4
1	A	281	ILE	20.2
1	A	184	VAL	19.9
1	A	196	PHE	19.5
1	A	180	PRO	19.2
1	A	243	SER	19.0
1	A	153	TYR	18.3
1	A	269	PHE	18.1
1	A	185	PRO	18.1
1	A	189	ALA	17.9
1	A	194	PHE	17.7
1	A	152	TYR	17.6
1	A	234	LEU	17.4
1	A	195	LEU	17.3
1	A	178	THR	17.2
1	A	233	LEU	17.2
1	A	182	PHE	16.9
1	A	117	GLY	16.9
1	A	155	VAL	16.9
1	A	188	PRO	16.7
1	A	242	SER	16.6
1	A	187	SER	16.5
1	A	145	ILE	16.4
1	A	240	ASP	16.2
1	A	149	ILE	16.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	118	SER	16.1
1	A	183	GLY	16.0
1	A	236	ASP	15.7
1	A	151	SER	15.7
1	A	265	ASP	15.6
1	A	174	PHE	15.4
1	A	181	ASP	15.3
1	A	146	SER	15.2
1	A	144	LEU	15.2
1	A	191	PHE	15.1
1	A	119	LEU	14.9
1	A	244	VAL	14.9
1	A	116	LYS	14.9
1	A	282	MET	14.5
1	A	154	THR	14.4
1	A	197	LYS	14.2
1	A	150	LYS	13.9
1	A	238	ARG	13.8
1	A	235	MET	13.8
1	A	115	GLU	13.3
1	A	221	ARG	13.2
1	A	170	GLU	12.7
1	A	237	LYS	12.6
1	A	199	ARG	12.4
1	A	129	GLU	12.3
1	A	186	GLU	12.1
1	A	157	GLN	12.1
1	A	105	ARG	12.0
1	A	262	GLN	11.1
1	A	90	ASN	11.0
1	A	239	LYS	10.4
1	A	1	MET	7.9
1	A	5	LYS	6.7
1	A	284	ASP	5.8
1	A	279	LYS	4.8
1	A	25	HIS	4.6
1	A	139	ASN	4.5
1	A	63	ASP	4.3
1	A	17	ALA	4.3
1	A	2	GLU	4.3
1	A	248	LYS	4.3
1	A	7	PHE	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	14	GLY	4.1
1	A	61	GLN	3.9
1	A	12	LYS	3.9
1	A	166	GLN	3.8
1	A	283	GLY	3.7
1	A	3	MET	3.6
1	A	264	ALA	3.6
1	A	78	GLN	3.5
1	A	47	ARG	3.5
1	A	276	GLU	3.4
1	A	4	GLU	3.4
1	A	41	LYS	3.4
1	A	10	ILE	3.3
1	A	246	ILE	3.3
1	A	193	ASN	3.2
1	A	33	ARG	3.1
1	A	226	CYS	3.1
1	A	165	THR	3.1
1	A	9	GLN	3.0
1	A	163	LEU	2.9
1	A	135	PHE	2.9
1	A	272	LEU	2.9
1	A	161	GLU	2.9
1	A	21	GLN	2.8
1	A	138	THR	2.8
1	A	278	ALA	2.8
1	A	275	ILE	2.8
1	A	211	VAL	2.8
1	A	208	HIS	2.7
1	A	24	ARG	2.7
1	A	8	GLU	2.6
1	A	207	GLU	2.6
1	A	13	SER	2.6
1	A	253	MET	2.5
1	A	23	ILE	2.5
1	A	206	PRO	2.5
1	A	204	LEU	2.4
1	A	134	ILE	2.4
1	A	18	ALA	2.4
1	A	22	ASP	2.4
1	A	147	GLU	2.4
1	A	73	LYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	16	TRP	2.3
1	A	42	ASN	2.3
1	A	19	ILE	2.2
1	A	177	THR	2.2
1	A	148	ASP	2.2
1	A	32	SER	2.1
1	A	251	LEU	2.1
1	A	137	ASP	2.1
1	A	58	LYS	2.1
1	A	37	LEU	2.1
1	A	40	ASN	2.0
1	A	38	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	WZY	A	402	13/13	0.56	0.56	34,40,41,45	26
2	TRS	A	401	8/8	0.77	0.20	32,48,58,69	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around WZY A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.