



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:50 AM UTC

PDB ID : 9GTY / pdb_00009gty
Title : RIPK1 in complex with AZ"320
Authors : Petersen, J.
Deposited on : 2024-09-18
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3689 (2.16-2.12)
Clashscore	190562	3812 (2.16-2.12)
Ramachandran outliers	187476	3773 (2.16-2.12)
Sidechain outliers	187428	3772 (2.16-2.12)
RSRZ outliers	180081	3691 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	313	12% 68% 11% • 19%
1	B	313	19% 67% 15% • 17%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4225 atoms, of which 12 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Receptor-interacting serine/threonine-protein kinase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	253	Total	C	N	O	S	0	0	0
			2006	1292	333	365	16			
1	B	259	Total	C	N	O	S	0	0	0
			2043	1314	339	374	16			

There are 46 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	MET	-	initiating methionine	UNP Q13546
A	-17	GLY	-	expression tag	UNP Q13546
A	-16	HIS	-	expression tag	UNP Q13546
A	-15	HIS	-	expression tag	UNP Q13546
A	-14	HIS	-	expression tag	UNP Q13546
A	-13	HIS	-	expression tag	UNP Q13546
A	-12	HIS	-	expression tag	UNP Q13546
A	-11	HIS	-	expression tag	UNP Q13546
A	-10	GLY	-	expression tag	UNP Q13546
A	-9	GLY	-	expression tag	UNP Q13546
A	-8	GLY	-	expression tag	UNP Q13546
A	-7	GLU	-	expression tag	UNP Q13546
A	-6	ASN	-	expression tag	UNP Q13546
A	-5	LEU	-	expression tag	UNP Q13546
A	-4	TYR	-	expression tag	UNP Q13546
A	-3	PHE	-	expression tag	UNP Q13546
A	-2	GLN	-	expression tag	UNP Q13546
A	-1	GLY	-	expression tag	UNP Q13546
A	0	SER	-	expression tag	UNP Q13546
A	34	ALA	CYS	engineered mutation	UNP Q13546
A	127	ALA	CYS	engineered mutation	UNP Q13546
A	233	ALA	CYS	engineered mutation	UNP Q13546
A	240	ALA	CYS	engineered mutation	UNP Q13546
B	-18	MET	-	initiating methionine	UNP Q13546
B	-17	GLY	-	expression tag	UNP Q13546

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	HIS	-	expression tag	UNP Q13546
B	-15	HIS	-	expression tag	UNP Q13546
B	-14	HIS	-	expression tag	UNP Q13546
B	-13	HIS	-	expression tag	UNP Q13546
B	-12	HIS	-	expression tag	UNP Q13546
B	-11	HIS	-	expression tag	UNP Q13546
B	-10	GLY	-	expression tag	UNP Q13546
B	-9	GLY	-	expression tag	UNP Q13546
B	-8	GLY	-	expression tag	UNP Q13546
B	-7	GLU	-	expression tag	UNP Q13546
B	-6	ASN	-	expression tag	UNP Q13546
B	-5	LEU	-	expression tag	UNP Q13546
B	-4	TYR	-	expression tag	UNP Q13546
B	-3	PHE	-	expression tag	UNP Q13546
B	-2	GLN	-	expression tag	UNP Q13546
B	-1	GLY	-	expression tag	UNP Q13546
B	0	SER	-	expression tag	UNP Q13546
B	34	ALA	CYS	engineered mutation	UNP Q13546
B	127	ALA	CYS	engineered mutation	UNP Q13546
B	233	ALA	CYS	engineered mutation	UNP Q13546
B	240	ALA	CYS	engineered mutation	UNP Q13546

- Molecule 2 is (5R)-5-[(7-chloro-1H-indol-3-yl)methyl]-3-methylimidazolidine-2,4-dione (CCD ID: RCM) (formula: C₁₃H₁₂ClN₃O₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Cl	N	O	0	0
			19	13	1	3	2		
2	B	1	Total	C	Cl	N	O	0	0
			19	13	1	3	2		

- Molecule 3 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	O	S	6	0
			10	2	6	1	1		
3	B	1	Total	C	H	O	S	6	0
			10	2	6	1	1		

- Molecule 4 is 7-[4-[3-(methylsulfonylmethyl)azetidin-1-yl]sulfonylphenyl]quinoline (CCD ID: A1IKX) (formula: $C_{20}H_{20}N_2O_4S_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	S	0	0
			28	20	2	4	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	45	Total O 45 45	0	0
5	B	45	Total O 45 45	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	47.70Å 93.48Å 125.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	62.77 – 2.15 62.77 – 2.15	Depositor EDS
% Data completeness (in resolution range)	74.8 (62.77-2.15) 74.8 (62.77-2.15)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 2.14Å)	Xtriage
Refinement program	BUSTER 2.11.8 (8-JUN-2022) PACIOREK	Depositor
R, R_{free}	0.249 , 0.294 0.237 , 0.281	Depositor DCC
R_{free} test set	1180 reflections (3.74%)	wwPDB-VP
Wilson B-factor (Å ²)	53.3	Xtriage
Anisotropy	0.047	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 67.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4225	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: RCM, A1IKX, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	1/2045 (0.0%)	1.03	5/2755 (0.2%)
1	B	0.73	1/2084 (0.0%)	1.09	4/2808 (0.1%)
All	All	0.72	2/4129 (0.0%)	1.06	9/5563 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	193	MET	SD-CE	-7.87	1.59	1.79
1	A	193	MET	SD-CE	-6.96	1.62	1.79

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	106	ALA	CA-C-N	9.10	138.93	121.54
1	B	106	ALA	C-N-CA	9.10	138.93	121.54
1	B	57	ASN	CA-CB-CG	7.07	119.67	112.60
1	B	96	GLU	N-CA-C	6.89	119.64	111.71
1	A	57	ASN	CA-CB-CG	6.37	118.97	112.60
1	A	95	MET	CA-C-N	5.09	128.83	120.63
1	A	95	MET	C-N-CA	5.09	128.83	120.63
1	A	56	HIS	CA-C-N	5.07	127.78	120.38
1	A	56	HIS	C-N-CA	5.07	127.78	120.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2006	0	2017	17	0
1	B	2043	0	2045	19	0
2	A	19	0	12	0	0
2	B	19	0	12	0	0
3	A	4	6	6	2	0
3	B	4	6	6	2	0
4	B	28	0	0	0	0
5	A	45	0	0	0	0
5	B	45	0	0	0	0
All	All	4213	12	4098	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:130:HIS:ND1	3:A:302:DMS:S	2.34	0.99
1:B:130:HIS:ND1	3:B:303:DMS:S	2.39	0.96
1:B:207:GLU:OE1	3:B:303:DMS:O	2.06	0.73
1:A:193:MET:HE2	1:A:198:LEU:HD23	1.75	0.69
1:A:207:GLU:OE1	3:A:302:DMS:O	2.12	0.68
1:B:236:GLN:HG3	1:B:237:LEU:N	2.08	0.68
1:B:193:MET:HE2	1:B:198:LEU:HD23	1.81	0.62
1:A:62:GLU:O	1:A:66:MET:HG3	2.03	0.58
1:A:68:ASN:ND2	1:A:71:ARG:HH21	2.02	0.57
1:B:18:LEU:HD21	1:B:37:ARG:HE	1.71	0.55
1:B:68:ASN:ND2	1:B:71:ARG:HH21	2.03	0.55
1:A:59:ALA:HB3	1:A:164:MET:HE2	1.90	0.54
1:B:236:GLN:HG3	1:B:237:LEU:H	1.73	0.54
1:B:130:HIS:NE2	1:B:210:ASP:OD2	2.44	0.51
1:A:130:HIS:NE2	1:A:210:ASP:OD2	2.44	0.50
1:B:197:HIS:CE1	1:B:205:PRO:HA	2.48	0.48
1:A:235:GLN:H	1:A:235:GLN:NE2	2.14	0.46
1:B:81:VAL:HG12	1:B:90:LEU:HD23	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:59:ALA:HB3	1:B:164:MET:HE2	1.99	0.44
1:B:97:LYS:HE2	1:B:148:ASN:HA	1.99	0.44
1:A:81:VAL:HG12	1:A:90:LEU:HD23	2.00	0.44
1:B:28:PHE:CZ	1:B:140:LYS:HE3	2.54	0.43
1:A:197:HIS:CE1	1:A:205:PRO:HA	2.53	0.43
1:A:83:ILE:HG23	1:A:88:TYR:HE1	1.84	0.43
1:A:193:MET:HE2	1:A:198:LEU:CD2	2.47	0.42
1:A:72:HIS:CD2	1:A:74:ARG:H	2.37	0.42
1:A:92:MET:HE1	1:A:157:LEU:CD2	2.49	0.42
1:B:72:HIS:CD2	1:B:74:ARG:H	2.38	0.42
1:B:146:VAL:HG22	1:B:152:ILE:HG22	2.01	0.42
1:B:235:GLN:NE2	1:B:235:GLN:H	2.17	0.42
1:A:236:GLN:HG3	1:A:237:LEU:N	2.35	0.41
1:A:83:ILE:HG23	1:A:88:TYR:CE1	2.56	0.41
1:B:124:GLU:OE1	1:B:286:ARG:NH2	2.50	0.41
1:B:83:ILE:HG23	1:B:88:TYR:HE1	1.85	0.41
1:A:195:PRO:HA	1:A:198:LEU:HD12	2.02	0.41
1:B:266:LEU:HG	1:B:276:PRO:HD3	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	245/313 (78%)	235 (96%)	8 (3%)	2 (1%)	16	9
1	B	253/313 (81%)	243 (96%)	8 (3%)	2 (1%)	16	9
All	All	498/626 (80%)	478 (96%)	16 (3%)	4 (1%)	16	9

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	107	GLU
1	A	107	GLU
1	A	85	GLU
1	B	231	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	216/271 (80%)	200 (93%)	16 (7%)	13	8
1	B	219/271 (81%)	205 (94%)	14 (6%)	16	11
All	All	435/542 (80%)	405 (93%)	30 (7%)	14	9

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	SER
1	A	31	VAL
1	A	33	LEU
1	A	46	THR
1	A	56	HIS
1	A	57	ASN
1	A	61	LEU
1	A	62	GLU
1	A	77	LYS
1	A	92	MET
1	A	204	LYS
1	A	235	GLN
1	A	236	GLN
1	A	239	MET
1	A	258	ARG
1	A	284	LYS
1	B	32	SER
1	B	56	HIS
1	B	57	ASN
1	B	61	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	62	GLU
1	B	66	MET
1	B	77	LYS
1	B	107	GLU
1	B	108	MET
1	B	239	MET
1	B	258	ARG
1	B	284	LYS
1	B	290	LEU
1	B	294	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	68	ASN
1	A	72	HIS
1	A	235	GLN
1	A	271	ASN
1	A	292	GLN
1	B	68	ASN
1	B	72	HIS
1	B	148	ASN
1	B	230	ASN
1	B	235	GLN
1	B	271	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	A1IKX	B	302	-	27,31,31	1.64	3 (11%)	42,47,47	1.62	8 (19%)
2	RCM	A	301	-	21,21,21	1.04	2 (9%)	26,31,31	1.54	5 (19%)
3	DMS	B	303	-	3,3,3	1.37	0	3,3,3	1.77	1 (33%)
3	DMS	A	302	-	3,3,3	1.39	0	3,3,3	1.59	1 (33%)
2	RCM	B	301	-	21,21,21	1.20	2 (9%)	26,31,31	1.44	4 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	A1IKX	B	302	-	-	5/19/29/29	0/4/4/4
2	RCM	A	301	-	-	0/4/20/20	0/3/3/3
2	RCM	B	301	-	-	0/4/20/20	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	302	A1IKX	S10-N8	5.95	1.71	1.63
2	B	301	RCM	C12-N3	-3.58	1.35	1.38
2	A	301	RCM	C12-N3	-3.34	1.35	1.38
4	B	302	A1IKX	C5-C6	-3.20	1.50	1.53
2	B	301	RCM	C14-N3	-2.28	1.34	1.38
4	B	302	A1IKX	O12-S10	2.28	1.45	1.43
2	A	301	RCM	C14-N3	-2.10	1.35	1.38

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	RCM	O1-C12-N3	5.39	128.41	124.02
2	B	301	RCM	O1-C12-N3	4.95	128.05	124.02
4	B	302	A1IKX	O12-S10-N8	3.91	110.37	106.69
4	B	302	A1IKX	C1-S2-C5	-3.78	98.83	103.12
4	B	302	A1IKX	C6-C5-S2	-3.23	106.56	114.55
4	B	302	A1IKX	O3-S2-O4	3.13	123.09	117.22
4	B	302	A1IKX	C13-S10-N8	-3.09	103.54	107.28
4	B	302	A1IKX	C6-C7-N8	2.68	90.58	88.38
2	A	301	RCM	C12-C11-N2	2.64	104.60	101.12
2	B	301	RCM	C4-C3-N1	2.55	133.82	129.51
2	B	301	RCM	C12-C11-N2	2.50	104.41	101.12
3	A	302	DMS	C2-S-C1	2.47	111.10	98.42
3	B	303	DMS	C2-S-C1	2.36	110.53	98.42
4	B	302	A1IKX	O12-S10-O11	2.30	123.18	119.59
2	B	301	RCM	C11-C12-N3	-2.26	105.24	107.72
2	A	301	RCM	C4-C3-N1	2.25	133.31	129.51
2	A	301	RCM	C11-C12-N3	-2.15	105.35	107.72
4	B	302	A1IKX	C6-C9-N8	2.15	90.15	88.38
2	A	301	RCM	O2-C14-N2	-2.14	122.95	126.84

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	302	A1IKX	C7-N8-S10-O12
4	B	302	A1IKX	C6-C5-S2-O4
4	B	302	A1IKX	C7-N8-S10-C13
4	B	302	A1IKX	C9-N8-S10-O12
4	B	302	A1IKX	C6-C5-S2-O3


There are no ring outliers.


2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	303	DMS	2	0
3	A	302	DMS	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	253/313 (80%)	1.04	39 (15%) 5 6	39, 63, 94, 118	0
1	B	259/313 (82%)	1.25	61 (23%) 2 2	41, 71, 123, 134	0
All	All	512/626 (81%)	1.14	100 (19%) 3 4	39, 67, 112, 134	0

All (100) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	20	SER	6.6
1	B	9	VAL	5.2
1	B	232	ILE	4.8
1	B	233	ALA	4.8
1	B	252	ILE	4.4
1	B	188	GLY	4.4
1	B	10	ILE	4.4
1	A	293	LEU	4.3
1	B	230	ASN	4.3
1	A	168	LEU	4.0
1	A	189	THR	4.0
1	B	231	ALA	3.9
1	B	21	ALA	3.9
1	B	168	LEU	3.9
1	B	255	TYR	3.8
1	A	201	VAL	3.7
1	A	252	ILE	3.7
1	A	9	VAL	3.6
1	B	31	VAL	3.6
1	B	8	ASN	3.6
1	B	56	HIS	3.5
1	A	58	GLU	3.4
1	B	41	LEU	3.4
1	B	60	LEU	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	237	LEU	3.4
1	A	18	LEU	3.3
1	B	83	ILE	3.2
1	A	23	LEU	3.2
1	B	23	LEU	3.2
1	B	112	LEU	3.2
1	A	88	TYR	3.1
1	B	253	THR	3.1
1	B	35	PHE	3.1
1	B	258	ARG	3.0
1	B	200	ASP	3.0
1	B	12	MET	3.0
1	A	188	GLY	2.9
1	A	61	LEU	2.9
1	A	8	ASN	2.9
1	A	255	TYR	2.8
1	B	40	GLY	2.8
1	B	57	ASN	2.7
1	B	18	LEU	2.7
1	A	56	HIS	2.7
1	A	109	SER	2.7
1	A	60	LEU	2.7
1	B	103	VAL	2.6
1	B	236	GLN	2.6
1	B	85	GLU	2.6
1	A	57	ASN	2.6
1	A	139	LEU	2.6
1	B	191	TYR	2.5
1	A	10	ILE	2.5
1	B	46	THR	2.5
1	A	83	ILE	2.5
1	B	249	VAL	2.4
1	B	66	MET	2.4
1	B	39	GLN	2.4
1	B	108	MET	2.4
1	B	59	ALA	2.4
1	A	15	SER	2.4
1	A	203	ALA	2.4
1	B	294	GLU	2.4
1	B	11	LYS	2.4
1	A	46	THR	2.4
1	A	191	TYR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	17	PHE	2.3
1	B	261	ILE	2.3
1	A	271	ASN	2.3
1	B	202	ASN	2.3
1	B	290	LEU	2.3
1	B	14	SER	2.3
1	B	119	ILE	2.2
1	B	58	GLU	2.2
1	A	20	SER	2.2
1	A	164	MET	2.2
1	B	272	PRO	2.2
1	A	231	ALA	2.2
1	A	192	TYR	2.2
1	B	228	TYR	2.2
1	B	22	GLU	2.1
1	B	150	PHE	2.1
1	B	243	SER	2.1
1	A	66	MET	2.1
1	A	202	ASN	2.1
1	A	41	LEU	2.1
1	B	33	LEU	2.1
1	B	19	GLU	2.1
1	B	250	ASP	2.1
1	B	28	PHE	2.1
1	A	166	SER	2.1
1	A	38	THR	2.1
1	A	86	GLY	2.1
1	B	244	GLY	2.1
1	A	167	LYS	2.1
1	A	251	ASP	2.0
1	B	248	ASP	2.0
1	B	251	ASP	2.0
1	A	190	LEU	2.0
1	B	24	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	A1IKX	B	302	28/28	0.84	0.13	72,81,95,95	0
2	RCM	A	301	19/19	0.86	0.11	47,49,52,55	0
2	RCM	B	301	19/19	0.90	0.10	53,53,55,57	0
3	DMS	A	302	4/4	0.91	0.21	20,23,30,33	6
3	DMS	B	303	4/4	0.93	0.19	28,28,30,32	6

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around A1IKX B 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around RCM A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around RCM B 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.