



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 18, 2026 – 12:38 PM UTC

PDB ID : 2GUO / pdb_00002guo
Title : Human Class I MHC HLA-A2 in complex with the native nonameric Melan-A/MART-1(27-35) peptide
Authors : Borbulevych, O.Y.; Baker, B.M.
Deposited on : 2006-05-01
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	2% 90% 8%
1	D	275	3% 91% 8%
2	B	100	2% 95% 5%
2	E	100	% 90% 8%
3	C	9	22% 78% 22%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	9	 78% 22%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6874 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, A-2 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	1	0
			2251	1405	410	427	9			
1	D	275	Total	C	N	O	S	0	1	0
			2251	1406	409	427	9			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			
2	E	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769
E	0	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called Peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O	0	0	0
			57	37	9	11			
3	F	9	Total	C	N	O	0	1	0
			61	41	9	11			

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		
4	E	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	E	1	Total	Na	0	0
			1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	195	Total	O	0	0
			195	195		
6	B	90	Total	O	0	0
			90	90		
6	C	7	Total	O	0	0
			7	7		
6	D	180	Total	O	0	0
			180	180		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	E	76	Total 76	O 76	0	0
6	F	7	Total 7	O 7	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class I histocompatibility antigen, A-2 alpha chain




- Molecule 1: HLA class I histocompatibility antigen, A-2 alpha chain




- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin



- Molecule 3: Peptide



- Molecule 3: Peptide

Chain F:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	83.96Å 58.37Å 89.42Å 90.00° 109.66° 90.00°	Depositor
Resolution (Å)	20.00 – 1.90 20.00 – 1.90	Depositor EDS
% Data completeness (in resolution range)	96.3 (20.00-1.90) 96.8 (20.00-1.90)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.64 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.187 , 0.244 0.187 , 0.243	Depositor DCC
R_{free} test set	3183 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	15.4	Xtriage
Anisotropy	0.379	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 36.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6874	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.00	1/2320 (0.0%)	1.13	3/3148 (0.1%)
1	D	0.96	1/2317 (0.0%)	1.10	6/3144 (0.2%)
2	B	0.99	0/860	1.03	2/1162 (0.2%)
2	E	0.97	0/860	1.01	2/1162 (0.2%)
3	C	1.08	0/56	0.87	0/74
3	F	1.23	0/64	1.05	0/85
All	All	0.98	2/6477 (0.0%)	1.09	13/8775 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	268	LYS	CE-NZ	8.20	1.74	1.49
1	A	135	ALA	CA-CB	-5.52	1.45	1.53

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	29	ASP	CA-CB-CG	7.73	120.33	112.60
1	D	30	ASP	CA-C-O	7.60	128.92	119.95
1	D	14	ARG	CA-C-N	7.49	129.21	119.84
1	D	14	ARG	C-N-CA	7.49	129.21	119.84
1	D	249	VAL	N-CA-CB	-6.88	101.57	111.21

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	71	THR	CA-C-N	6.80	126.76	119.76
2	B	71	THR	C-N-CA	6.80	126.76	119.76
1	D	30	ASP	O-C-N	-6.66	114.07	122.26
1	A	249	VAL	N-CA-CB	-6.15	102.59	111.21
2	E	71	THR	CA-C-N	5.81	125.75	119.76
2	E	71	THR	C-N-CA	5.81	125.75	119.76
1	A	176	LYS	N-CA-C	5.79	117.59	111.28
1	A	222	GLU	CB-CG-CD	5.58	122.08	112.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	136[B]	ALA	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2251	0	2098	13	0
1	D	2251	0	2100	9	0
2	B	837	0	803	1	0
2	E	837	0	803	3	0
3	C	57	0	67	1	0
3	F	61	0	74	1	0
4	A	12	0	16	0	0
4	D	6	0	8	1	0
4	E	6	0	8	0	0
5	E	1	0	0	0	0
6	A	195	0	0	2	0
6	B	90	0	0	0	0
6	C	7	0	0	0	0
6	D	180	0	0	2	0
6	E	76	0	0	0	0
6	F	7	0	0	0	0
All	All	6874	0	5977	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:268:LYS:CE	1:D:268:LYS:NZ	1.73	1.51
1:D:268:LYS:NZ	1:D:268:LYS:CD	2.53	0.71
1:A:19:GLU:OE1	1:A:75:ARG:NH1	2.26	0.69
1:A:218:GLN:HE21	1:A:221:GLY:HA2	1.61	0.65
1:A:108:ARG:NH1	1:D:174:ASN:OD1	2.22	0.65
1:A:65:ARG:NH1	6:A:726:HOH:O	2.33	0.61
1:D:157:ARG:NH1	6:D:633:HOH:O	2.38	0.56
1:A:218:GLN:NE2	1:A:221:GLY:HA2	2.20	0.56
1:A:70:HIS:HA	3:C:6:ILE:HD11	1.89	0.54
1:A:108:ARG:NH2	6:A:792:HOH:O	2.45	0.49
1:A:249:VAL:HG22	1:A:257:TYR:CZ	2.48	0.49
1:A:110:LEU:HD11	1:D:169:ARG:HD3	1.96	0.48
1:A:173:GLU:O	1:A:176:LYS:NZ	2.40	0.48
1:A:196:ASP:OD1	1:A:196:ASP:N	2.48	0.47
2:E:48:LYS:H	2:E:48:LYS:HG2	1.35	0.46
1:D:77:ASP:OD1	3:F:8:THR:OG1	2.25	0.45
1:A:51:TRP:O	1:A:54:GLN:NE2	2.49	0.44
4:D:604:GOL:O3	6:D:638:HOH:O	2.21	0.44
1:D:249:VAL:HG22	1:D:257:TYR:CZ	2.55	0.42
2:E:6:LYS:N	2:E:6:LYS:HD3	2.34	0.42
1:D:218:GLN:NE2	1:D:221:GLY:HA2	2.36	0.41
1:A:126:LEU:HG	1:A:130:LEU:HA	2.02	0.40
2:B:75:LYS:HB2	2:B:75:LYS:HE3	1.86	0.40
1:D:117:ALA:HB2	2:E:60:TRP:CE2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/275 (100%)	270 (98%)	4 (2%)	0	100	100
1	D	274/275 (100%)	265 (97%)	8 (3%)	1 (0%)	30	22
2	B	98/100 (98%)	98 (100%)	0	0	100	100
2	E	98/100 (98%)	98 (100%)	0	0	100	100
3	C	7/9 (78%)	7 (100%)	0	0	100	100
3	F	8/9 (89%)	8 (100%)	0	0	100	100
All	All	759/768 (99%)	746 (98%)	12 (2%)	1 (0%)	48	40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	15	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/231 (100%)	221 (95%)	11 (5%)	23	15
1	D	231/231 (100%)	220 (95%)	11 (5%)	23	15
2	B	95/95 (100%)	92 (97%)	3 (3%)	34	27
2	E	95/95 (100%)	87 (92%)	8 (8%)	10	4
3	C	5/5 (100%)	4 (80%)	1 (20%)	1	0
3	F	6/5 (120%)	5 (83%)	1 (17%)	2	0
All	All	664/662 (100%)	629 (95%)	35 (5%)	20	12

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	ARG
1	A	35	ARG
1	A	110	LEU
1	A	113	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	156	LEU
1	A	215	LEU
1	A	222	GLU
1	A	225	THR
1	A	249	VAL
1	A	255	GLN
1	A	272	LEU
2	B	0	MET
2	B	48	LYS
2	B	89	GLN
3	C	7	LEU
1	D	89	GLU
1	D	113	TYR
1	D	156	LEU
1	D	182	THR
1	D	215	LEU
1	D	222	GLU
1	D	223	ASP
1	D	225	THR
1	D	249	VAL
1	D	255	GLN
1	D	272	LEU
2	E	0	MET
2	E	6	LYS
2	E	47	GLU
2	E	48	LYS
2	E	74	GLU
2	E	75	LYS
2	E	89	GLN
2	E	92	ILE
3	F	6	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	115	GLN
1	A	218	GLN
1	A	255	GLN
1	A	262	GLN
1	D	72	GLN
1	D	197	HIS
1	D	218	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	226	GLN
1	D	255	GLN
1	D	262	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	A	602	-	5,5,5	0.26	0	5,5,5	0.70	0
4	GOL	A	603	-	5,5,5	0.44	0	5,5,5	0.82	0
4	GOL	D	604	-	5,5,5	0.30	0	5,5,5	0.73	0
4	GOL	E	601	-	5,5,5	0.32	0	5,5,5	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	602	-	-	4/4/4/4	-
4	GOL	A	603	-	-	2/4/4/4	-
4	GOL	D	604	-	-	2/4/4/4	-
4	GOL	E	601	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	602	GOL	O1-C1-C2-C3
4	A	602	GOL	C1-C2-C3-O3
4	A	602	GOL	O2-C2-C3-O3
4	D	604	GOL	C1-C2-C3-O3
4	A	602	GOL	O1-C1-C2-O2
4	E	601	GOL	O1-C1-C2-O2
4	E	601	GOL	O1-C1-C2-C3
4	D	604	GOL	O2-C2-C3-O3
4	A	603	GOL	O2-C2-C3-O3
4	A	603	GOL	O1-C1-C2-C3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	604	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/275 (100%)	0.36	5 (1%) 67 71	6, 14, 22, 26	1 (0%)
1	D	275/275 (100%)	0.39	9 (3%) 49 52	6, 13, 22, 30	1 (0%)
2	B	100/100 (100%)	0.24	2 (2%) 65 69	8, 14, 22, 24	0
2	E	100/100 (100%)	0.29	1 (1%) 79 82	8, 14, 22, 26	0
3	C	9/9 (100%)	1.10	2 (22%) 2 2	17, 20, 29, 31	0
3	F	9/9 (100%)	0.66	0 100 100	15, 20, 27, 27	1 (11%)
All	All	768/768 (100%)	0.36	19 (2%) 58 62	6, 14, 22, 31	3 (0%)

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	1	GLY	6.0
3	C	6	ILE	2.7
1	D	89	GLU	2.7
2	B	77	GLU	2.5
1	A	1	GLY	2.4
1	A	264	GLU	2.4
1	D	18	GLY	2.4
2	E	48	LYS	2.3
1	D	113	TYR	2.3
1	D	222	GLU	2.3
1	D	223	ASP	2.2
1	A	196	ASP	2.2
1	D	136[A]	ALA	2.2
1	D	232	GLU	2.2
1	D	196	ASP	2.1
1	A	19	GLU	2.1
3	C	4	ILE	2.1
2	B	34	ASP	2.1
1	A	2	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	A	603	6/6	0.71	0.15	47,48,48,50	0
4	GOL	D	604	6/6	0.76	0.15	39,39,41,41	0
4	GOL	E	601	6/6	0.80	0.15	45,46,48,51	0
4	GOL	A	602	6/6	0.81	0.13	30,35,36,39	0
5	NA	E	605	1/1	0.93	0.08	25,25,25,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.