



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:29 PM UTC

PDB ID : 4GU4 / pdb_00004gu4
Title : Crystal structure of the T1L reovirus attachment protein sigma1 in complex with alpha-2,3-sialyllactose
Authors : Reiss, K.; Stehle, T.
Deposited on : 2012-08-29
Resolution : 3.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1085 (3.54-3.46)
Clashscore	190562	1140 (3.54-3.46)
Ramachandran outliers	187476	1113 (3.54-3.46)
Sidechain outliers	187428	1114 (3.54-3.46)
RSRZ outliers	180081	1084 (3.54-3.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	219	
1	B	219	
1	C	219	
2	D	3	
2	E	3	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	2	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4912 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Outer capsid protein sigma-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	205	Total	C	N	O	S	0	0	0
			1589	1016	265	301	7			
1	B	205	Total	C	N	O	S	0	0	0
			1607	1024	267	309	7			
1	C	205	Total	C	N	O	S	0	0	0
			1598	1020	267	304	7			

There are 27 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	252	GLY	-	expression tag	UNP P04506
A	253	VAL	-	expression tag	UNP P04506
A	254	LEU	-	expression tag	UNP P04506
A	255	ASN	-	expression tag	UNP P04506
A	256	GLN	-	expression tag	UNP P04506
A	257	GLY	-	expression tag	UNP P04506
A	258	VAL	-	expression tag	UNP P04506
A	259	THR	-	expression tag	UNP P04506
A	260	SER	-	expression tag	UNP P04506
B	252	GLY	-	expression tag	UNP P04506
B	253	VAL	-	expression tag	UNP P04506
B	254	LEU	-	expression tag	UNP P04506
B	255	ASN	-	expression tag	UNP P04506
B	256	GLN	-	expression tag	UNP P04506
B	257	GLY	-	expression tag	UNP P04506
B	258	VAL	-	expression tag	UNP P04506
B	259	THR	-	expression tag	UNP P04506
B	260	SER	-	expression tag	UNP P04506
C	252	GLY	-	expression tag	UNP P04506
C	253	VAL	-	expression tag	UNP P04506
C	254	LEU	-	expression tag	UNP P04506
C	255	ASN	-	expression tag	UNP P04506
C	256	GLN	-	expression tag	UNP P04506

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	257	GLY	-	expression tag	UNP P04506
C	258	VAL	-	expression tag	UNP P04506
C	259	THR	-	expression tag	UNP P04506
C	260	SER	-	expression tag	UNP P04506

- Molecule 2 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	3	Total	C	N	O	0	0	0
			43	23	1	19			
2	E	3	Total	C	N	O	0	0	0
			43	23	1	19			

- Molecule 3 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	F	2	Total	C	N	O	0	0	0
			32	17	1	14			

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Outer capsid protein sigma-1

Chain A: 




- Molecule 1: Outer capsid protein sigma-1

Chain B: 



- Molecule 1: Outer capsid protein sigma-1

Chain C: 



- Molecule 2: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranose

Chain D: 



- Molecule 2: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranose

Chain E:  67% 33%


EGF1
GAL2
SIA3

- Molecule 3: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose

Chain F:  50% 50%


GAL3
SIA2

4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	149.43Å 149.43Å 165.19Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.91 – 3.50 48.91 – 3.50	Depositor EDS
% Data completeness (in resolution range)	97.8 (48.91-3.50) 97.7 (48.91-3.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.46 (at 3.01Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.186 , 0.197 0.200 , 0.210	Depositor DCC
R_{free} test set	4195 reflections (9.75%)	wwPDB-VP
Wilson B-factor (Å ²)	76.6	Xtriage
Anisotropy	0.427	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 73.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.026 for -h,-k,l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4912	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, GAL, SIA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	0/1630	1.02	2/2231 (0.1%)
1	B	0.77	0/1648	1.03	4/2254 (0.2%)
1	C	0.71	0/1639	1.01	1/2242 (0.0%)
All	All	0.75	0/4917	1.02	7/6727 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	268	SER	N-CA-C	6.76	118.48	108.60
1	B	289	ASP	CA-C-N	6.05	128.38	120.28
1	B	289	ASP	C-N-CA	6.05	128.38	120.28
1	A	303	ARG	N-CA-C	-5.42	100.88	109.76
1	A	307	THR	CB-CA-C	5.17	118.28	109.75
1	B	307	THR	CB-CA-C	5.15	118.25	109.75
1	B	375	ASN	CA-CB-CG	5.06	117.66	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1589	0	1508	25	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1607	0	1543	11	0
1	C	1598	0	1525	21	0
2	D	43	0	37	0	0
2	E	43	0	37	0	0
3	F	32	0	28	0	0
All	All	4912	0	4678	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:409:LEU:HB2	1:A:433:VAL:HG13	1.50	0.91
1:A:420:ASN:O	1:A:421:ASN:CB	2.30	0.80
1:A:409:LEU:HB2	1:A:433:VAL:CG1	2.12	0.79
1:A:420:ASN:O	1:A:421:ASN:HB2	1.82	0.77
1:A:311:TYR:HD2	1:B:324:VAL:HB	1.62	0.64
1:C:296:ILE:HG22	1:C:297:ASN:ND2	2.18	0.59
1:A:311:TYR:CD2	1:B:324:VAL:HB	2.40	0.57
1:A:272:PRO:HG3	1:B:295:ILE:HD12	1.91	0.52
1:A:433:VAL:HG11	1:A:465:TYR:HB3	1.92	0.51
1:A:428:PHE:O	1:A:429:ARG:HD2	2.12	0.49
1:A:275:VAL:HG22	1:A:284:LEU:HD13	1.95	0.48
1:A:324:VAL:HB	1:C:311:TYR:CD2	2.49	0.48
1:C:419:LEU:CD2	1:C:424:ARG:HG2	2.44	0.48
1:A:274:THR:O	1:A:284:LEU:HD12	2.14	0.47
1:B:428:PHE:O	1:B:429:ARG:HD2	2.14	0.47
1:C:428:PHE:O	1:C:429:ARG:HD2	2.14	0.47
1:A:277:GLU:HA	1:A:281:VAL:O	2.15	0.46
1:C:420:ASN:HB2	1:C:453:HIS:HA	1.97	0.46
1:C:269:VAL:HG21	1:C:275:VAL:HG23	1.98	0.45
1:C:271:PRO:HA	1:C:272:PRO:C	2.42	0.45
1:B:311:TYR:CD2	1:C:324:VAL:HB	2.53	0.44
1:A:412:THR:HG22	1:A:463:ILE:HD12	1.99	0.44
1:A:308:LEU:HG	1:C:306:LEU:HD13	1.99	0.44
1:A:420:ASN:O	1:A:421:ASN:HB3	2.17	0.44
1:B:333:ARG:HB3	1:B:360:LYS:HB2	2.00	0.44
1:C:412:THR:HG22	1:C:463:ILE:HD12	1.99	0.44
1:B:412:THR:HG22	1:B:463:ILE:HD12	1.99	0.44
1:C:333:ARG:HB3	1:C:360:LYS:HB2	1.98	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:419:LEU:HD22	1:C:424:ARG:HG2	1.99	0.43
1:C:289:ASP:HB3	1:C:292:ASP:HB2	1.99	0.43
1:B:292:ASP:CG	1:C:302:LEU:HD12	2.43	0.43
1:C:419:LEU:HB2	1:C:455:GLN:HG3	2.01	0.43
1:B:277:GLU:HB2	1:B:282:ARG:HE	1.84	0.43
1:B:429:ARG:NH1	1:B:449:THR:O	2.52	0.42
1:A:333:ARG:HB3	1:A:360:LYS:HB2	2.00	0.42
1:C:429:ARG:NH1	1:C:449:THR:O	2.52	0.42
1:A:270:LEU:HD13	1:B:285:SER:HB3	2.02	0.42
1:A:271:PRO:HA	1:A:272:PRO:C	2.45	0.42
1:A:284:LEU:HD12	1:A:285:SER:H	1.85	0.42
1:C:386:ASN:OD1	1:C:388:VAL:HG13	2.20	0.41
1:A:286:PHE:HB3	1:C:272:PRO:HD2	2.01	0.41
1:A:284:LEU:HD12	1:A:284:LEU:HA	1.79	0.41
1:A:401:GLU:HA	1:A:402:PRO:HD3	1.98	0.41
1:C:387:PHE:O	1:C:391:LEU:CD2	2.69	0.41
1:C:401:GLU:HA	1:C:402:PRO:HD3	1.97	0.41
1:A:407:ASP:HA	1:A:408:PRO:HA	1.95	0.40
1:C:269:VAL:HG11	1:C:284:LEU:HD13	2.02	0.40
1:A:429:ARG:NH1	1:A:449:THR:O	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	203/219 (93%)	197 (97%)	5 (2%)	1 (0%)	24	57
1	B	203/219 (93%)	200 (98%)	3 (2%)	0	100	100
1	C	203/219 (93%)	202 (100%)	1 (0%)	0	100	100
All	All	609/657 (93%)	599 (98%)	9 (2%)	1 (0%)	43	74

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	421	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/192 (88%)	167 (99%)	2 (1%)	63	73
1	B	177/192 (92%)	175 (99%)	2 (1%)	65	74
1	C	172/192 (90%)	170 (99%)	2 (1%)	63	73
All	All	518/576 (90%)	512 (99%)	6 (1%)	63	73

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	290	THR
1	A	367	LEU
1	B	282	ARG
1	B	283	THR
1	C	283	THR
1	C	285	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	340	GLN
1	A	421	ASN
1	A	445	ASN
1	B	321	ASN
1	B	340	GLN
1	C	297	ASN
1	C	321	ASN
1	C	322	ASN
1	C	340	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	445	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BGC	D	1	2	12,12,12	0.29	0	17,17,17	0.39	0
2	GAL	D	2	2	11,11,12	0.28	0	15,15,17	0.81	1 (6%)
2	SIA	D	3	2	20,20,21	0.70	0	21,28,31	1.30	3 (14%)
2	BGC	E	1	2	12,12,12	0.22	0	17,17,17	0.53	0
2	GAL	E	2	2	11,11,12	0.30	0	15,15,17	0.64	0
2	SIA	E	3	2	20,20,21	0.73	0	21,28,31	1.38	4 (19%)
3	GAL	F	1	3	12,12,12	0.31	0	17,17,17	0.45	0
3	SIA	F	2	3	20,20,21	0.76	0	21,28,31	1.37	4 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	D	1	2	-	0/2/22/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GAL	D	2	2	-	0/2/19/22	0/1/1/1
2	SIA	D	3	2	-	4/18/34/38	0/1/1/1
2	BGC	E	1	2	-	1/2/22/22	0/1/1/1
2	GAL	E	2	2	-	1/2/19/22	0/1/1/1
2	SIA	E	3	2	-	5/18/34/38	0/1/1/1
3	GAL	F	1	3	-	1/2/22/22	0/1/1/1
3	SIA	F	2	3	-	4/18/34/38	0/1/1/1

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	2	SIA	O6-C2-C1	2.92	113.22	107.72
2	D	3	SIA	O6-C2-C1	2.63	112.69	107.72
2	E	3	SIA	C6-C5-N5	-2.52	106.88	110.91
2	E	3	SIA	O1B-C1-C2	2.50	119.22	112.71
2	D	3	SIA	C6-C5-N5	-2.50	106.92	110.91
2	D	2	GAL	C1-O5-C5	2.47	115.49	112.19
2	D	3	SIA	O8-C8-C7	2.45	114.98	109.25
2	E	3	SIA	O8-C8-C7	2.43	114.94	109.25
3	F	2	SIA	O8-C8-C7	2.29	114.61	109.25
3	F	2	SIA	O9-C9-C8	-2.22	106.48	111.16
3	F	2	SIA	C6-C5-N5	-2.10	107.56	110.91
2	E	3	SIA	O9-C9-C8	-2.08	106.79	111.16

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	3	SIA	C6-C7-C8-C9
2	D	3	SIA	C6-C7-C8-O8
2	D	3	SIA	O7-C7-C8-C9
2	D	3	SIA	O7-C7-C8-O8
2	E	3	SIA	C6-C7-C8-C9
2	E	3	SIA	C6-C7-C8-O8
2	E	3	SIA	O7-C7-C8-C9
2	E	3	SIA	O7-C7-C8-O8
3	F	2	SIA	C6-C7-C8-C9
3	F	2	SIA	C6-C7-C8-O8
3	F	2	SIA	O7-C7-C8-C9

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	F	2	SIA	O7-C7-C8-O8
2	E	1	BGC	O5-C5-C6-O6
3	F	1	GAL	O5-C5-C6-O6
2	E	3	SIA	O1A-C1-C2-O6
2	E	2	GAL	O5-C5-C6-O6


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	205/219 (93%)	-0.07	7 (3%)	48	27	55, 77, 109, 124	0
1	B	205/219 (93%)	-0.11	4 (1%)	65	39	56, 79, 116, 135	0
1	C	205/219 (93%)	0.03	8 (3%)	43	23	56, 91, 112, 121	0
All	All	615/657 (93%)	-0.05	19 (3%)	51	29	55, 81, 112, 135	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	278	ALA	5.1
1	A	277	GLU	4.1
1	A	421	ASN	3.8
1	C	277	GLU	3.7
1	B	265	LYS	3.6
1	A	469	VAL	3.5
1	C	469	VAL	3.0
1	B	469	VAL	2.9
1	C	278	ALA	2.9
1	C	423	GLN	2.8
1	A	279	SER	2.7
1	C	279	SER	2.6
1	C	265	LYS	2.6
1	C	282	ARG	2.3
1	C	267	ASP	2.3
1	A	265	LYS	2.3
1	B	306	LEU	2.1
1	B	307	THR	2.1
1	A	306	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BGC	E	1	12/12	0.57	0.13	119,124,131,131	0
2	BGC	D	1	12/12	0.75	0.12	118,124,126,127	0
3	GAL	F	1	12/12	0.78	0.11	110,123,126,126	0
2	GAL	E	2	11/12	0.88	0.11	98,110,116,118	0
2	GAL	D	2	11/12	0.92	0.08	97,104,111,114	0
3	SIA	F	2	20/21	0.93	0.12	77,89,107,108	0
2	SIA	D	3	20/21	0.95	0.09	82,88,104,106	0
2	SIA	E	3	20/21	0.95	0.10	65,76,107,107	0


The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.