



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 18, 2026 – 04:41 PM UTC

PDB ID : 4GU8 / pdb_00004gu8
Title : Crystal Structure of Burkholderia oklahomensis agglutinin (BOA)
Authors : Whitley, M.J.; Furey, W.; Gronenborn, A.M.
Deposited on : 2012-08-29
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	279	94% 6%
1	B	279	95% 5%
1	C	279	% 95% ...
1	D	279	96% .
1	E	279	% 93% 6% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	279	 96%
1	G	279	 95%
1	H	279	 95%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	DTT	E	301	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 17085 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Burkholderia oklahomensis agglutinin (BOA).


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	279	Total	C	N	O	S	0	0	0
			2046	1270	365	401	10			
1	B	279	Total	C	N	O	S	0	0	0
			2046	1270	365	401	10			
1	C	275	Total	C	N	O	S	0	0	0
			2016	1253	361	393	9			
1	D	279	Total	C	N	O	S	0	1	0
			2050	1273	366	401	10			
1	E	275	Total	C	N	O	S	0	0	0
			2016	1253	361	393	9			
1	F	276	Total	C	N	O	S	0	1	0
			2029	1261	363	395	10			
1	G	275	Total	C	N	O	S	0	0	0
			2016	1253	361	393	9			
1	H	276	Total	C	N	O	S	0	1	0
			2029	1261	363	395	10			

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is 2,3-DIHYDROXY-1,4-DITHIOBUTANE (CCD ID: DTT) (formula: $C_4H_{10}O_2S_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	E	1	Total	C	O	S	0	0
			8	4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	131	Total	O	0	0
			131	131		
4	B	109	Total	O	0	0
			109	109		
4	C	118	Total	O	0	0
			118	118		
4	D	117	Total	O	0	0
			117	117		
4	E	87	Total	O	0	0
			87	87		
4	F	98	Total	O	0	0
			98	98		
4	G	68	Total	O	0	0
			68	68		
4	H	71	Total	O	0	0
			71	71		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Burkholderia oklahomensis agglutinin (BOA)

Chain A: 




- Molecule 1: Burkholderia oklahomensis agglutinin (BOA)

Chain B: 




- Molecule 1: Burkholderia oklahomensis agglutinin (BOA)

Chain C: 




- Molecule 1: Burkholderia oklahomensis agglutinin (BOA)

Chain D: 




- Molecule 1: Burkholderia oklahomensis agglutinin (BOA)

Chain E: 



- Molecule 1: Burkholderia oklahomensis agglutinin (BOA)



• Molecule 1: Burkholderia oklahomensis agglutinin (BOA)



• Molecule 1: Burkholderia oklahomensis agglutinin (BOA)



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	119.72Å 130.58Å 149.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	18.41 – 2.40 18.41 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.5 (18.41-2.40) 97.6 (18.41-2.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.03 (at 2.40Å)	Xtriage
Refinement program	BUSTER-TNT, BUSTER 2.10.0	Depositor
R, R_{free}	0.194 , 0.229 0.189 , 0.227	Depositor DCC
R_{free} test set	4601 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	46.0	Xtriage
Anisotropy	0.742	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 37.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	17085	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, DTT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.90	0/2100	1.07	2/2858 (0.1%)
1	B	0.84	0/2100	1.12	5/2858 (0.2%)
1	C	0.83	1/2070 (0.0%)	1.08	0/2819
1	D	0.82	0/2107	1.09	5/2869 (0.2%)
1	E	0.83	0/2070	1.11	2/2819 (0.1%)
1	F	0.79	0/2086	1.06	2/2840 (0.1%)
1	G	0.77	0/2070	1.06	0/2819
1	H	0.81	2/2086 (0.1%)	1.09	3/2840 (0.1%)
All	All	0.82	3/16689 (0.0%)	1.08	19/22722 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	272	MET	SD-CE	6.17	1.95	1.79
1	C	226	HIS	CA-C	5.30	1.57	1.52
1	H	74	THR	CA-C	5.08	1.54	1.52

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	191	ASN	CA-CB-CG	9.89	122.49	112.60
1	B	191	ASN	CA-CB-CG	9.87	122.47	112.60
1	A	191	ASN	CA-CB-CG	8.59	121.19	112.60
1	E	154	GLY	CA-C-N	7.35	130.13	120.28
1	E	154	GLY	C-N-CA	7.35	130.13	120.28
1	H	191	ASN	CA-CB-CG	6.62	119.22	112.60
1	F	171	ASP	N-CA-C	-5.49	103.05	110.68
1	A	62	VAL	N-CA-CB	5.43	117.22	110.05
1	F	183	ASP	CA-CB-CG	5.28	117.88	112.60
1	B	174	VAL	CA-C-N	-5.23	116.34	122.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	174	VAL	C-N-CA	-5.23	116.34	122.14
1	H	105	ASN	CA-C-N	5.22	129.04	121.26
1	H	105	ASN	C-N-CA	5.22	129.04	121.26
1	B	105	ASN	CA-C-N	5.19	129.28	121.40
1	B	105	ASN	C-N-CA	5.19	129.28	121.40
1	D	174	VAL	CA-C-N	-5.17	115.65	122.16
1	D	174	VAL	C-N-CA	-5.17	115.65	122.16
1	D	87	GLY	CA-C-N	5.05	127.55	120.28
1	D	87	GLY	C-N-CA	5.05	127.55	120.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2046	0	1910	13	0
1	B	2046	0	1910	8	0
1	C	2016	0	1880	8	0
1	D	2050	0	1916	6	0
1	E	2016	0	1880	13	0
1	F	2029	0	1898	4	0
1	G	2016	0	1880	6	0
1	H	2029	0	1898	4	0
2	A	18	0	24	3	0
2	D	12	0	16	4	0
3	E	8	0	10	9	0
4	A	131	0	0	0	0
4	B	109	0	0	0	0
4	C	118	0	0	1	0
4	D	117	0	0	0	0
4	E	87	0	0	0	0
4	F	98	0	0	0	0
4	G	68	0	0	0	0
4	H	71	0	0	0	0
All	All	17085	0	15222	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:110:ALA:HB1	2:A:301:GOL:H2	1.33	1.05
1:A:198:GLY:HA3	2:D:302:GOL:H12	1.48	0.95
1:A:249:GLY:HA2	2:D:301:GOL:H12	1.48	0.93
1:E:209:GLY:HA3	3:E:301:DTT:H2	1.69	0.75
1:D:182:THR:HG23	2:D:302:GOL:H2	1.75	0.68
1:A:149:GLU:HG3	2:A:303:GOL:H2	1.77	0.67
1:E:210:ASN:ND2	3:E:301:DTT:H3	2.13	0.64
1:E:210:ASN:OD1	3:E:301:DTT:H12	1.97	0.64
1:E:210:ASN:H	3:E:301:DTT:H41	1.63	0.62
1:E:210:ASN:CG	3:E:301:DTT:H12	2.23	0.62
1:E:238:GLY:N	3:E:301:DTT:S1	2.69	0.60
1:G:275:PRO:O	1:G:276:GLN:HB2	2.03	0.59
1:C:17:GLN:NE2	1:C:20:GLY:O	2.36	0.58
1:A:40:ASN:O	1:A:62:VAL:HG23	2.04	0.57
1:E:104:VAL:HG22	3:E:301:DTT:C2	2.36	0.56
1:E:236:ARG:O	3:E:301:DTT:H11	2.06	0.55
1:B:25:TRP:CG	1:B:137:LYS:HZ2	2.25	0.54
1:A:221:THR:HB	1:D:221:THR:HB	1.90	0.53
1:C:152:TRP:HB3	1:C:267:ILE:HG22	1.90	0.53
1:B:93:HIS:HB3	1:C:258:SER:HB2	1.90	0.52
1:E:46:ILE:HG21	1:E:69:PHE:CD1	2.44	0.52
1:C:218:TRP:CD1	1:C:226:HIS:HE2	2.27	0.52
1:B:152:TRP:HB3	1:B:267:ILE:HG22	1.93	0.51
1:G:152:TRP:HB3	1:G:267:ILE:HG22	1.94	0.50
1:H:152:TRP:HB3	1:H:267:ILE:HG22	1.94	0.49
1:D:152:TRP:HB3	1:D:267:ILE:HG22	1.93	0.49
1:A:152:TRP:HB3	1:A:267:ILE:HG22	1.95	0.49
1:H:71:ALA:HB1	1:H:79:TYR:HB3	1.95	0.48
1:B:71:ALA:HB1	1:B:79:TYR:HB3	1.95	0.48
1:A:42:VAL:HG13	1:A:62:VAL:HG22	1.97	0.47
1:A:71:ALA:HB1	1:A:79:TYR:HB3	1.97	0.47
1:B:53:ARG:HH11	1:B:53:ARG:HG2	1.80	0.46
1:F:71:ALA:HB1	1:F:79:TYR:HB3	1.97	0.46
1:B:84:GLN:HB2	1:B:92:TRP:CD2	2.52	0.45
1:D:71:ALA:HB1	1:D:79:TYR:HB3	1.98	0.45
1:C:71:ALA:HB1	1:C:79:TYR:HB3	1.99	0.45
1:G:258:SER:HB2	1:H:93:HIS:HB3	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:226:HIS:HD2	4:C:325:HOH:O	1.99	0.45
1:G:71:ALA:HB1	1:G:79:TYR:HB3	1.98	0.45
1:E:50:ASP:HB2	1:H:196:GLY:HA3	1.99	0.44
1:A:3:LYS:O	1:A:9:ASN:HB3	2.18	0.44
1:A:266:PRO:HG3	1:C:53:ARG:HH22	1.83	0.44
1:E:71:ALA:HB1	1:E:79:TYR:HB3	2.00	0.43
1:B:84:GLN:HB2	1:B:92:TRP:CE3	2.53	0.43
1:E:104:VAL:HG22	3:E:301:DTT:H2	2.00	0.43
1:C:17:GLN:HG2	1:C:25:TRP:CZ2	2.54	0.43
1:G:192:MET:HE3	1:G:192:MET:HB2	1.87	0.43
1:F:246:ILE:HD12	1:F:255:PHE:CD2	2.54	0.43
1:E:14:VAL:HG22	1:E:138:SER:HB3	2.01	0.42
1:A:3:LYS:NZ	1:A:275:PRO:O	2.48	0.42
1:B:-2:SER:N	1:F:92:TRP:O	2.52	0.42
1:D:258:SER:OG	2:D:301:GOL:H11	2.19	0.42
1:D:93:HIS:HB3	1:F:258:SER:HB2	2.02	0.41
1:G:201:GLY:O	1:G:216:ASN:HA	2.20	0.41
1:A:103:ARG:HD3	2:A:302:GOL:H31	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	277/279 (99%)	271 (98%)	6 (2%)	0	100	100
1	B	277/279 (99%)	273 (99%)	4 (1%)	0	100	100
1	C	273/279 (98%)	268 (98%)	5 (2%)	0	100	100
1	D	278/279 (100%)	273 (98%)	5 (2%)	0	100	100
1	E	273/279 (98%)	269 (98%)	4 (2%)	0	100	100
1	F	275/279 (99%)	269 (98%)	6 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	273/279 (98%)	267 (98%)	6 (2%)	0	100	100
1	H	275/279 (99%)	269 (98%)	5 (2%)	1 (0%)	30	43
All	All	2201/2232 (99%)	2159 (98%)	41 (2%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	155	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	206/206 (100%)	205 (100%)	1 (0%)	81	91
1	B	206/206 (100%)	205 (100%)	1 (0%)	81	91
1	C	202/206 (98%)	201 (100%)	1 (0%)	81	91
1	D	207/206 (100%)	207 (100%)	0	100	100
1	E	202/206 (98%)	199 (98%)	3 (2%)	57	77
1	F	204/206 (99%)	204 (100%)	0	100	100
1	G	202/206 (98%)	201 (100%)	1 (0%)	81	91
1	H	204/206 (99%)	204 (100%)	0	100	100
All	All	1633/1648 (99%)	1626 (100%)	7 (0%)	84	92

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	62	VAL
1	B	276	GLN
1	C	17	GLN
1	E	2	THR
1	E	76	SER
1	E	178	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	G	56	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (27) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	ASN
1	A	210	ASN
1	A	276	GLN
1	B	12	GLN
1	B	210	ASN
1	B	226	HIS
1	C	12	GLN
1	C	210	ASN
1	C	276	GLN
1	D	12	GLN
1	D	16	ASN
1	D	189	ASN
1	D	191	ASN
1	E	77	ASN
1	E	93	HIS
1	E	107	GLN
1	E	140	GLN
1	F	160	HIS
1	F	189	ASN
1	F	210	ASN
1	G	83	ASN
1	G	150	ASN
1	G	189	ASN
1	G	210	ASN
1	H	12	GLN
1	H	45	ASN
1	H	151	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	D	301	-	5,5,5	0.40	0	5,5,5	0.61	0
3	DTT	E	301	-	7,7,7	1.19	0	4,8,8	1.93	2 (50%)
2	GOL	A	301	-	5,5,5	0.70	0	5,5,5	1.03	0
2	GOL	A	303	-	5,5,5	0.69	0	5,5,5	0.47	0
2	GOL	D	302	-	5,5,5	0.68	0	5,5,5	0.77	0
2	GOL	A	302	-	5,5,5	0.44	0	5,5,5	1.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	D	301	-	-	2/4/4/4	-
3	DTT	E	301	-	-	5/8/8/8	-
2	GOL	A	301	-	-	3/4/4/4	-
2	GOL	A	303	-	-	3/4/4/4	-
2	GOL	D	302	-	-	2/4/4/4	-
2	GOL	A	302	-	-	4/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	301	DTT	O2-C2-C3	-2.39	104.56	109.57
3	E	301	DTT	O3-C3-C2	-2.10	105.16	109.57

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	302	GOL	O1-C1-C2-C3
2	A	303	GOL	C1-C2-C3-O3
2	A	303	GOL	O2-C2-C3-O3
2	D	301	GOL	C1-C2-C3-O3
2	D	301	GOL	O2-C2-C3-O3
2	D	302	GOL	C1-C2-C3-O3
3	E	301	DTT	C1-C2-C3-O3
3	E	301	DTT	C1-C2-C3-C4
3	E	301	DTT	O2-C2-C3-O3
3	E	301	DTT	O2-C2-C3-C4
2	A	301	GOL	C1-C2-C3-O3
2	A	302	GOL	C1-C2-C3-O3
2	A	301	GOL	O1-C1-C2-O2
2	A	302	GOL	O1-C1-C2-O2
2	A	302	GOL	O2-C2-C3-O3
2	D	302	GOL	O2-C2-C3-O3
2	A	303	GOL	O1-C1-C2-O2
2	A	301	GOL	O1-C1-C2-C3
3	E	301	DTT	S1-C1-C2-O2

There are no ring outliers.

6 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	301	GOL	2	0
3	E	301	DTT	9	0
2	A	301	GOL	1	0
2	A	303	GOL	1	0
2	D	302	GOL	2	0
2	A	302	GOL	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	279/279 (100%)	-0.57	0 100 100	31, 42, 71, 88	0
1	B	279/279 (100%)	-0.39	1 (0%) 88 86	35, 49, 67, 94	0
1	C	275/279 (98%)	-0.45	2 (0%) 84 81	35, 47, 73, 102	0
1	D	279/279 (100%)	-0.32	0 100 100	35, 53, 81, 95	1 (0%)
1	E	275/279 (98%)	-0.09	4 (1%) 72 68	33, 55, 87, 102	0
1	F	276/279 (98%)	-0.34	4 (1%) 73 69	32, 54, 82, 112	1 (0%)
1	G	275/279 (98%)	0.04	1 (0%) 88 86	44, 62, 92, 112	0
1	H	276/279 (98%)	-0.09	1 (0%) 88 86	40, 64, 85, 99	1 (0%)
All	All	2214/2232 (99%)	-0.28	13 (0%) 85 83	31, 53, 82, 112	3 (0%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	152	TRP	3.5
1	C	153	GLY	2.7
1	C	155	SER	2.6
1	B	85	TRP	2.4
1	E	24	PRO	2.2
1	G	224	PRO	2.2
1	F	218	TRP	2.2
1	E	120	ALA	2.1
1	F	105	ASN	2.1
1	E	20	GLY	2.1
1	E	105	ASN	2.1
1	F	2	THR	2.0
1	F	152	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	A	301	6/6	0.59	0.23	46,50,55,56	6
2	GOL	D	301	6/6	0.71	0.16	57,58,59,60	6
2	GOL	D	302	6/6	0.77	0.20	38,41,43,45	6
3	DTT	E	301	8/8	0.78	0.25	45,49,51,55	8
2	GOL	A	303	6/6	0.80	0.20	36,46,47,48	6
2	GOL	A	302	6/6	0.86	0.11	62,67,68,71	0

6.5 Other polymers [i](#)

There are no such residues in this entry.