



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 12:39 AM UTC

PDB ID : 4GUT / pdb_00004gut
Title : Crystal structure of LSD2-NPAC
Authors : Chen, F.; Dong, Z.; Fang, J.; Yang, Y.; Li, Z.; Xu, Y.; Yang, H.; Wang, P.;
Fang, R.; Shi, Y.; Xu, Y.
Deposited on : 2012-08-29
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	776	
2	B	124	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	PGE	B	301	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6390 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lysine-specific histone demethylase 1B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	740	Total	C	N	O	S	0	2	0
			5861	3740	998	1081	42			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	PRO	-	expression tag	UNP Q8NB78
A	48	LEU	-	expression tag	UNP Q8NB78
A	49	GLY	-	expression tag	UNP Q8NB78
A	50	SER	-	expression tag	UNP Q8NB78


- Molecule 2 is a protein called Putative oxidoreductase GLYR1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	12	Total	C	N	O	0	0	0
			105	69	19	17			

There are 7 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	145	PRO	-	expression tag	UNP Q49A26
B	146	LEU	-	expression tag	UNP Q49A26
B	147	GLY	-	expression tag	UNP Q49A26
B	148	SER	-	expression tag	UNP Q49A26
B	149	PRO	-	expression tag	UNP Q49A26
B	150	GLU	-	expression tag	UNP Q49A26
B	151	PHE	-	expression tag	UNP Q49A26

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0
			53	27	9	15	2	

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	3	Total 3	Zn 3	0	0

- Molecule 6 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total 10	C 6	O 4	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	342	Total 342	O 342	0	0
7	B	4	Total 4	O 4	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.00Å 89.76Å 86.69Å 90.00° 104.96° 90.00°	Depositor
Resolution (Å)	35.92 – 2.00 35.92 – 2.00	Depositor EDS
% Data completeness (in resolution range)	90.2 (35.92-2.00) 95.1 (35.92-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.67 (at 2.00Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, R_{free}	0.196 , 0.207 0.197 , 0.209	Depositor DCC
R_{free} test set	2995 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å ²)	28.2	Xtriage
Anisotropy	0.279	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6390	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, PGE, GOL, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	0/6016	0.80	0/8149
2	B	1.18	0/110	0.71	0/149
All	All	0.64	0/6126	0.79	0/8298

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5861	0	5777	57	0
2	B	105	0	91	24	0
3	A	53	0	29	3	0
4	A	12	0	16	3	0
5	A	3	0	0	0	0
6	B	10	0	14	25	0
7	A	342	0	0	1	0
7	B	4	0	0	0	0
All	All	6390	0	5927	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:217:PHE:CD1	6:B:301:PGE:C2	2.32	1.13
2:B:217:PHE:CD1	6:B:301:PGE:H2	1.83	1.11
2:B:217:PHE:HD1	6:B:301:PGE:H2	0.98	1.10
2:B:217:PHE:HD1	6:B:301:PGE:C2	1.71	1.03
2:B:217:PHE:CE1	6:B:301:PGE:C3	2.43	1.01
2:B:217:PHE:HB3	6:B:301:PGE:O1	1.63	0.99
2:B:217:PHE:HE1	6:B:301:PGE:C3	1.82	0.91
1:A:422:TRP:CZ3	4:A:903:GOL:H2	2.12	0.85
1:A:149:LYS:HD2	1:A:189:GLU:HB2	1.59	0.83
2:B:217:PHE:CD1	6:B:301:PGE:O2	2.32	0.83
1:A:454:LEU:HD21	1:A:585:LYS:HG3	1.60	0.82
1:A:422:TRP:HZ3	4:A:903:GOL:H2	1.44	0.81
2:B:217:PHE:HB3	6:B:301:PGE:C1	2.10	0.80
1:A:51:ARG:HG2	1:A:51:ARG:HH11	1.50	0.77
2:B:217:PHE:CE1	6:B:301:PGE:H32	2.20	0.75
1:A:352:ILE:O	1:A:356:MET:HG3	1.88	0.73
2:B:217:PHE:CE1	6:B:301:PGE:H3	2.24	0.72
2:B:217:PHE:HE1	6:B:301:PGE:H32	1.53	0.71
1:A:495:SER:HA	1:A:498:ARG:HG2	1.76	0.68
2:B:217:PHE:HE1	6:B:301:PGE:H3	1.56	0.67
1:A:306:MET:HE3	1:A:345:CYS:HA	1.76	0.67
2:B:217:PHE:CE1	6:B:301:PGE:O2	2.46	0.67
1:A:766:ALA:HA	3:A:901:FAD:HM81	1.78	0.66
1:A:493:VAL:HG11	1:A:516:ILE:HG21	1.79	0.65
1:A:285:ARG:HD3	6:B:301:PGE:H1	1.79	0.64
1:A:498:ARG:HD2	7:A:1440:HOH:O	1.97	0.64
1:A:550:ASN:HB3	1:A:553:GLN:HE22	1.63	0.63
1:A:76:ASN:H	1:A:76:ASN:HD22	1.45	0.62
2:B:217:PHE:CD1	6:B:301:PGE:C3	2.83	0.61
2:B:217:PHE:CB	6:B:301:PGE:C1	2.79	0.61
1:A:725:ASP:H	1:A:728:GLN:HE21	1.49	0.60
1:A:53:CYS:SG	1:A:84:HIS:HB2	2.41	0.59
1:A:114:LYS:HG3	1:A:125:PRO:HB2	1.85	0.58
1:A:551:LEU:HA	1:A:554:VAL:HG13	1.86	0.57
1:A:285:ARG:HG2	6:B:301:PGE:O4	2.06	0.56
1:A:756:ARG:N	4:A:903:GOL:O1	2.30	0.54
1:A:204:MET:HA	1:A:204:MET:HE2	1.91	0.53
1:A:99:TYR:CE2	1:A:131:MET:HE1	2.45	0.52
2:B:217:PHE:CB	6:B:301:PGE:H12	2.39	0.52
2:B:217:PHE:CG	6:B:301:PGE:H12	2.44	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:70:SER:HB2	1:A:98:HIS:HB2	1.90	0.52
1:A:722:ARG:HG3	1:A:723:THR:HG23	1.92	0.52
1:A:550:ASN:HD22	1:A:716:GLU:HG3	1.75	0.52
2:B:217:PHE:CD1	6:B:301:PGE:C1	2.92	0.52
2:B:214:ASP:HB2	2:B:216:HIS:HD2	1.74	0.51
1:A:51:ARG:HG2	1:A:51:ARG:NH1	2.24	0.51
1:A:205:LEU:CD1	1:A:207:LEU:O	2.60	0.50
1:A:471:GLY:HA2	1:A:739:GLU:O	2.12	0.49
1:A:72:ARG:O	1:A:94:GLU:HG3	2.12	0.48
1:A:725:ASP:H	1:A:728:GLN:NE2	2.09	0.48
1:A:584[B]:GLU:HG2	1:A:585:LYS:N	2.29	0.47
1:A:316:ALA:O	1:A:320:THR:HG23	2.15	0.46
1:A:736:THR:HG22	1:A:740:LEU:HD22	1.98	0.46
1:A:493:VAL:CG1	1:A:516:ILE:HG21	2.45	0.46
1:A:93:ASN:O	1:A:96:PHE:HB3	2.16	0.46
1:A:410:VAL:HG13	1:A:592:ILE:HG12	1.97	0.45
1:A:151:ARG:HD2	1:A:169:CYS:SG	2.57	0.45
1:A:493:VAL:HG13	1:A:516:ILE:HD13	1.98	0.45
1:A:51:ARG:HH11	1:A:51:ARG:CG	2.26	0.45
2:B:214:ASP:HA	2:B:215:PRO:HD3	1.83	0.44
1:A:114:LYS:O	1:A:118:THR:HB	2.18	0.44
1:A:276:ASN:H	6:B:301:PGE:H52	1.81	0.44
1:A:102:SER:HA	1:A:107:TYR:CG	2.52	0.44
1:A:803:GLN:O	3:A:901:FAD:H1'2	2.19	0.43
1:A:454:LEU:HD13	1:A:581:VAL:HG12	1.99	0.43
1:A:626:THR:HG22	1:A:793:ALA:HB3	1.99	0.43
1:A:92:CYS:SG	1:A:94:GLU:HG2	2.59	0.43
1:A:436:ALA:HA	3:A:901:FAD:C4X	2.49	0.42
1:A:149:LYS:CD	1:A:189:GLU:HB2	2.41	0.42
1:A:205:LEU:HD11	1:A:207:LEU:O	2.18	0.42
1:A:306:MET:CE	1:A:345:CYS:HA	2.48	0.42
2:B:217:PHE:CD1	6:B:301:PGE:H32	2.52	0.42
2:B:217:PHE:CB	6:B:301:PGE:O1	2.51	0.42
2:B:217:PHE:CG	6:B:301:PGE:C1	3.03	0.42
1:A:83:TYR:O	1:A:90:HIS:HA	2.19	0.42
1:A:66:PHE:CD2	1:A:66:PHE:C	2.98	0.41
1:A:71:GLU:H	1:A:71:GLU:CD	2.28	0.41
1:A:352:ILE:O	1:A:356:MET:CG	2.63	0.41
1:A:100:TYR:HD2	1:A:101:ARG:HE	1.69	0.41
1:A:130:PHE:HD2	1:A:131:MET:HE2	1.85	0.41
1:A:801:PHE:O	1:A:807:GLY:HA3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	736/776 (95%)	718 (98%)	18 (2%)	0	100	100
2	B	10/124 (8%)	9 (90%)	1 (10%)	0	100	100
All	All	746/900 (83%)	727 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	637/662 (96%)	601 (94%)	36 (6%)	18	15
2	B	12/106 (11%)	12 (100%)	0	100	100
All	All	649/768 (84%)	613 (94%)	36 (6%)	19	17

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	ARG
1	A	71	GLU
1	A	76	ASN
1	A	80	SER
1	A	85	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	94	GLU
1	A	101	ARG
1	A	118	THR
1	A	120	ASN
1	A	122	LYS
1	A	146	GLU
1	A	151	ARG
1	A	168	ARG
1	A	183	ASP
1	A	194	LEU
1	A	283	CYS
1	A	294	LEU
1	A	296	GLU
1	A	308	LEU
1	A	344	ARG
1	A	361	LEU
1	A	382	ASN
1	A	410	VAL
1	A	467	LEU
1	A	493	VAL
1	A	542	ASN
1	A	554	VAL
1	A	574	LEU
1	A	576	THR
1	A	585	LYS
1	A	680	PHE
1	A	693	LEU
1	A	707	SER
1	A	721	VAL
1	A	730	LEU
1	A	740	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	76	ASN
1	A	103	HIS
1	A	174	ASN
1	A	274	GLN
1	A	276	ASN
1	A	402	HIS
1	A	542	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	553	GLN
1	A	593	GLN
1	A	706	HIS
1	A	728	GLN
2	B	216	HIS
2	B	218	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	A	903	-	5,5,5	0.39	0	5,5,5	0.58	0
4	GOL	A	902	-	5,5,5	0.21	0	5,5,5	0.47	0
3	FAD	A	901	-	58,58,58	2.28	19 (32%)	85,89,89	1.51	14 (16%)
6	PGE	B	301	-	9,9,9	0.52	0	8,8,8	1.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	903	-	-	0/4/4/4	-
4	GOL	A	902	-	-	2/4/4/4	-
3	FAD	A	901	-	-	0/34/50/50	0/6/6/6
6	PGE	B	301	-	-	2/7/7/7	-

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	901	FAD	C5A-N7A	-5.80	1.28	1.39
3	A	901	FAD	P-O2P	-4.67	1.33	1.55
3	A	901	FAD	C8A-N9A	-4.48	1.29	1.37
3	A	901	FAD	PA-O2A	-4.19	1.36	1.55
3	A	901	FAD	O4B-C4B	-4.05	1.36	1.45
3	A	901	FAD	P-O1P	-3.65	1.38	1.50
3	A	901	FAD	C4A-N9A	-3.63	1.30	1.37
3	A	901	FAD	C5X-N5	-3.52	1.33	1.39
3	A	901	FAD	O3'-C3'	-3.36	1.34	1.43
3	A	901	FAD	PA-O1A	-3.34	1.39	1.50
3	A	901	FAD	O5'-C5'	-3.27	1.32	1.44
3	A	901	FAD	O4'-C4'	-3.06	1.36	1.43
3	A	901	FAD	O2'-C2'	-3.05	1.36	1.43
3	A	901	FAD	PA-O3P	-2.88	1.56	1.59
3	A	901	FAD	O3B-C3B	-2.73	1.36	1.43
3	A	901	FAD	PA-O5B	-2.52	1.49	1.59
3	A	901	FAD	C1'-C2'	-2.36	1.49	1.52
3	A	901	FAD	C4X-N5	2.34	1.35	1.30
3	A	901	FAD	O2-C2	-2.30	1.19	1.24

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	901	FAD	C5A-C4A-N3A	-5.17	119.60	126.72
3	A	901	FAD	N3A-C2A-N1A	-3.40	123.44	128.58
3	A	901	FAD	C4A-C5A-N7A	-3.36	106.74	110.58
3	A	901	FAD	N3A-C4A-N9A	3.21	132.63	127.17
3	A	901	FAD	C2A-N3A-C4A	3.12	119.44	111.83
3	A	901	FAD	C9A-C5X-N5	-3.08	119.18	122.45
3	A	901	FAD	C5A-N7A-C8A	3.04	108.23	103.45
3	A	901	FAD	O4B-C1B-C2B	-2.60	101.06	106.62
3	A	901	FAD	C5X-C9A-N10	2.45	120.18	117.97

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	901	FAD	O5'-C5'-C4'	2.43	115.84	109.36
3	A	901	FAD	N9A-C8A-N7A	-2.40	110.52	113.94
3	A	901	FAD	O5B-C5B-C4B	2.15	116.32	108.99
3	A	901	FAD	C10-C4X-N5	-2.11	120.50	124.81
3	A	901	FAD	C10-N1-C2	2.09	121.38	116.85

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	902	GOL	O1-C1-C2-C3
4	A	902	GOL	O1-C1-C2-O2
6	B	301	PGE	O2-C3-C4-O3
6	B	301	PGE	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 31 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	903	GOL	3	0
3	A	901	FAD	3	0
6	B	301	PGE	25	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	740/776 (95%)	0.29	44 (5%) 28 27	18, 35, 65, 93	2 (0%)
2	B	12/124 (9%)	0.94	3 (25%) 2 1	27, 36, 76, 84	0
All	All	752/900 (83%)	0.30	47 (6%) 26 24	18, 35, 65, 93	2 (0%)

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	48	LEU	5.8
1	A	83	TYR	4.8
1	A	100	TYR	3.9
1	A	47	PRO	3.8
1	A	96	PHE	3.6
1	A	82	TRP	3.4
1	A	111	THR	3.3
2	B	217	PHE	3.2
1	A	206	ILE	3.2
1	A	57	GLY	3.1
1	A	200	TRP	3.0
1	A	56	ALA	2.9
1	A	107	TYR	2.9
1	A	199	HIS	2.8
1	A	112	THR	2.8
1	A	101	ARG	2.8
1	A	59	THR	2.8
1	A	103	HIS	2.8
1	A	173	PRO	2.8
1	A	49	GLY	2.7
1	A	91	PHE	2.7
1	A	79	THR	2.7
2	B	215	PRO	2.7
1	A	87	CYS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	276	ASN	2.6
1	A	122	LYS	2.5
1	A	118	THR	2.5
1	A	116	ILE	2.5
1	A	207	LEU	2.4
1	A	124	GLU	2.4
1	A	526	ILE	2.3
1	A	174	ASN	2.3
1	A	117	TRP	2.3
1	A	50	SER	2.3
1	A	120	ASN	2.2
2	B	216	HIS	2.2
1	A	235	THR	2.2
1	A	264	GLY	2.2
1	A	371	ALA	2.2
1	A	99	TYR	2.2
1	A	110	TYR	2.2
1	A	62	CYS	2.2
1	A	224	TYR	2.1
1	A	114	LYS	2.1
1	A	84	HIS	2.1
1	A	85	LEU	2.1
1	A	821	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	PGE	B	301	10/10	0.66	0.16	55,61,64,64	0
4	GOL	A	903	6/6	0.85	0.17	32,37,37,42	0
4	GOL	A	902	6/6	0.85	0.13	32,33,36,43	0
3	FAD	A	901	53/53	0.96	0.06	16,21,26,27	0
5	ZN	A	904	1/1	0.97	0.07	47,47,47,47	0
5	ZN	A	906	1/1	0.98	0.03	44,44,44,44	0
5	ZN	A	905	1/1	0.99	0.04	37,37,37,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.