



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 10:13 AM UTC

PDB ID : 5GVN / pdb_00005gvn
Title : Plasmodium vivax SHMT bound with PLP-glycine and GS653
Authors : Chitnumsub, P.; Jaruwat, A.; Leartsakulpanich, U.; Schwertz, G.
Deposited on : 2016-09-06
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	442	
1	B	442	
1	C	442	

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 11053 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine hydroxymethyltransferase, putative.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	442	Total	C	N	O	S	0	0	0
			3458	2186	600	655	17			
1	B	442	Total	C	N	O	S	0	0	0
			3458	2186	600	655	17			
1	C	442	Total	C	N	O	S	0	0	0
			3458	2186	600	655	17			

- Molecule 2 is N-GLYCINE-[3-HYDROXY-2-METHYL-5-PHOSPHONOOXYMETHYL-PYRIDIN-4-YL-METHANE] (CCD ID: PLG) (formula: C₁₀H₁₅N₂O₇P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			20	10	2	7	1		
2	B	1	Total	C	N	O	P	0	0
			20	10	2	7	1		
2	C	1	Total	C	N	O	P	0	0
			20	10	2	7	1		

- Molecule 3 is 3-[3-[3-[(4 {S})-6-azanyl-5-cyano-3-methyl-4-propan-2-yl-2 {H}-pyrano [2,3-c]pyrazol-4-yl]-5-fluoranyl-phenyl]phenyl]propanoic acid (CCD ID: G6F) (formula: $C_{26}H_{25}FN_4O_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	0	0
			34	26	1	4	3		
3	B	1	Total	C	F	N	O	0	0
			34	26	1	4	3		
3	C	1	Total	C	F	N	O	0	0
			34	26	1	4	3		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		
4	C	1	Total	Cl	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	172	Total	O	0	0
			172	172		
5	B	169	Total	O	0	0
			169	169		
5	C	174	Total	O	0	0
			174	174		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serine hydroxymethyltransferase, putative



- Molecule 1: Serine hydroxymethyltransferase, putative



- Molecule 1: Serine hydroxymethyltransferase, putative



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	101.85Å 58.80Å 235.61Å 90.00° 89.97° 90.00°	Depositor
Resolution (Å)	50.01 – 2.30 50.01 – 2.30	Depositor EDS
% Data completeness (in resolution range)	97.5 (50.01-2.30) 97.7 (50.01-2.30)	Depositor EDS
R_{merge}	0.02	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	9.61 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.219 , 0.282 (Not available) , 0.269	Depositor DCC
R_{free} test set	6186 reflections (9.83%)	wwPDB-VP
Wilson B-factor (Å ²)	27.4	Xtriage
Anisotropy	0.053	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 17.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.020 for -1/2*h-3/2*k,-1/2*h+1/2*k,-l 0.020 for -1/2*h+3/2*k,1/2*h+1/2*k,-l 0.477 for 1/2*h-3/2*k,-1/2*h-1/2*k,-l 0.478 for 1/2*h+3/2*k,1/2*h-1/2*k,-l 0.018 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11053	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: G6F, PLG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.77	0/3521	0.94	0/4754
1	B	0.77	0/3521	0.94	1/4754 (0.0%)
1	C	0.77	0/3521	0.94	0/4754
All	All	0.77	0/10563	0.94	1/14262 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	125	CYS	N-CA-C	5.14	118.33	111.75

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3458	0	3477	20	0
1	B	3458	0	3477	23	0
1	C	3458	0	3477	24	0
2	A	20	0	11	4	0
2	B	20	0	11	1	0
2	C	20	0	11	2	0
3	A	34	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	34	0	0	0	0
3	C	34	0	0	0	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	A	172	0	0	0	0
5	B	169	0	0	1	0
5	C	174	0	0	1	0
All	All	11053	0	10464	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:167:ARG:NH1	1:A:200:GLU:OE1	2.22	0.71
1:C:383:LYS:HB2	1:C:385:LYS:HE2	1.82	0.61
1:B:341:LYS:NZ	5:B:601:HOH:O	2.37	0.58
1:C:193:GLN:O	1:C:197:ILE:HG13	2.04	0.57
1:A:327:LEU:HD12	1:A:327:LEU:C	2.31	0.56
1:B:174:LYS:HD2	1:B:201:VAL:CG1	2.36	0.54
1:C:427:GLN:HE21	1:C:427:GLN:HA	1.73	0.54
1:B:120:MET:SD	1:B:169:MET:HE3	2.49	0.52
1:A:112:LEU:HD11	1:A:230:VAL:HG21	1.91	0.52
1:B:195:ARG:HB2	1:B:227:HIS:HB3	1.92	0.51
1:A:109:LEU:O	1:A:113:VAL:HG22	2.11	0.50
1:C:122:MET:HE1	1:C:161:VAL:HG22	1.93	0.50
1:B:235:THR:O	1:B:241:GLY:N	2.44	0.49
1:C:195:ARG:HB2	1:C:227:HIS:HB3	1.94	0.49
1:A:353:LEU:C	1:A:353:LEU:HD12	2.37	0.49
1:C:226:LEU:HB2	1:C:227:HIS:CD2	2.48	0.48
1:A:229:ASP:OD1	1:A:253:ARG:HD2	2.13	0.48
1:B:234:THR:HB	1:B:236:HIS:CE1	2.49	0.47
1:B:385:LYS:HE3	1:B:385:LYS:H	1.79	0.47
1:C:237:LYS:CE	2:C:501:PLG:H4A2	2.44	0.47
1:C:353:LEU:HD12	1:C:353:LEU:C	2.40	0.47
1:A:249:PHE:HB3	1:A:261:ILE:HD12	1.97	0.46
1:B:6:PRO:O	1:B:10:ILE:HG22	2.15	0.46
1:C:235:THR:O	1:C:241:GLY:N	2.46	0.46
1:C:7:LEU:HA	1:C:10:ILE:HG22	1.98	0.46
1:C:122:MET:HE3	1:C:153:TYR:CE1	2.51	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:234:THR:HB	1:A:236:HIS:CE1	2.51	0.45
1:B:113:VAL:HG12	1:B:176:LYS:HB3	1.97	0.45
1:B:153:TYR:HB3	1:B:169:MET:HE1	1.98	0.45
1:A:183:THR:HB	2:A:501:PLG:O3	2.16	0.45
1:B:113:VAL:CG1	1:B:177:VAL:HG23	2.47	0.45
1:B:385:LYS:H	1:B:385:LYS:CE	2.29	0.45
1:B:398:ILE:O	1:B:402:VAL:HG23	2.17	0.44
1:C:113:VAL:CG1	1:C:177:VAL:HG23	2.48	0.44
1:C:427:GLN:HA	1:C:427:GLN:NE2	2.32	0.44
1:A:237:LYS:HE2	2:A:501:PLG:H4A2	1.99	0.44
1:C:229:ASP:OD1	1:C:253:ARG:HD2	2.18	0.44
1:B:153:TYR:HB3	1:B:169:MET:CE	2.48	0.43
1:A:237:LYS:CE	2:A:501:PLG:H4A2	2.49	0.43
1:A:416:LYS:O	1:A:420:PRO:HD3	2.19	0.43
1:B:7:LEU:HA	1:B:10:ILE:CG2	2.47	0.43
1:B:237:LYS:CE	2:B:501:PLG:H4A2	2.48	0.43
1:C:183:THR:HB	2:C:501:PLG:O3	2.19	0.43
1:A:41:ALA:HB2	1:B:10:ILE:HG21	2.00	0.43
1:A:184:SER:HA	1:A:327:LEU:HD21	1.99	0.43
1:A:226:LEU:HB2	1:A:227:HIS:CD2	2.53	0.43
1:B:122:MET:HE1	1:B:161:VAL:HG22	2.01	0.43
1:A:122:MET:HE3	1:A:153:TYR:CE1	2.54	0.43
1:C:102:SER:HA	1:C:131:THR:HG21	2.01	0.43
1:C:251:LYS:HG2	5:C:764:HOH:O	2.19	0.42
1:C:374:THR:N	1:C:375:PRO:CD	2.82	0.42
1:C:234:THR:HB	1:C:236:HIS:CE1	2.55	0.42
1:C:97:GLN:N	1:C:98:PRO:CD	2.83	0.42
1:C:291:PRO:O	1:C:295:GLU:HG3	2.19	0.42
1:B:195:ARG:HD2	1:B:227:HIS:O	2.20	0.41
1:C:120:MET:HG2	1:C:169:MET:HE3	2.01	0.41
1:A:400:ILE:HG21	1:A:425:LEU:HD21	2.02	0.41
1:B:59:PRO:HA	1:B:71:ASP:OD1	2.19	0.41
1:A:195:ARG:HD2	1:A:227:HIS:O	2.20	0.41
1:B:3:ASN:ND2	1:B:10:ILE:HD13	2.36	0.41
1:B:374:THR:N	1:B:375:PRO:CD	2.84	0.41
1:B:226:LEU:HB2	1:B:227:HIS:CD2	2.56	0.41
1:C:398:ILE:O	1:C:402:VAL:HG23	2.21	0.41
1:A:374:THR:N	1:A:375:PRO:CD	2.84	0.40
1:C:89:ASP:OD1	1:C:89:ASP:C	2.64	0.40
1:A:34:SER:OG	2:A:501:PLG:O	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	440/442 (100%)	429 (98%)	10 (2%)	1 (0%)	43	55
1	B	440/442 (100%)	424 (96%)	15 (3%)	1 (0%)	43	55
1	C	440/442 (100%)	427 (97%)	13 (3%)	0	100	100
All	All	1320/1326 (100%)	1280 (97%)	38 (3%)	2 (0%)	43	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	237	LYS
1	A	237	LYS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	381/381 (100%)	358 (94%)	23 (6%)	17	25
1	B	381/381 (100%)	356 (93%)	25 (7%)	15	21
1	C	381/381 (100%)	356 (93%)	25 (7%)	15	21
All	All	1143/1143 (100%)	1070 (94%)	73 (6%)	16	23

All (73) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	54	TYR
1	A	62	ARG
1	A	86	ASN
1	A	90	GLU
1	A	100	SER
1	A	136	ASP
1	A	138	LYS
1	A	183	THR
1	A	201	VAL
1	A	240	ARG
1	A	244	SER
1	A	252	LYS
1	A	254	ASN
1	A	260	LYS
1	A	327	LEU
1	A	340	SER
1	A	352	SER
1	A	362	VAL
1	A	385	LYS
1	A	406	GLU
1	A	427	GLN
1	A	430	GLN
1	B	21	GLU
1	B	54	TYR
1	B	61	LYS
1	B	62	ARG
1	B	100	SER
1	B	138	LYS
1	B	139	LYS
1	B	174	LYS
1	B	183	THR
1	B	184	SER
1	B	201	VAL
1	B	240	ARG
1	B	244	SER
1	B	252	LYS
1	B	253	ARG
1	B	254	ASN
1	B	261	ILE
1	B	327	LEU
1	B	352	SER
1	B	355	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	363	ASP
1	B	385	LYS
1	B	411	LYS
1	B	427	GLN
1	B	430	GLN
1	C	21	GLU
1	C	54	TYR
1	C	62	ARG
1	C	100	SER
1	C	139	LYS
1	C	151	LYS
1	C	167	ARG
1	C	168	GLU
1	C	172	SER
1	C	174	LYS
1	C	183	THR
1	C	184	SER
1	C	196	GLN
1	C	197	ILE
1	C	252	LYS
1	C	261	ILE
1	C	327	LEU
1	C	352	SER
1	C	362	VAL
1	C	385	LYS
1	C	406	GLU
1	C	411	LYS
1	C	413	VAL
1	C	427	GLN
1	C	430	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	356	ASN
1	A	426	GLN
1	B	9	GLN
1	B	315	GLN
1	B	427	GLN
1	C	227	HIS
1	C	315	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	405	GLN
1	C	427	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	G6F	B	502	-	35,37,37	2.96	4 (11%)	48,55,55	2.85	15 (31%)
2	PLG	A	501	-	20,20,20	2.58	3 (15%)	26,28,28	3.12	13 (50%)
2	PLG	B	501	-	20,20,20	2.66	4 (20%)	26,28,28	3.10	13 (50%)
3	G6F	A	502	-	35,37,37	2.85	4 (11%)	48,55,55	2.70	11 (22%)
2	PLG	C	501	-	20,20,20	2.77	3 (15%)	26,28,28	2.80	10 (38%)
3	G6F	C	502	-	35,37,37	3.06	4 (11%)	48,55,55	2.74	13 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	G6F	B	502	-	-	1/21/43/43	0/4/4/4
2	PLG	A	501	-	-	3/12/12/12	0/1/1/1
2	PLG	B	501	-	-	3/12/12/12	0/1/1/1
3	G6F	A	502	-	-	1/21/43/43	0/4/4/4
2	PLG	C	501	-	-	3/12/12/12	0/1/1/1
3	G6F	C	502	-	-	1/21/43/43	0/4/4/4

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	502	G6F	C8-C3	-12.35	1.26	1.42
3	B	502	G6F	C8-C3	-11.86	1.27	1.42
3	A	502	G6F	C8-C3	-11.42	1.27	1.42
3	C	502	G6F	O1-C2	8.71	1.45	1.37
2	C	501	PLG	C3-C2	8.54	1.49	1.41
3	B	502	G6F	C10-C5	8.31	1.50	1.38
3	C	502	G6F	C10-C5	8.19	1.49	1.38
2	B	501	PLG	C3-C2	8.08	1.49	1.41
3	B	502	G6F	O1-C2	7.99	1.45	1.37
3	A	502	G6F	C10-C5	7.93	1.49	1.38
3	A	502	G6F	O1-C2	7.74	1.44	1.37
2	A	501	PLG	C3-C2	7.44	1.48	1.41
2	A	501	PLG	C5-C4	6.61	1.49	1.40
2	C	501	PLG	C5-C4	6.50	1.49	1.40
2	B	501	PLG	C5-C4	6.41	1.49	1.40
2	C	501	PLG	C3-C4	4.70	1.46	1.40
2	A	501	PLG	C3-C4	4.19	1.46	1.40
2	B	501	PLG	C3-C4	3.78	1.45	1.40
3	B	502	G6F	N11-N12	2.77	1.42	1.36
3	C	502	G6F	N11-N12	2.45	1.41	1.36
3	A	502	G6F	N11-N12	2.41	1.41	1.36
2	B	501	PLG	OXT-C	-2.31	1.23	1.30

All (75) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	502	G6F	O1-C6-C5	-11.98	118.87	125.76
3	A	502	G6F	O1-C6-C5	-11.08	119.39	125.76
3	C	502	G6F	O1-C6-C5	-10.97	119.45	125.76
3	B	502	G6F	O1-C6-N12	9.45	126.44	119.13
3	C	502	G6F	O1-C6-N12	9.39	126.39	119.13

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	G6F	O1-C6-N12	9.30	126.33	119.13
2	C	501	PLG	OP4-C5A-C5	9.12	126.44	109.36
2	A	501	PLG	OP4-C5A-C5	8.25	124.81	109.36
2	B	501	PLG	OP4-C5A-C5	7.96	124.28	109.36
3	C	502	G6F	O1-C2-N7	6.48	114.61	109.99
2	A	501	PLG	C4A-N-CA	5.74	119.41	112.72
3	B	502	G6F	O1-C2-N7	5.72	114.07	109.99
2	A	501	PLG	C4A-C4-C5	5.68	125.94	119.75
2	B	501	PLG	C4A-C4-C5	5.48	125.71	119.75
3	A	502	G6F	O1-C2-N7	5.22	113.71	109.99
2	C	501	PLG	C4A-N-CA	5.16	118.72	112.72
2	B	501	PLG	C4A-N-CA	5.05	118.60	112.72
2	B	501	PLG	C5A-C5-C4	-4.86	112.43	122.67
2	A	501	PLG	C5A-C5-C4	-4.64	112.89	122.67
3	C	502	G6F	C3-C2-N7	-4.37	123.18	127.50
2	B	501	PLG	C6-C5-C4	4.29	121.30	118.06
2	B	501	PLG	C5A-C5-C6	4.23	126.26	119.36
2	A	501	PLG	C5A-C5-C6	4.17	126.15	119.36
2	C	501	PLG	OP4-P-OP1	-4.06	95.46	106.44
2	A	501	PLG	C6-C5-C4	3.83	120.95	118.06
2	C	501	PLG	C3-C4-C5	-3.82	115.27	118.73
3	B	502	G6F	C3-C2-N7	-3.81	123.73	127.50
2	C	501	PLG	C4A-C4-C5	3.79	123.88	119.75
3	B	502	G6F	C8-C3-C2	3.74	122.48	117.67
2	B	501	PLG	C3-C4-C5	-3.53	115.53	118.73
3	A	502	G6F	C3-C2-N7	-3.46	124.08	127.50
2	A	501	PLG	C3-C4-C5	-3.43	115.62	118.73
3	A	502	G6F	C16-C17-C18	3.23	121.82	118.35
2	B	501	PLG	C4-C4A-N	3.21	117.42	111.50
3	B	502	G6F	C6-O1-C2	3.04	120.21	115.95
3	B	502	G6F	C16-C17-C18	3.02	121.60	118.35
2	A	501	PLG	C4-C4A-N	2.99	117.02	111.50
2	B	501	PLG	OP2-P-OP4	-2.94	99.01	106.67
3	A	502	G6F	C6-O1-C2	2.93	120.06	115.95
3	C	502	G6F	C16-C17-C18	2.92	121.49	118.35
3	B	502	G6F	C14-C4-C3	-2.90	105.47	110.22
2	B	501	PLG	O3-C3-C2	2.89	123.58	117.58
3	A	502	G6F	C36-C14-C4	2.81	115.23	112.31
3	B	502	G6F	C36-C14-C4	2.80	115.22	112.31
3	C	502	G6F	C14-C4-C3	-2.79	105.66	110.22
3	B	502	G6F	C17-C18-C19	-2.78	120.12	123.50
3	C	502	G6F	C17-C18-C19	-2.78	120.13	123.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	502	G6F	C10-C5-C6	-2.75	102.01	103.38
3	A	502	G6F	C17-C18-C19	-2.74	120.17	123.50
3	B	502	G6F	C10-C5-C6	-2.70	102.04	103.38
2	A	501	PLG	O3-C3-C2	2.67	123.11	117.58
2	C	501	PLG	C6-C5-C4	2.64	120.06	118.06
2	C	501	PLG	C4-C4A-N	2.63	116.35	111.50
3	A	502	G6F	C10-C5-C6	-2.59	102.09	103.38
2	C	501	PLG	OP3-P-OP4	2.57	113.38	106.67
2	B	501	PLG	OP3-P-OP2	2.56	117.40	107.80
2	A	501	PLG	C6-N1-C2	2.55	123.82	119.20
2	B	501	PLG	C6-N1-C2	2.53	123.78	119.20
3	C	502	G6F	C36-C14-C4	2.43	114.83	112.31
3	C	502	G6F	C13-C10-C5	-2.43	128.88	132.49
3	B	502	G6F	C13-C10-N11	2.42	125.63	121.81
3	A	502	G6F	C15-C14-C4	2.42	114.82	112.31
3	C	502	G6F	C13-C10-N11	2.39	125.58	121.81
3	C	502	G6F	C6-O1-C2	2.38	119.28	115.95
3	B	502	G6F	C5-C10-N11	-2.31	105.00	106.04
3	A	502	G6F	C8-C3-C2	2.28	120.61	117.67
3	C	502	G6F	C15-C14-C4	2.27	114.67	112.31
2	A	501	PLG	OP4-P-OP1	-2.25	100.36	106.44
2	C	501	PLG	O3-C3-C2	2.17	122.07	117.58
2	B	501	PLG	C2A-C2-N1	2.16	121.72	117.64
2	A	501	PLG	OXT-C-CA	2.15	120.98	112.81
3	B	502	G6F	C13-C10-C5	-2.11	129.36	132.49
2	A	501	PLG	C2A-C2-N1	2.10	121.59	117.64
3	B	502	G6F	C15-C14-C4	2.07	114.46	112.31
2	C	501	PLG	C6-N1-C2	2.02	122.86	119.20

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	PLG	C5A-OP4-P-OP1
2	A	501	PLG	C5A-OP4-P-OP2
2	A	501	PLG	C5A-OP4-P-OP3
2	B	501	PLG	C5A-OP4-P-OP1
2	B	501	PLG	C5A-OP4-P-OP2
2	B	501	PLG	C5A-OP4-P-OP3
2	C	501	PLG	C5-C4-C4A-N
2	C	501	PLG	C3-C4-C4A-N
3	B	502	G6F	C42-C46-C47-C48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	502	G6F	C42-C46-C47-C48
2	C	501	PLG	C4-C5-C5A-OP4
3	C	502	G6F	C42-C46-C47-C48


There are no ring outliers.

3 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	PLG	4	0
2	B	501	PLG	1	0
2	C	501	PLG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand G6F B 502




Ligand PLG A 501




Ligand PLG B 501




Ligand G6F A 502



Ligand PLG C 501



Ligand G6F C 502



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	442/442 (100%)	-1.05	0 100 100	15, 28, 50, 77	0
1	B	442/442 (100%)	-1.05	0 100 100	14, 28, 51, 77	0
1	C	442/442 (100%)	-1.06	0 100 100	15, 28, 50, 75	0
All	All	1326/1326 (100%)	-1.05	0 100 100	14, 28, 51, 77	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	G6F	B	502	34/34	0.98	0.05	29,32,56,61	0
3	G6F	C	502	34/34	0.98	0.06	28,31,54,58	0
2	PLG	C	501	20/20	0.99	0.04	19,22,24,24	0
3	G6F	A	502	34/34	0.99	0.05	27,33,59,63	0
2	PLG	A	501	20/20	0.99	0.04	19,22,25,26	0

Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PLG	B	501	20/20	0.99	0.04	18,22,25,26	0
4	CL	A	503	1/1	0.99	0.04	33,33,33,33	0
4	CL	C	503	1/1	0.99	0.04	32,32,32,32	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around G6F B 502:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around G6F C 502:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PLG C 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around G6F A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PLG A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around PLG B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.