



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 1, 2026 – 01:57 PM UTC

PDB ID : 6GVC / pdb_00006gvc
Title : Structure of ArhGAP12 bound to G-Actin
Authors : Mouilleron, S.; Treisman, R.; Diring, J.
Deposited on : 2018-06-20
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	377	
1	B	377	
1	C	377	
1	D	377	
2	Q	231	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	R	231	2% 81% 13% 6%
2	S	231	5% 77% 17% 6%
2	T	231	3% 81% 15% •

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 18354 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Actin, alpha skeletal muscle.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	359	Total	C	N	O	S	0	0	0
			2760	1758	454	529	19			
1	A	356	Total	C	N	O	S	0	2	0
			2755	1748	465	523	19			
1	C	359	Total	C	N	O	S	0	0	0
			2762	1756	460	527	19			
1	D	356	Total	C	N	O	S	0	1	0
			2723	1733	453	518	19			

- Molecule 2 is a protein called Rho GTPase-activating protein 12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	R	216	Total	C	N	O	S	0	0	0
			1704	1096	299	304	5			
2	Q	223	Total	C	N	O	S	0	0	0
			1757	1129	313	310	5			
2	S	217	Total	C	N	O	S	0	0	0
			1700	1093	301	301	5			
2	T	222	Total	C	N	O	S	0	1	0
			1743	1121	309	308	5			

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
R	561	PRO	-	expression tag	UNP Q8C0D4
R	562	GLY	-	expression tag	UNP Q8C0D4
R	563	ILE	-	expression tag	UNP Q8C0D4
R	564	PRO	-	expression tag	UNP Q8C0D4
R	565	GLY	-	expression tag	UNP Q8C0D4
R	566	SER	-	expression tag	UNP Q8C0D4
R	567	THR	-	expression tag	UNP Q8C0D4
Q	561	PRO	-	expression tag	UNP Q8C0D4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
Q	562	GLY	-	expression tag	UNP Q8C0D4
Q	563	ILE	-	expression tag	UNP Q8C0D4
Q	564	PRO	-	expression tag	UNP Q8C0D4
Q	565	GLY	-	expression tag	UNP Q8C0D4
Q	566	SER	-	expression tag	UNP Q8C0D4
Q	567	THR	-	expression tag	UNP Q8C0D4
S	561	PRO	-	expression tag	UNP Q8C0D4
S	562	GLY	-	expression tag	UNP Q8C0D4
S	563	ILE	-	expression tag	UNP Q8C0D4
S	564	PRO	-	expression tag	UNP Q8C0D4
S	565	GLY	-	expression tag	UNP Q8C0D4
S	566	SER	-	expression tag	UNP Q8C0D4
S	567	THR	-	expression tag	UNP Q8C0D4
T	561	PRO	-	expression tag	UNP Q8C0D4
T	562	GLY	-	expression tag	UNP Q8C0D4
T	563	ILE	-	expression tag	UNP Q8C0D4
T	564	PRO	-	expression tag	UNP Q8C0D4
T	565	GLY	-	expression tag	UNP Q8C0D4
T	566	SER	-	expression tag	UNP Q8C0D4
T	567	THR	-	expression tag	UNP Q8C0D4

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		
3	R	1	Total	C	O	0	0
			4	2	2		
3	Q	1	Total	C	O	0	0
			4	2	2		
3	S	1	Total	C	O	0	0
			4	2	2		
3	S	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is LATRUNCULIN B (CCD ID: LAB) (formula: $C_{20}H_{29}NO_5S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	S	0	0
			27	20	1	5	1		
4	A	1	Total	C	N	O	S	0	0
			27	20	1	5	1		
4	C	1	Total	C	N	O	S	0	0
			27	20	1	5	1		
4	D	1	Total	C	N	O	S	0	0
			27	20	1	5	1		

- # ATP

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).




- Molecule 7 is water.



Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	35	Total 35	O 35	0	0
7	C	29	Total 29	O 29	0	0
7	D	17	Total 17	O 17	0	0
7	R	9	Total 9	O 9	0	0
7	Q	16	Total 16	O 16	0	0
7	S	22	Total 22	O 22	0	0
7	T	15	Total 15	O 15	0	0

- Molecule 1: Actin, alpha skeletal muscle









• Molecule 2: Rho GTPase-activating protein 12




• Molecule 2: Rho GTPase-activating protein 12




• Molecule 2: Rho GTPase-activating protein 12



• Molecule 2: Rho GTPase-activating protein 12





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	101.65Å 130.24Å 109.30Å 90.00° 111.17° 90.00°	Depositor
Resolution (Å)	54.13 – 2.60 54.13 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.4 (54.13-2.60) 99.4 (54.13-2.60)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.69 (at 2.61Å)	Xtriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, R_{free}	0.207 , 0.251 0.209 , 0.252	Depositor DCC
R_{free} test set	4064 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	47.8	Xtriage
Anisotropy	0.266	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 51.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	18354	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ATP, LAB, EDO, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.16	0/2817	0.32	0/3823
1	B	0.18	0/2821	0.33	0/3831
1	C	0.18	0/2823	0.35	0/3833
1	D	0.15	0/2785	0.30	0/3786
2	Q	0.18	0/1791	0.38	0/2422
2	R	0.16	0/1736	0.31	0/2351
2	S	0.18	0/1731	0.33	0/2342
2	T	0.18	0/1777	0.35	0/2406
All	All	0.17	0/18281	0.33	0/24794

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2755	0	2685	18	0
1	B	2760	0	2693	26	0
1	C	2762	0	2683	31	1
1	D	2723	0	2632	19	0
2	Q	1757	0	1770	17	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	R	1704	0	1696	15	0
2	S	1700	0	1692	22	0
2	T	1743	0	1747	23	0
3	A	4	0	6	0	0
3	B	4	0	6	0	0
3	C	4	0	6	0	0
3	D	4	0	6	0	0
3	Q	4	0	6	1	0
3	R	4	0	6	0	0
3	S	8	0	12	1	0
4	A	27	0	29	2	0
4	B	27	0	29	2	0
4	C	27	0	29	0	0
4	D	27	0	29	3	0
5	A	31	0	12	0	0
5	B	31	0	12	0	0
5	C	31	0	12	0	0
5	D	31	0	12	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
7	A	35	0	0	0	0
7	B	39	0	0	1	0
7	C	29	0	0	0	0
7	D	17	0	0	1	0
7	Q	16	0	0	1	0
7	R	9	0	0	0	0
7	S	22	0	0	1	0
7	T	15	0	0	0	0
All	All	18354	0	17810	165	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 165 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:208:ILE:HD11	1:C:240:TYR:HE2	1.49	0.77
2:S:730:ARG:HG2	2:S:785:LEU:HD21	1.69	0.75
1:B:61:LYS:NZ	1:A:83:GLU:OE2	2.25	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:R:596:ASP:HB3	2:R:601:SER:HB2	1.74	0.67
2:Q:597:GLN:HE22	3:Q:801:EDO:H12	1.61	0.66

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:364:GLU:O	2:Q:581:ARG:NH1[2_545]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	352/377 (93%)	344 (98%)	8 (2%)	0	100	100
1	B	355/377 (94%)	350 (99%)	5 (1%)	0	100	100
1	C	355/377 (94%)	347 (98%)	8 (2%)	0	100	100
1	D	351/377 (93%)	343 (98%)	8 (2%)	0	100	100
2	Q	219/231 (95%)	210 (96%)	7 (3%)	2 (1%)	14	30
2	R	212/231 (92%)	205 (97%)	6 (3%)	1 (0%)	24	46
2	S	213/231 (92%)	206 (97%)	6 (3%)	1 (0%)	24	46
2	T	219/231 (95%)	210 (96%)	9 (4%)	0	100	100
All	All	2276/2432 (94%)	2215 (97%)	57 (2%)	4 (0%)	43	66

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	Q	666	GLU
2	R	666	GLU
2	Q	654	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	S	666	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	291/320 (91%)	291 (100%)	0	100	100
1	B	293/320 (92%)	293 (100%)	0	100	100
1	C	290/320 (91%)	289 (100%)	1 (0%)	86	94
1	D	284/320 (89%)	284 (100%)	0	100	100
2	Q	188/209 (90%)	185 (98%)	3 (2%)	55	79
2	R	181/209 (87%)	180 (99%)	1 (1%)	78	91
2	S	178/209 (85%)	177 (99%)	1 (1%)	78	91
2	T	186/209 (89%)	185 (100%)	1 (0%)	81	92
All	All	1891/2116 (89%)	1884 (100%)	7 (0%)	84	93

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	Q	688	THR
2	Q	782	LEU
2	T	701	GLN
2	S	783	LEU
2	Q	658	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
2	Q	722	GLN
2	T	669	HIS
2	Q	776	GLN
2	T	716	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	T	624	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 4 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	LAB	D	402	-	28,29,29	1.59	2 (7%)	29,41,41	1.68	7 (24%)
3	EDO	C	401	-	3,3,3	0.45	0	2,2,2	0.35	0
3	EDO	D	401	-	3,3,3	0.44	0	2,2,2	0.37	0
4	LAB	B	402	-	28,29,29	1.58	2 (7%)	29,41,41	1.66	5 (17%)
5	ATP	A	403	6	32,33,33	1.42	4 (12%)	48,52,52	1.71	9 (18%)
4	LAB	A	402	-	28,29,29	1.59	2 (7%)	29,41,41	1.60	7 (24%)
3	EDO	A	401	-	3,3,3	0.42	0	2,2,2	0.38	0
3	EDO	R	801	-	3,3,3	0.47	0	2,2,2	0.27	0
5	ATP	C	403	6	32,33,33	1.48	7 (21%)	48,52,52	1.75	9 (18%)
3	EDO	S	801	-	3,3,3	0.46	0	2,2,2	0.34	0
5	ATP	D	403	6	32,33,33	1.46	6 (18%)	48,52,52	1.72	10 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	B	401	-	3,3,3	0.39	0	2,2,2	0.51	0
5	ATP	B	403	6	32,33,33	1.50	7 (21%)	48,52,52	1.70	8 (16%)
4	LAB	C	402	-	28,29,29	1.57	2 (7%)	29,41,41	1.71	6 (20%)
3	EDO	S	802	-	3,3,3	0.45	0	2,2,2	0.31	0
3	EDO	Q	801	-	3,3,3	0.39	0	2,2,2	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	LAB	D	402	-	-	5/21/49/49	0/2/3/3
3	EDO	C	401	-	-	0/1/1/1	-
3	EDO	D	401	-	-	0/1/1/1	-
4	LAB	B	402	-	-	5/21/49/49	0/2/3/3
5	ATP	A	403	6	-	2/22/38/38	0/3/3/3
4	LAB	A	402	-	-	5/21/49/49	0/2/3/3
3	EDO	A	401	-	-	0/1/1/1	-
3	EDO	R	801	-	-	0/1/1/1	-
5	ATP	C	403	6	-	1/22/38/38	0/3/3/3
3	EDO	S	801	-	-	0/1/1/1	-
5	ATP	D	403	6	-	1/22/38/38	0/3/3/3
3	EDO	B	401	-	-	0/1/1/1	-
5	ATP	B	403	6	-	1/22/38/38	0/3/3/3
4	LAB	C	402	-	-	5/21/49/49	0/2/3/3
3	EDO	S	802	-	-	0/1/1/1	-
3	EDO	Q	801	-	-	0/1/1/1	-

The worst 5 of 32 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	402	LAB	O2-C1	5.51	1.46	1.34
4	C	402	LAB	O2-C1	5.47	1.45	1.34
4	A	402	LAB	O2-C1	5.42	1.45	1.34
4	B	402	LAB	O2-C1	5.33	1.45	1.34
4	B	402	LAB	C18-S1	-5.23	1.66	1.77

The worst 5 of 61 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	403	ATP	C5-C4-N3	-5.50	119.15	126.72
5	A	403	ATP	C5-C4-N3	-5.44	119.23	126.72
5	C	403	ATP	C5-C4-N3	-5.43	119.24	126.72
5	D	403	ATP	C5-C4-N3	-5.33	119.37	126.72
5	D	403	ATP	N3-C4-N9	4.53	134.88	127.17

There are no chirality outliers.

5 of 25 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	402	LAB	C9-C10-C11-C12
4	A	402	LAB	C9-C10-C11-O3
4	C	402	LAB	C9-C10-C11-O3
4	D	402	LAB	C9-C10-C11-O3
4	C	402	LAB	C11-C10-C9-C8


There are no ring outliers.

5 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	402	LAB	3	0
4	B	402	LAB	2	0
4	A	402	LAB	2	0
3	S	801	EDO	1	0
3	Q	801	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand LAB D 402




Bond lengths



Bond angles




Torsions




Rings


Ligand LAB B 402




Bond lengths




Bond angles




Torsions




Rings




Ligand LAB A 402




Bond lengths




Bond angles





Torsions




Rings









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	356/377 (94%)	-0.02	8 (2%) 62 57	21, 45, 81, 113	2 (0%)
1	B	359/377 (95%)	-0.09	5 (1%) 73 69	30, 45, 75, 99	0
1	C	359/377 (95%)	0.05	8 (2%) 62 57	28, 48, 91, 151	0
1	D	356/377 (94%)	0.08	8 (2%) 62 57	23, 49, 90, 127	1 (0%)
2	Q	223/231 (96%)	0.29	7 (3%) 51 45	32, 53, 83, 106	0
2	R	216/231 (93%)	0.24	5 (2%) 61 55	28, 53, 78, 101	0
2	S	217/231 (93%)	0.32	11 (5%) 33 28	37, 55, 77, 90	0
2	T	222/231 (96%)	0.28	7 (3%) 50 44	28, 54, 80, 92	1 (0%)
All	All	2308/2432 (94%)	0.11	59 (2%) 57 51	21, 49, 84, 151	4 (0%)

The worst 5 of 59 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	60	SER	5.5
1	A	60	SER	5.0
2	S	656	GLU	4.7
2	S	655	ASP	4.3
1	C	61	LYS	4.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	S	801	4/4	0.64	0.25	83,84,84,85	0
3	EDO	R	801	4/4	0.84	0.17	50,52,52,52	0
3	EDO	D	401	4/4	0.88	0.15	51,52,52,52	0
3	EDO	S	802	4/4	0.88	0.13	59,60,60,60	0
4	LAB	C	402	27/27	0.88	0.13	56,63,67,68	0
4	LAB	D	402	27/27	0.89	0.12	64,66,67,67	0
4	LAB	B	402	27/27	0.91	0.12	49,54,59,60	0
3	EDO	C	401	4/4	0.91	0.13	54,54,55,55	0
3	EDO	Q	801	4/4	0.91	0.19	77,77,77,77	0
3	EDO	B	401	4/4	0.92	0.08	42,42,43,44	0
4	LAB	A	402	27/27	0.92	0.11	45,49,54,55	0
3	EDO	A	401	4/4	0.94	0.09	46,46,46,47	0
5	ATP	C	403	31/31	0.95	0.07	44,47,51,52	0
5	ATP	D	403	31/31	0.95	0.08	44,47,52,54	0
6	MG	B	404	1/1	0.95	0.17	39,39,39,39	0
5	ATP	B	403	31/31	0.96	0.06	36,43,47,48	0
5	ATP	A	403	31/31	0.97	0.07	39,42,47,47	0
6	MG	D	404	1/1	0.97	0.07	43,43,43,43	0
6	MG	C	404	1/1	0.99	0.04	47,47,47,47	0
6	MG	A	404	1/1	0.99	0.07	45,45,45,45	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around LAB C 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around LAB D 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around LAB B 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around LAB A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ATP C 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ATP D 403:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ATP B 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ATP A 403:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.