



wwPDB NMR Structure Validation Summary Report ⓘ

Mar 9, 2026 – 04:56 PM UTC

PDB ID : 1GX7 / pdb_00001gx7
Title : Best model of the electron transfer complex between cytochrome c3 and [Fe]-hydrogenase
Authors : Elantak, L.; Morelli, X.; Bornet, O.; Hatchikian, C.; Czjzek, M.; Dolla, A.; Guerlesquin, F.
Deposited on : 2002-03-28

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR, THEORETICAL MODEL

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	371	60% 32% 6% .
2	D	88	66% 28% 5% .
3	E	107	58% 28% 14%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA and RNA chains that are outliers for geometric criteria:

Mol	Chain	Compound	Res	Total models with violations	
				Chirality	Geometry
5	A	PDT	4	-	1

2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 5509 atoms, of which 968 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called PERIPLASMIC [FE] HYDROGENASE LARGE SUBUNIT.

Mol	Chain	Residues	Atoms						Trace
1	A	371	Total	C	H	N	O	S	0
			3394	1783	587	465	527	32	


- Molecule 2 is a protein called PERIPLASMIC [FE] HYDROGENASE SMALL SUBUNIT.

Mol	Chain	Residues	Atoms						Trace
2	D	88	Total	C	H	N	O	S	0
			877	457	164	123	132	1	

- Molecule 3 is a protein called CYTOCHROME C3.

Mol	Chain	Residues	Atoms						Trace
3	E	107	Total	C	H	N	O	S	0
			1011	495	201	152	152	11	

- Molecule 4 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms		
4	A	1	Total	Fe	S
			8	4	4

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		
4	A	1	Total	Fe	S
			8	4	4
4	A	1	Total	Fe	S
			8	4	4

- Molecule 5 is 1,3-PROPANEDITHIOL (CCD ID: PDT) (formula: $C_3H_8S_2$).




Mol	Chain	Residues	Atoms		
5	A	1	Total	C	S
			5	3	2

- Molecule 6 is FE (II) ION (CCD ID: FE2) (formula: Fe).


Mol	Chain	Residues	Atoms	
6	A	2	Total	Fe
			2	2

- Molecule 7 is CYANIDE ION (CCD ID: CYN) (formula: CN).



Mol	Chain	Residues	Atoms		
7	A	1	Total	C	N
			2	1	1
7	A	1	Total	C	N
			2	1	1

- Molecule 8 is CARBON MONOXIDE (CCD ID: CMO) (formula: CO).



Mol	Chain	Residues	Atoms		
8	A	1	Total	C	O
			2	1	1
8	A	1	Total	C	O
			2	1	1


-
- The chemical structure of HEC (Hydroxyethylchlorin) is a complex macrocyclic molecule. It features a central iron atom (Fe) coordinated by four nitrogen atoms (N) in a porphyrin-like ring. The central iron atom is also coordinated by a central hydrogen atom (H). The structure includes various side chains and functional groups, such as hydroxyl groups (OH) and carboxylic acid groups (COOH). The atoms are labeled with letters and numbers, indicating their positions within the molecule. The overall structure is highly symmetrical and complex, with many atoms and bonds represented.

Mol	Chain	Residues	Atoms					
9	E	1	Total 47	C 34	Fe 1	H 4	N 4	O 4
9	E	1	Total 47	C 34	Fe 1	H 4	N 4	O 4
9	E	1	Total 47	C 34	Fe 1	H 4	N 4	O 4
9	E	1	Total 47	C 34	Fe 1	H 4	N 4	O 4


4 Residue-property plots

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


• Molecule 1: PERIPLASMIC [FE] HYDROGENASE LARGE SUBUNIT



• Molecule 2: PERIPLASMIC [FE] HYDROGENASE SMALL SUBUNIT



• Molecule 3: CYTOCHROME C3



5 Refinement protocol and experimental data overview ⓘ

The models were refined using the following method: ?.

Of the ? calculated structures, 1 were deposited, based on the following criterion: ?.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	

No chemical shift data was provided.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CYN, CMO, PDT, HEC, SF4, FE2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.68	0/2874 (0.0%)	1.79	75/3885 (1.9%)
2	D	0.67	0/733 (0.0%)	1.74	18/988 (1.8%)
3	E	0.70	0/827 (0.0%)	1.92	35/1101 (3.2%)
All	All	0.68	0/4434 (0.0%)	1.81	128/5974 (2.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	2	20
2	D	1	4
3	E	0	6
All	All	3	30

There are no bond-length outliers.

5 of 128 angle outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	378	CYS	N-CA-C	-9.00	100.12	108.22
1	A	331	ASP	N-CA-C	-8.98	95.68	109.24
1	A	100	VAL	N-CA-C	-8.86	95.71	108.11
3	E	43	TYR	N-CA-C	-8.82	94.86	109.07
3	E	58	LYS	N-CA-C	-8.39	96.23	109.23

All chiral outliers are listed below.

Mol	Chain	Res	Type	Atoms
1	A	50	ILE	CB

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	A	63	ILE	CB
2	D	39	ILE	CB

5 of 29 planar outliers are listed below.

Mol	Chain	Res	Type	Group
1	A	36	ILE	Peptide
1	A	37	GLY	Peptide
1	A	68	ASN	Peptide
1	A	71	GLN	Peptide
1	A	82	TYR	Sidechain

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	2807	587	2765	54
2	D	713	164	703	10
3	E	810	201	792	9
4	A	24	0	0	6
5	A	5	0	6	12
6	A	2	0	0	2
7	A	4	0	0	7
8	A	4	0	0	7
9	E	172	16	120	5
All	All	4541	968	4386	70

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

5 of 70 clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:238:LYS:HG2	1:A:257:THR:HB	0.90	1.44
1:A:232:MET:SD	8:A:9:CMO:C	0.81	2.68
1:A:296:PHE:HB3	5:A:4:PDT:H11	0.80	1.51
1:A:232:MET:SD	7:A:7:CYN:C	0.78	2.70
1:A:104:ALA:HB2	1:A:139:PHE:HB3	0.75	1.58

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	369/371 (99%)	311 (84%)	49 (13%)	9 (2%)	7	44
2	D	86/88 (98%)	75 (87%)	10 (12%)	1 (1%)	13	61
3	E	105/107 (98%)	76 (72%)	20 (19%)	9 (9%)	1	12
All	All	560/566 (99%)	462 (82%)	79 (14%)	19 (3%)	4	34

5 of 19 Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	33	ALA
1	A	57	PRO
1	A	61	PRO
1	A	67	ILE
1	A	198	SER

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	297/297 (100%)	265 (89%)	32 (11%)	8	52
2	D	76/76 (100%)	69 (91%)	7 (9%)	11	56
3	E	88/88 (100%)	78 (89%)	10 (11%)	8	50
All	All	461/461 (100%)	412 (89%)	49 (11%)	9	52

5 of 49 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	29	GLN
1	A	40	THR
1	A	42	SER
1	A	47	THR
1	A	58	HIS

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.6 Ligand geometry ⓘ

Of 14 ligands modelled in this entry, 2 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
4	SF4	A	2	1	0,12,12	0.00	-
5	PDT	A	4	6	4,4,4	2.29	2 (50%)
8	CMO	A	10	7,6	0,1,1	0.00	-
9	HEC	E	111	3	46,50,50	1.48	2 (4%)
8	CMO	A	9	7,6	0,1,1	0.00	-
4	SF4	A	1	1	0,12,12	0.00	-
9	HEC	E	110	3	46,50,50	1.45	2 (4%)
4	SF4	A	3	1	0,12,12	0.00	-
9	HEC	E	109	3	46,50,50	1.44	2 (4%)

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
7	CYN	A	8	6,8	1,1,1	0.64	0 (0%)
7	CYN	A	7	6,8	1,1,1	0.49	0 (0%)
9	HEC	E	112	3	46,50,50	1.48	2 (4%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
4	SF4	A	2	1	-	-	-
5	PDT	A	4	6	3,3,3	1.43	0 (0%)
8	CMO	A	10	7,6	-	-	-
9	HEC	E	111	3	58,82,82	1.65	4 (6%)
8	CMO	A	9	7,6	-	-	-
4	SF4	A	1	1	-	-	-
9	HEC	E	110	3	58,82,82	1.54	2 (3%)
4	SF4	A	3	1	-	-	-
9	HEC	E	109	3	58,82,82	1.54	4 (6%)
7	CYN	A	8	6,8	-	-	-
7	CYN	A	7	6,8	-	-	-
9	HEC	E	112	3	58,82,82	1.66	3 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SF4	A	2	1	-	-	0,6,5,5
5	PDT	A	4	6	-	0,2,2,2	-
9	HEC	E	111	3	-	0,14,54,54	-
9	HEC	E	110	3	-	0,14,54,54	-
4	SF4	A	1	1	-	-	0,6,5,5
9	HEC	E	109	3	-	0,14,54,54	-
4	SF4	A	3	1	-	-	0,6,5,5
9	HEC	E	112	3	-	0,14,54,54	-

5 of 10 bond outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	E	111	HEC	CAC-C3C	5.99	1.54	1.35
9	E	110	HEC	CAC-C3C	5.85	1.54	1.35
9	E	112	HEC	CAB-C3B	5.85	1.54	1.35
9	E	112	HEC	CAC-C3C	5.84	1.54	1.35
9	E	111	HEC	CAB-C3B	5.84	1.54	1.35

5 of 13 angle outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	E	112	HEC	CBB-CAB-C3B	7.90	111.64	127.43
9	E	110	HEC	CBC-CAC-C3C	7.56	112.31	127.43
9	E	111	HEC	CBB-CAB-C3B	7.29	112.87	127.43
9	E	109	HEC	CBB-CAB-C3B	7.21	113.03	127.43
9	E	112	HEC	CBC-CAC-C3C	7.04	113.36	127.43


There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand HEC E 111




Bond lengths



Bond angles




Torsions




Rings


Ligand HEC E 110




Bond lengths



Bond angles




Torsions




Rings


Ligand HEC E 109




Bond lengths




Bond angles



Torsions



Rings



6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided