



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:41 PM UTC

PDB ID : 5GX7 / pdb_00005gx7
Title : Crystal structure of solute-binding protein complexed with unsaturated chondroitin disaccharide with a sulfate group at C-6 position of GalNAc
Authors : Oiki, S.; Mikami, B.; Murata, K.; Hashimoto, W.
Deposited on : 2016-09-15
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	483	2% 92% 5% ..
1	B	483	3% 89% 8% ..
1	C	483	3% 90% 8% .
2	D	2	100%
2	E	2	50% 50%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	2	 100%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 12349 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Extracellular solute-binding protein family 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	474	Total	C	N	O	S	0	0	0
			3833	2466	633	728	6			
1	B	474	Total	C	N	O	S	0	0	0
			3833	2466	633	728	6			
1	C	474	Total	C	N	O	S	0	0	0
			3833	2466	633	728	6			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	18	MET	-	expression tag	UNP D1AWE0
B	18	MET	-	expression tag	UNP D1AWE0
C	18	MET	-	expression tag	UNP D1AWE0

- Molecule 2 is an oligosaccharide called 4-deoxy-alpha-L-threo-hex-4-enopyranuronic acid-(1-3)-2-acetamido-2-deoxy-6-O-sulfo-beta-D-galactopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	2	Total	C	N	O	S	0	0	0
			30	14	1	14	1			
2	E	2	Total	C	N	O	S	0	0	0
			30	14	1	14	1			
2	F	2	Total	C	N	O	S	0	0	0
			30	14	1	14	1			

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Ca 1	0	0
3	B	1	Total 1	Ca 1	0	0
3	C	1	Total 1	Ca 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	294	Total 294	O 294	0	0
4	B	260	Total 260	O 260	0	0
4	C	203	Total 203	O 203	0	0

- Molecule 1: Extracellular solute-binding protein family 1



M61
GCD2

- Molecule 2: 4-deoxy-alpha-L-threo-hex-4-enopyranuronic acid-(1-3)-2-acetamido-2-deoxy-6-O-sulfo-beta-D-galactopyranose

Chain E:  50% 50%M61
GCD2

- Molecule 2: 4-deoxy-alpha-L-threo-hex-4-enopyranuronic acid-(1-3)-2-acetamido-2-deoxy-6-O-sulfo-beta-D-galactopyranose

Chain F:  100%M61
GCD2

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	80.28Å 112.44Å 166.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.67 – 1.99 45.67 – 1.99	Depositor EDS
% Data completeness (in resolution range)	97.2 (45.67-1.99) 97.3 (45.67-1.99)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.21 (at 2.00Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.197 , 0.249 (Not available) , 0.238	Depositor DCC
R_{free} test set	5094 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	22.0	Xtriage
Anisotropy	0.479	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 27.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	12349	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NG6, CA, GCD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.91	1/3923 (0.0%)	0.87	2/5308 (0.0%)
1	B	0.87	1/3923 (0.0%)	0.85	2/5308 (0.0%)
1	C	0.86	1/3923 (0.0%)	0.88	0/5308
All	All	0.88	3/11769 (0.0%)	0.87	4/15924 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	127	ARG	CZ-NH2	-6.81	1.24	1.33
1	C	127	ARG	CZ-NH2	-6.38	1.25	1.33
1	A	68	THR	CB-CG2	-6.29	1.31	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	442	LYS	CB-CG-CD	-7.08	95.02	111.30
1	A	409	TYR	OH-CZ-CE2	-5.34	103.86	119.90
1	B	442	LYS	CG-CD-CE	5.19	123.24	111.30
1	A	409	TYR	CE1-CZ-OH	5.13	135.30	119.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3833	0	3788	22	0
1	B	3833	0	3788	30	0
1	C	3833	0	3788	25	0
2	D	30	0	13	0	0
2	E	30	0	14	3	0
2	F	30	0	13	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	A	294	0	0	1	0
4	B	260	0	0	4	0
4	C	203	0	0	3	0
All	All	12349	0	11404	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:387:THR:HG22	1:B:390:ALA:H	1.18	1.05
1:C:253:LYS:HE3	4:C:883:HOH:O	1.79	0.81
1:B:442:LYS:HE2	1:B:443:GLU:OE1	1.85	0.75
1:C:382:ASN:OD1	1:C:386:LYS:HG3	1.90	0.72
1:B:127:ARG:NH2	1:B:131:ASP:OD1	2.27	0.67
1:B:293:ARG:HH11	1:B:293:ARG:HG3	1.58	0.66
1:B:387:THR:CG2	1:B:390:ALA:H	2.04	0.66
1:B:141:MET:HE3	1:B:328:GLY:HA3	1.78	0.65
1:C:141:MET:HE3	1:C:328:GLY:HA3	1.76	0.65
1:C:445:SER:O	1:C:449:VAL:HG22	1.98	0.64
1:B:387:THR:HG22	1:B:390:ALA:N	2.02	0.63
1:A:113:GLN:HG2	1:A:114:HIS:CD2	2.33	0.63
1:B:135:VAL:HG11	1:B:440:TYR:O	2.00	0.61
1:C:438:LEU:HA	4:C:852:HOH:O	2.00	0.61
1:C:111:ILE:HA	1:C:115:ALA:HB3	1.83	0.60
1:A:68:THR:HG22	4:A:920:HOH:O	2.00	0.59
1:C:135:VAL:HG11	1:C:440:TYR:O	2.03	0.59
1:C:36:HIS:CE1	1:C:400:ARG:HB2	2.37	0.59
1:A:387:THR:HG22	1:A:389:LEU:N	2.18	0.58
1:A:372:LYS:HE3	1:A:373:PRO:HD2	1.87	0.57
1:B:293:ARG:HG3	1:B:293:ARG:NH1	2.17	0.56
1:C:229:ASN:C	1:C:231:MET:H	2.13	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:36:HIS:CE1	2:E:1:NG6:C8	2.89	0.56
1:B:293:ARG:HG2	4:B:770:HOH:O	2.06	0.55
1:A:294:ASN:HD21	1:A:385:GLY:CA	2.21	0.54
1:B:123:GLU:O	1:C:126:PRO:HG2	2.08	0.53
1:A:149:PHE:HA	1:A:152:LYS:HE3	1.91	0.53
1:B:377:ASP:OD1	1:B:381:LYS:HE2	2.09	0.53
1:A:111:ILE:HA	1:A:115:ALA:HB3	1.92	0.52
1:B:387:THR:HG23	4:B:903:HOH:O	2.09	0.52
1:B:36:HIS:HE1	2:E:1:NG6:C8	2.24	0.51
1:C:50:LYS:HE2	1:C:54:ASP:OD1	2.09	0.51
1:A:231:MET:HE3	1:A:497:ASN:OD1	2.12	0.50
1:C:41:LEU:HD11	1:C:48:PHE:CE2	2.47	0.49
1:B:207:ASN:ND2	1:B:210:LYS:HE2	2.28	0.48
1:A:387:THR:HG22	1:A:390:ALA:H	1.77	0.48
1:A:387:THR:HG22	1:A:389:LEU:H	1.77	0.48
1:C:28:LYS:HD2	4:C:723:HOH:O	2.12	0.48
1:A:143:PRO:HB3	1:A:325:TYR:CD2	2.49	0.48
1:A:294:ASN:HD21	1:A:385:GLY:HA2	1.79	0.48
1:A:148:TYR:O	1:A:152:LYS:HG3	2.13	0.48
1:B:206:ASN:OD1	1:B:210:LYS:HE3	2.14	0.48
1:B:108:GLU:HG2	1:B:109:ASP:N	2.28	0.47
1:C:229:ASN:O	1:C:231:MET:N	2.48	0.47
1:B:99:GLY:HA2	1:B:104:LEU:HB2	1.96	0.46
1:B:423:LYS:NZ	1:B:427:ASP:OD2	2.44	0.46
1:C:289:SER:HB2	1:C:418:ALA:HB2	1.98	0.46
1:B:393:ARG:HH22	2:E:1:NG6:C8	2.28	0.46
1:A:113:GLN:HG2	1:A:114:HIS:NE2	2.31	0.46
1:A:100:ILE:CG2	1:A:442:LYS:HG3	2.46	0.46
1:C:223:PRO:HA	1:C:236:PRO:HG3	1.97	0.45
1:C:323:THR:HG22	1:C:434:GLU:HG2	1.98	0.45
1:B:141:MET:CE	1:B:328:GLY:HA3	2.44	0.45
1:B:472:ASP:HB3	1:B:475:LYS:HG3	1.99	0.44
1:A:239:LYS:HE3	4:B:919:HOH:O	2.16	0.44
1:B:36:HIS:HD2	4:B:830:HOH:O	1.99	0.44
1:C:387:THR:HG23	1:C:390:ALA:CB	2.47	0.44
1:A:148:TYR:CZ	1:A:152:LYS:HG2	2.52	0.44
1:C:50:LYS:HE2	1:C:54:ASP:CG	2.42	0.44
1:A:113:GLN:O	1:A:113:GLN:HG3	2.17	0.43
1:A:227:GLU:HG2	1:A:228:GLU:N	2.32	0.43
1:C:442:LYS:H	1:C:442:LYS:HG3	1.58	0.43
1:C:166:LYS:HB2	1:C:166:LYS:HE3	1.87	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:131:ASP:HB3	1:A:325:TYR:CE1	2.55	0.41
1:C:120:LYS:HE2	1:C:124:GLU:OE2	2.19	0.41
1:A:148:TYR:O	1:A:152:LYS:HE3	2.21	0.41
1:B:55:THR:O	1:B:56:ASN:HB2	2.21	0.41
1:C:112:ASN:O	1:C:119:LYS:NZ	2.54	0.41
1:B:490:SER:HA	1:B:493:GLU:OE2	2.21	0.41
1:C:141:MET:CE	1:C:328:GLY:HA3	2.47	0.41
1:A:442:LYS:HE3	1:A:443:GLU:OE2	2.21	0.40
1:B:205:ALA:HB1	1:B:210:LYS:HB2	2.03	0.40
1:C:95:LEU:HA	1:C:95:LEU:HD12	1.82	0.40
1:B:111:ILE:HA	1:B:115:ALA:HB3	2.03	0.40
1:B:422:TYR:O	1:B:426:MET:HG2	2.21	0.40
1:B:337:ASP:CG	1:B:340:SER:HB3	2.47	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	472/483 (98%)	463 (98%)	8 (2%)	1 (0%)	43	42
1	B	472/483 (98%)	459 (97%)	12 (2%)	1 (0%)	43	42
1	C	472/483 (98%)	459 (97%)	11 (2%)	2 (0%)	30	27
All	All	1416/1449 (98%)	1381 (98%)	31 (2%)	4 (0%)	36	35

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	442	LYS
1	A	80	GLU
1	C	437	ILE
1	B	437	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	408/416 (98%)	406 (100%)	2 (0%)	81	87
1	B	408/416 (98%)	404 (99%)	4 (1%)	68	75
1	C	408/416 (98%)	407 (100%)	1 (0%)	87	92
All	All	1224/1248 (98%)	1217 (99%)	7 (1%)	78	85

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	387	THR
1	A	409	TYR
1	B	152	LYS
1	B	387	THR
1	B	409	TYR
1	B	441	THR
1	C	409	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	65	GLN
1	A	229	ASN
1	A	294	ASN
1	A	398	GLN
1	A	405	GLN
1	B	36	HIS
1	B	65	GLN
1	B	405	GLN
1	C	65	GLN
1	C	97	ASN
1	C	150	ASN
1	C	155	GLN
1	C	398	GLN
1	C	405	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NG6	D	1	2	19,19,19	1.47	2 (10%)	27,28,28	2.03	8 (29%)
2	GCD	D	2	2	10,11,12	2.59	4 (40%)	12,15,17	1.91	1 (8%)
2	NG6	E	1	2	19,19,19	1.21	3 (15%)	27,28,28	2.22	5 (18%)
2	GCD	E	2	2	10,11,12	1.48	2 (20%)	12,15,17	0.98	1 (8%)
2	NG6	F	1	2	19,19,19	1.52	3 (15%)	27,28,28	1.62	5 (18%)
2	GCD	F	2	2	10,11,12	1.83	2 (20%)	12,15,17	1.52	3 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NG6	D	1	2	-	0/10/30/30	0/1/1/1
2	GCD	D	2	2	-	0/4/17/20	0/1/1/1
2	NG6	E	1	2	-	5/10/30/30	0/1/1/1
2	GCD	E	2	2	-	0/4/17/20	0/1/1/1
2	NG6	F	1	2	-	3/10/30/30	0/1/1/1
2	GCD	F	2	2	-	0/4/17/20	0/1/1/1

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2	GCD	O5-C5	5.43	1.44	1.37
2	D	1	NG6	O2S-S	4.70	1.65	1.45
2	F	1	NG6	O2S-S	4.15	1.63	1.45
2	D	2	GCD	C3-C4	3.91	1.55	1.50
2	F	2	GCD	O5-C5	3.38	1.41	1.37
2	F	1	NG6	C8-C7	-3.20	1.43	1.50
2	D	1	NG6	O6-S	-3.01	1.48	1.56
2	F	2	GCD	C3-C4	2.84	1.54	1.50
2	F	1	NG6	O7-C7	2.63	1.29	1.23
2	D	2	GCD	O5-C1	-2.59	1.41	1.45
2	E	1	NG6	O3S-S	2.59	1.66	1.50
2	D	2	GCD	O6A-C6	2.49	1.28	1.22
2	E	2	GCD	O5-C5	2.41	1.40	1.37
2	E	2	GCD	C4-C5	2.41	1.37	1.33
2	E	1	NG6	C8-C7	-2.28	1.45	1.50
2	E	1	NG6	O1-C1	2.09	1.46	1.39

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	1	NG6	O6-C6-C5	-8.39	92.61	107.57
2	D	1	NG6	O6-C6-C5	-5.56	97.66	107.57
2	D	1	NG6	O6-S-O2S	-5.41	90.40	106.92
2	D	2	GCD	O5-C5-C4	-5.11	120.25	124.94
2	F	1	NG6	C6-C5-C4	-3.52	104.68	112.07
2	E	1	NG6	O3S-S-O6	3.35	114.08	106.37
2	E	1	NG6	O6-S-O2S	-3.35	96.70	106.92
2	F	1	NG6	O6-C6-C5	3.25	113.37	107.57
2	D	1	NG6	O3S-S-O2S	3.13	119.51	108.56
2	F	2	GCD	O5-C5-C4	-2.88	122.29	124.94
2	F	2	GCD	C3-C4-C5	2.53	125.73	121.45
2	F	1	NG6	O3S-S-O6	2.47	112.06	106.37
2	E	1	NG6	O7-C7-N2	2.47	126.35	121.98
2	F	1	NG6	O1-C1-O5	-2.47	103.08	110.41
2	E	1	NG6	O2S-S-O1S	2.44	121.64	112.24
2	D	1	NG6	O1-C1-C2	-2.34	104.35	109.22
2	D	1	NG6	O6-S-O1S	2.34	114.06	106.92
2	F	1	NG6	C1-O5-C5	-2.17	109.46	113.65
2	F	2	GCD	O5-C5-C6	2.14	115.79	111.85
2	D	1	NG6	C1-O5-C5	-2.09	109.60	113.65
2	E	2	GCD	O6B-C6-C5	2.07	118.76	114.06
2	D	1	NG6	O5-C1-C2	-2.06	107.44	109.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	NG6	O3-C3-C2	-2.03	105.54	109.58

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	1	NG6	C8-C7-N2-C2
2	E	1	NG6	O7-C7-N2-C2
2	F	1	NG6	C8-C7-N2-C2
2	F	1	NG6	O7-C7-N2-C2
2	E	1	NG6	C6-O6-S-O2S
2	E	1	NG6	C6-O6-S-O3S
2	F	1	NG6	C5-C6-O6-S
2	E	1	NG6	C6-O6-S-O1S


There are no ring outliers.

1 monomer is involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	1	NG6	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.


Oligosaccharide Chain D




Bond lengths




Bond angles




Torsions



Rings





5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	474/483 (98%)	-0.13	9 (1%) 66 66	13, 20, 34, 56	0
1	B	474/483 (98%)	-0.01	16 (3%) 48 47	15, 22, 41, 60	3 (0%)
1	C	474/483 (98%)	0.15	16 (3%) 48 47	14, 23, 46, 62	2 (0%)
All	All	1422/1449 (98%)	0.01	41 (2%) 53 52	13, 21, 41, 62	5 (0%)

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	27	PRO	8.7
1	C	437	ILE	5.5
1	B	437	ILE	4.9
1	A	437	ILE	4.8
1	B	436	PRO	4.4
1	C	500	PHE	4.0
1	A	81	GLY	3.7
1	A	385	GLY	3.6
1	B	27	PRO	3.6
1	C	438	LEU	3.5
1	B	499	ALA	3.3
1	C	436	PRO	3.3
1	B	229	ASN	3.1
1	C	383	PRO	3.0
1	C	231	MET	3.0
1	C	499	ALA	3.0
1	B	293	ARG	2.9
1	C	27	PRO	2.9
1	B	438	LEU	2.9
1	B	384	ASP	2.8
1	B	440	TYR	2.7
1	A	387	THR	2.6
1	B	383	PRO	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	500	PHE	2.5
1	B	385	GLY	2.4
1	B	442	LYS	2.4
1	C	442	LYS	2.4
1	C	135	VAL	2.4
1	C	441	THR	2.4
1	B	500	PHE	2.4
1	B	441	THR	2.4
1	B	231	MET	2.3
1	A	436	PRO	2.2
1	C	379	VAL	2.2
1	C	370	ASP	2.2
1	A	384	ASP	2.1
1	B	443	GLU	2.1
1	A	112	ASN	2.1
1	C	364	SER	2.0
1	C	367	THR	2.0
1	C	449	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NG6	E	1	19/19	0.90	0.09	9,17,38,48	0
2	NG6	F	1	19/19	0.91	0.08	13,21,35,39	0
2	NG6	D	1	19/19	0.93	0.07	13,16,38,40	0
2	GCD	F	2	11/12	0.95	0.06	16,18,20,20	0
2	GCD	D	2	11/12	0.96	0.05	15,16,17,18	0
2	GCD	E	2	11/12	0.96	0.05	15,17,18,19	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain D:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CA	A	603	1/1	0.97	0.04	22,22,22,22	0
3	CA	C	603	1/1	0.97	0.04	25,25,25,25	0
3	CA	B	603	1/1	1.00	0.02	18,18,18,18	0

6.5 Other polymers [i](#)

There are no such residues in this entry.