



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:51 PM UTC

PDB ID : 3GYN / pdb_00003gyn
Title : Crystal structure of HCV NS5B polymerase with a novel monocyclic dihydropyridinone inhibitor
Authors : Zhao, Q.; Showalter, R.E.; Han, Q.; Kissinger, C.R.
Deposited on : 2009-04-04
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	578	7% 89% 7% .
1	B	578	8% 89% 7% .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8944 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called RNA-directed RNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	557	Total	C	N	O	S	0	0	0
			4334	2735	764	804	31			
1	B	557	Total	C	N	O	S	0	0	0
			4334	2735	764	804	31			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	544	GLN	ARG	engineered mutation	UNP P26663
A	571	LEU	-	expression tag	UNP P26663
A	572	GLU	-	expression tag	UNP P26663
A	573	HIS	-	expression tag	UNP P26663
A	574	HIS	-	expression tag	UNP P26663
A	575	HIS	-	expression tag	UNP P26663
A	576	HIS	-	expression tag	UNP P26663
A	577	HIS	-	expression tag	UNP P26663
A	578	HIS	-	expression tag	UNP P26663
B	544	GLN	ARG	engineered mutation	UNP P26663
B	571	LEU	-	expression tag	UNP P26663
B	572	GLU	-	expression tag	UNP P26663
B	573	HIS	-	expression tag	UNP P26663
B	574	HIS	-	expression tag	UNP P26663
B	575	HIS	-	expression tag	UNP P26663
B	576	HIS	-	expression tag	UNP P26663
B	577	HIS	-	expression tag	UNP P26663
B	578	HIS	-	expression tag	UNP P26663

- Molecule 2 is N-{3-[(5R)-1-cyclopentyl-4-hydroxy-5-methyl-5-(3-methylbutyl)-2-oxo-1,2,5,6-tetrahydropyridin-3-yl]-1,1-dioxido-4H-1,2,4-benzothiadiazin-7-yl}methanesulfonamide (CCD ID: B42) (formula: C₂₄H₃₄N₄O₆S₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			36	24	4	6	2		
2	B	1	Total	C	N	O	S	0	0
			36	24	4	6	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	98	Total	O	0	0
			98	98		
3	B	106	Total	O	0	0
			106	106		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RNA-directed RNA polymerase



• Molecule 1: RNA-directed RNA polymerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	85.86Å 106.39Å 126.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.15 – 2.15 34.15 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.7 (34.15-2.15) 99.9 (34.15-2.15)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.37 (at 2.14Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.225 , 0.276 0.229 , 0.275	Depositor DCC
R_{free} test set	3231 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	30.7	Xtriage
Anisotropy	0.328	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 30.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8944	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 53.65 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.0897e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: B42

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/4428	0.78	0/6009
1	B	0.50	0/4428	0.80	0/6009
All	All	0.50	0/8856	0.79	0/12018

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4334	0	4354	25	0
1	B	4334	0	4354	22	0
2	A	36	0	34	1	0
2	B	36	0	34	2	0
3	A	98	0	0	1	0
3	B	106	0	0	0	0
All	All	8944	0	8776	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:451:CYS:HB2	3:A:677:HOH:O	1.81	0.79
1:B:12:THR:HG21	1:B:270:LYS:HE2	1.69	0.75
1:A:51:LYS:HE3	1:A:156:PRO:HG2	1.72	0.72
1:B:308:LEU:HB2	1:B:311:CYS:SG	2.33	0.68
1:A:308:LEU:HB3	1:A:311:CYS:SG	2.38	0.63
1:A:22:PRO:HG2	1:A:401:ARG:HG3	1.86	0.57
1:A:313:MET:HG2	1:A:322:VAL:HG22	1.86	0.57
1:A:110:ASN:HD22	1:A:110:ASN:C	2.13	0.55
1:A:381:VAL:HG11	1:A:474:LEU:CD2	2.37	0.55
1:A:381:VAL:HG11	1:A:474:LEU:HD21	1.89	0.55
1:A:83:LEU:HB2	1:A:173:MET:HA	1.90	0.54
1:B:219:TYR:HB3	1:B:320:LEU:HD23	1.90	0.54
1:B:160:ILE:HA	1:B:282:SER:OG	2.08	0.52
1:A:82:LEU:HD13	1:A:249:ALA:HB2	1.92	0.51
1:A:144:VAL:HB	1:A:394:ARG:HG2	1.91	0.51
1:A:172:LYS:HE3	1:A:560:ILE:HD13	1.92	0.51
1:B:308:LEU:CB	1:B:311:CYS:SG	2.98	0.51
1:B:22:PRO:HG2	1:B:401:ARG:HG3	1.92	0.50
1:B:187:MET:HE2	1:B:293:LEU:HD23	1.94	0.49
1:A:409:LEU:HD21	1:A:454:ILE:HB	1.95	0.49
1:A:410:GLY:C	2:A:579:B42:H11	2.39	0.48
1:A:160:ILE:HA	1:A:282:SER:OG	2.14	0.48
1:A:233:ILE:O	1:A:236:GLU:HG2	2.14	0.47
1:B:346:TYR:O	1:B:347:SER:HB3	2.15	0.47
1:B:172:LYS:HE3	1:B:560:ILE:HD13	1.97	0.46
1:B:227:THR:HB	1:B:347:SER:O	2.14	0.46
1:A:219:TYR:HB3	1:A:320:LEU:HD23	1.97	0.46
1:A:77:THR:O	1:B:24:ASN:HB2	2.17	0.45
1:B:514:GLN:HB2	1:B:518:ALA:HB3	1.99	0.44
1:A:429:PHE:O	1:A:433:LEU:HG	2.16	0.44
1:B:24:ASN:ND2	1:B:400:ALA:HA	2.32	0.44
1:B:76:SER:HA	1:B:242:CYS:O	2.17	0.43
1:B:83:LEU:HB2	1:B:173:MET:HA	1.99	0.43
1:B:144:VAL:HB	1:B:394:ARG:HG2	2.00	0.43
1:A:277:ARG:HG2	1:A:279:CYS:H	1.84	0.43
1:A:268:ASN:HB3	1:A:274:CYS:SG	2.58	0.43
1:A:308:LEU:CB	1:A:311:CYS:SG	3.07	0.43
1:B:508:ARG:HH22	1:B:534:LEU:HD13	1.83	0.42
1:B:414:MET:HG3	2:B:579:B42:H13A	2.01	0.42
1:B:429:PHE:O	1:B:433:LEU:HG	2.19	0.42
1:B:521:CYS:O	1:B:525:LEU:HB2	2.20	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:197:PRO:O	1:A:201:VAL:HG23	2.20	0.41
1:B:466:LEU:HD22	1:B:551:PHE:HE2	1.85	0.41
1:A:539:ILE:HA	1:A:540:PRO:HD3	1.94	0.40
2:B:579:B42:N25	2:B:579:B42:O34	2.53	0.40
1:B:527:ASN:O	1:B:533:LYS:HD3	2.22	0.40
1:A:346:TYR:O	1:A:347:SER:HB3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	553/578 (96%)	539 (98%)	14 (2%)	0	100	100
1	B	553/578 (96%)	544 (98%)	9 (2%)	0	100	100
All	All	1106/1156 (96%)	1083 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	475/493 (96%)	471 (99%)	4 (1%)	73	79
1	B	475/493 (96%)	472 (99%)	3 (1%)	78	84

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	950/986 (96%)	943 (99%)	7 (1%)	76	82

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	110	ASN
1	A	160	ILE
1	A	440	GLU
1	A	461	GLN
1	B	160	ILE
1	B	461	GLN
1	B	544	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	110	ASN
1	A	117	ASN
1	A	330	GLN
1	B	117	ASN
1	B	142	ASN
1	B	330	GLN
1	B	411	ASN
1	B	461	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	B42	A	579	-	39,39,39	2.45	6 (15%)	44,61,61	1.83	8 (18%)
2	B42	B	579	-	39,39,39	2.35	6 (15%)	44,61,61	1.81	9 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	B42	A	579	-	-	1/15/64/64	0/4/4/4
2	B42	B	579	-	-	3/15/64/64	0/4/4/4

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	579	B42	C9-N26	9.74	1.42	1.30
2	B	579	B42	C9-N26	9.13	1.41	1.30
2	A	579	B42	S36-N28	7.32	1.72	1.63
2	B	579	B42	S36-N28	7.10	1.72	1.63
2	B	579	B42	C9-N25	-4.89	1.30	1.36
2	A	579	B42	C9-N25	-4.68	1.30	1.36
2	A	579	B42	S35-N26	4.33	1.69	1.60
2	B	579	B42	S35-N26	4.04	1.69	1.60
2	A	579	B42	C7-C10	3.23	1.51	1.44
2	B	579	B42	C7-C10	2.92	1.51	1.44
2	A	579	B42	C10-N27	2.78	1.40	1.35
2	B	579	B42	C10-N27	2.42	1.39	1.35

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	579	B42	C21-S36-N28	-5.93	99.89	106.56
2	B	579	B42	O31-S35-O30	5.40	122.44	115.90
2	B	579	B42	C21-S36-N28	-5.27	100.63	106.56
2	A	579	B42	O31-S35-C6	-4.76	104.56	108.82
2	A	579	B42	C6-C4-N25	4.44	124.09	121.00
2	A	579	B42	O31-S35-O30	4.40	121.23	115.90
2	B	579	B42	C6-C4-N25	4.05	123.81	121.00
2	B	579	B42	O31-S35-C6	-3.95	105.27	108.82
2	B	579	B42	C18-C17-C15	-3.14	106.04	109.44
2	A	579	B42	C6-S35-N26	-3.05	100.68	105.10
2	B	579	B42	C6-S35-N26	-3.05	100.69	105.10
2	A	579	B42	O29-C10-C7	-2.41	120.01	125.16
2	A	579	B42	O33-S36-N28	2.40	111.90	107.10
2	B	579	B42	O32-S36-C21	-2.39	104.48	108.26
2	B	579	B42	O29-C10-C7	-2.37	120.09	125.16
2	A	579	B42	O32-S36-C21	-2.27	104.66	108.26
2	B	579	B42	O33-S36-N28	2.17	111.43	107.10

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	579	B42	C5-N28-S36-O33
2	B	579	B42	C5-N28-S36-O32
2	A	579	B42	C5-N28-S36-O33
2	B	579	B42	C5-N28-S36-C21


There are no ring outliers.


2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	579	B42	1	0
2	B	579	B42	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	557/578 (96%)	0.67	40 (7%) 21 25	31, 41, 59, 94	0
1	B	557/578 (96%)	0.75	48 (8%) 16 17	31, 43, 63, 91	0
All	All	1114/1156 (96%)	0.71	88 (7%) 18 20	31, 42, 61, 94	0

All (88) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	534	LEU	6.9
1	A	549	GLY	6.1
1	A	545	LEU	5.8
1	B	534	LEU	5.4
1	B	541	ALA	5.1
1	A	537	THR	5.1
1	B	545	LEU	5.0
1	B	532	THR	4.9
1	A	541	ALA	4.8
1	A	532	THR	4.6
1	A	536	LEU	4.6
1	A	533	LYS	4.6
1	A	535	LYS	4.6
1	B	535	LYS	4.4
1	A	547	LEU	4.4
1	A	542	ALA	4.4
1	B	530	VAL	4.4
1	A	540	PRO	4.0
1	A	539	ILE	4.0
1	B	539	ILE	4.0
1	B	547	LEU	3.9
1	B	497	LEU	3.6
1	B	14	CYS	3.6
1	A	513	SER	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	543	SER	3.5
1	B	542	ALA	3.5
1	B	540	PRO	3.4
1	A	14	CYS	3.2
1	A	543	SER	3.2
1	B	549	GLY	3.2
1	B	550	TRP	3.1
1	A	550	TRP	3.1
1	B	376	ALA	3.1
1	B	95	HIS	3.1
1	B	533	LYS	3.0
1	B	381	VAL	3.0
1	A	378	GLY	2.9
1	A	538	PRO	2.9
1	B	538	PRO	2.9
1	B	544	GLN	2.9
1	A	311	CYS	2.9
1	A	531	LYS	2.8
1	A	110	ASN	2.8
1	B	531	LYS	2.8
1	A	530	VAL	2.8
1	B	377	SER	2.8
1	A	381	VAL	2.7
1	B	111	LEU	2.7
1	B	148	GLN	2.7
1	B	536	LEU	2.7
1	B	330	GLN	2.6
1	B	101	PHE	2.6
1	A	95	HIS	2.6
1	B	160	ILE	2.6
1	B	505	ARG	2.6
1	B	100	LYS	2.6
1	B	382	TYR	2.6
1	B	102	GLY	2.6
1	B	32	ARG	2.6
1	B	516	GLY	2.5
1	B	352	ASP	2.5
1	B	537	THR	2.5
1	B	402	HIS	2.5
1	B	8	GLY	2.4
1	A	94	PRO	2.4
1	A	32	ARG	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	469	LEU	2.3
1	B	528	TRP	2.3
1	A	512	LEU	2.3
1	B	15	ALA	2.3
1	B	548	SER	2.3
1	A	43	ARG	2.3
1	B	378	GLY	2.3
1	A	16	ALA	2.2
1	A	376	ALA	2.2
1	B	331	GLU	2.2
1	A	402	HIS	2.2
1	A	501	ARG	2.2
1	B	105	ALA	2.1
1	A	431	SER	2.1
1	A	8	GLY	2.1
1	A	15	ALA	2.1
1	A	352	ASP	2.1
1	A	51	LYS	2.1
1	B	97	ALA	2.1
1	B	311	CYS	2.1
1	A	102	GLY	2.1
1	B	213	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	B42	A	579	36/36	0.96	0.08	35,37,39,40	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	B42	B	579	36/36	0.96	0.08	37,40,41,41	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around B42 A 579:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around B42 B 579:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.