



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 11:08 PM UTC

PDB ID : 4GYW / pdb_00004gyw
Title : Crystal structure of human O-GlcNAc Transferase in complex with UDP and a glycopeptide
Authors : Lazarus, M.B.; Jiang, J.; Gloster, T.M.; Zandberg, W.F.; Vocadlo, D.J.; Walker, S.
Deposited on : 2012-09-05
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	723	10% 91% 5% .
1	C	723	11% 87% 6% 7%
2	B	14	14% 86% 7% 7%
2	D	14	14% 79% 21%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 12156 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UDP-N-acetylglucosamine--peptide N-acetylglucosaminyltransferase 110 kDa subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	695	Total	C	N	O	S	0	8	0
			5536	3514	966	1019	37			
1	C	674	Total	C	N	O	S	0	7	0
			5367	3417	934	978	38			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	309	GLY	-	expression tag	UNP O15294
A	310	PRO	-	expression tag	UNP O15294
A	311	GLY	-	expression tag	UNP O15294
A	312	SER	-	expression tag	UNP O15294
C	309	GLY	-	expression tag	UNP O15294
C	310	PRO	-	expression tag	UNP O15294
C	311	GLY	-	expression tag	UNP O15294
C	312	SER	-	expression tag	UNP O15294

- Molecule 2 is a protein called Casein kinase II subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	14	Total	C	N	O	S	0	0	0
			95	58	15	20	2			
2	D	14	Total	C	N	O	S	0	0	0
			95	58	15	20	2			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	13	TYR	-	expression tag	UNP P68400
D	13	TYR	-	expression tag	UNP P68400

-
- The diagram illustrates the chemical structure of UDP (Uridine Diphosphate). It consists of a uracil base (a six-membered ring with two nitrogen atoms, N1 and N3) attached to a ribose sugar (a five-membered ring). The ribose sugar is linked to a diphosphate group (two phosphate groups connected by an oxygen atom). The uracil base is labeled with atoms N1, N3, C2, C4, C5, and C6. The ribose sugar is labeled with atoms C1', C2', C3', C4', and C5'. The diphosphate group is labeled with atoms O1A, O1B, O2A, O2B, O3A, and O3B. The structure is shown in a 3D representation with wedges and dashes indicating stereochemistry.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 25	C 9	N 2	O 12	P 2	0	0
3	C	1	Total 25	C 9	N 2	O 12	P 2	0	0

-
- Diagram illustrating the Lewis structure of the sulfate ion (SO_4^{2-}). The central sulfur atom (S) is bonded to four oxygen atoms (O). The top and bottom oxygen atoms are double-bonded to sulfur, while the left and right oxygen atoms are single-bonded to sulfur. Each single-bonded oxygen atom carries a negative charge (O^-). The sulfur atom has a formal charge of +6, indicated by the 'S' below it. The oxygen atoms are labeled O1, O2, O3, and O4 in green text.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	D	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	487	Total	O	0	0
			487	487		
6	B	19	Total	O	0	0
			19	19		
6	C	450	Total	O	0	0
			450	450		
6	D	14	Total	O	0	0
			14	14		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: UDP-N-acetylglucosamine--peptide N-acetylglucosaminyltransferase 110 kDa subunit





- Molecule 1: UDP-N-acetylglucosamine--peptide N-acetylglucosaminyltransferase 110 kDa subunit



- Molecule 2: Casein kinase II subunit alpha



● Molecule 2: Casein kinase II subunit alpha



4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	99.00Å 137.77Å 153.57Å 90.00° 102.90° 90.00°	Depositor
Resolution (Å)	43.90 – 1.70 43.90 – 1.70	Depositor EDS
% Data completeness (in resolution range)	94.2 (43.90-1.70) 94.2 (43.90-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.24 (at 1.70Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7_650)	Depositor
R, R_{free}	0.187 , 0.205 0.181 , 0.199	Depositor DCC
R_{free} test set	10341 reflections (4.59%)	wwPDB-VP
Wilson B-factor (Å ²)	18.9	Xtriage
Anisotropy	0.149	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 44.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12156	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, NAG, UDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/5688	0.69	2/7713 (0.0%)
1	C	0.38	0/5515	0.69	4/7478 (0.1%)
2	B	0.43	0/97	0.63	0/131
2	D	0.37	0/97	0.65	0/131
All	All	0.38	0/11397	0.69	6/15453 (0.0%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	858	VAL	CA-C-N	5.47	124.93	119.24
1	C	858	VAL	C-N-CA	5.47	124.93	119.24
1	C	608	ILE	CA-C-N	5.29	125.10	119.28
1	C	608	ILE	C-N-CA	5.29	125.10	119.28
1	A	608	ILE	CA-C-N	5.28	124.91	119.05
1	A	608	ILE	C-N-CA	5.28	124.91	119.05

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5536	0	5520	22	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	5367	0	5363	24	0
2	B	95	0	87	2	0
2	D	95	0	87	3	0
3	A	25	0	11	0	0
3	C	25	0	11	0	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0
4	D	5	0	0	0	0
5	B	14	0	13	0	0
5	D	14	0	13	1	0
6	A	487	0	0	5	0
6	B	19	0	0	0	0
6	C	450	0	0	3	0
6	D	14	0	0	0	0
All	All	12156	0	11105	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:726:LEU:CD2	1:A:819:VAL:HG22	2.26	0.66
2:B:25:MET:HE2	2:B:25:MET:HA	1.80	0.64
2:D:25:MET:HA	2:D:25:MET:HE2	1.81	0.63
1:A:706:LYS:HE3	6:A:1654:HOH:O	2.04	0.58
1:A:986:LYS:HG3	1:A:990[B]:GLN:OE1	2.04	0.56
1:C:443:ARG:NH2	6:C:1621:HOH:O	2.36	0.56
1:C:726[A]:LEU:CD2	1:C:819:VAL:HG22	2.37	0.55
1:C:607:GLN:NE2	6:C:1582:HOH:O	2.39	0.54
1:C:986:LYS:HG3	1:C:990[B]:GLN:OE1	2.09	0.53
1:A:558:HIS:CG	1:A:559:PRO:HD2	2.46	0.50
1:A:376:GLU:HG2	6:A:1424:HOH:O	2.12	0.50
1:A:402:GLN:O	1:A:406:GLN:HG2	2.12	0.49
1:C:772:ILE:HG23	1:C:789:ILE:HD13	1.94	0.48
1:A:325:ASN:O	1:A:329:GLU:HG2	2.14	0.47
1:C:337:VAL:O	1:C:341:ARG:HG3	2.14	0.47
1:A:770:ASN:ND2	6:A:1682:HOH:O	2.47	0.47
1:A:496:HIS:CE1	2:B:23:ALA:HB1	2.51	0.46
1:A:954:GLY:O	1:A:986:LYS:HE2	2.16	0.46
1:C:516:ARG:NH2	1:C:519:ASN:OD1	2.48	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:362:GLN:HG2	6:A:1639:HOH:O	2.15	0.46
1:A:941:LEU:HD23	1:A:941:LEU:C	2.41	0.45
1:C:501:MET:HE1	5:D:101:NAG:H81	1.98	0.45
1:C:402:GLN:O	1:C:406:GLN:HG2	2.17	0.45
1:A:359:SER:O	1:A:363:GLN:HG3	2.17	0.44
1:C:941:LEU:C	1:C:941:LEU:HD23	2.43	0.44
1:C:517:HIS:NE2	6:C:1569:HOH:O	2.31	0.44
1:C:347:PHE:CE1	1:C:349:GLU:HB2	2.53	0.44
1:A:516:ARG:NH2	6:A:1680:HOH:O	2.51	0.43
1:C:496:HIS:CE1	2:D:23:ALA:HB1	2.53	0.43
1:C:558:HIS:CG	1:C:559:PRO:HD2	2.55	0.42
1:C:408:TYR:CZ	1:C:424:ASN:HB3	2.55	0.42
1:C:629:MET:O	1:C:654:GLY:HA3	2.20	0.42
1:A:332:ASN:CG	1:A:335:GLU:HB2	2.44	0.42
1:A:486:GLN:OE1	1:A:493:PRO:HA	2.19	0.42
1:A:515[A]:GLU:OE2	1:A:519:ASN:ND2	2.51	0.42
1:C:361:LEU:HD13	1:C:369:GLU:HB3	2.02	0.41
1:C:576:PHE:CZ	1:C:1007:GLU:HB3	2.55	0.41
1:C:844:ASP:HB2	1:C:845:PRO:HD2	2.01	0.41
1:A:844:ASP:HB2	1:A:845:PRO:HD2	2.01	0.41
1:A:603:ILE:N	1:A:603:ILE:HD12	2.35	0.41
1:C:418:PHE:CE1	1:C:420:ASP:HB2	2.55	0.41
1:C:873:GLU:N	1:C:874:PRO:HD2	2.36	0.41
1:A:850[B]:MET:HE2	1:A:850[B]:MET:HB2	1.89	0.41
1:A:629:MET:O	1:A:654:GLY:HA3	2.21	0.40
1:C:388:TYR:O	1:C:407:CYS:HB3	2.21	0.40
1:C:633:THR:O	2:D:24:ASN:HB3	2.21	0.40
1:C:357:LEU:O	1:C:361:LEU:HG	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	698/723 (96%)	686 (98%)	12 (2%)	0	100	100
1	C	675/723 (93%)	663 (98%)	12 (2%)	0	100	100
2	B	12/14 (86%)	12 (100%)	0	0	100	100
2	D	12/14 (86%)	12 (100%)	0	0	100	100
All	All	1397/1474 (95%)	1373 (98%)	24 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	606/618 (98%)	601 (99%)	5 (1%)	73	65
1	C	586/618 (95%)	585 (100%)	1 (0%)	87	84
2	B	11/11 (100%)	10 (91%)	1 (9%)	9	2
2	D	11/11 (100%)	11 (100%)	0	100	100
All	All	1214/1258 (96%)	1207 (99%)	7 (1%)	78	72

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	525	ILE
1	A	527	VAL
1	A	744	VAL
1	A	914	THR
1	A	926	VAL
2	B	25	MET
1	C	926	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	321	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	356	ASN
1	A	363	GLN
1	A	364	GLN
1	A	802	GLN
1	A	881	GLN
1	C	607	GLN
1	C	691	HIS
1	C	784	GLN
1	C	805	ASN
1	C	1012	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	B	101	2	14,14,15	0.52	0	17,19,21	0.81	1 (5%)
4	SO4	A	1102	-	4,4,4	0.25	0	6,6,6	0.07	0
3	UDP	C	1101	-	25,26,26	1.03	2 (8%)	38,40,40	1.45	4 (10%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	D	102	-	4,4,4	0.27	0	6,6,6	0.13	0
4	SO4	B	102	-	4,4,4	0.24	0	6,6,6	0.06	0
3	UDP	A	1101	-	25,26,26	1.04	1 (4%)	38,40,40	1.39	4 (10%)
5	NAG	D	101	2	14,14,15	0.43	0	17,19,21	0.82	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	UDP	C	1101	-	-	4/16/32/32	0/2/2/2
3	UDP	A	1101	-	-	4/16/32/32	0/2/2/2
5	NAG	B	101	2	-	0/6/23/26	0/1/1/1
5	NAG	D	101	2	-	1/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1101	UDP	PA-O3A	2.22	1.61	1.59
3	C	1101	UDP	C6-C5	2.07	1.39	1.35
3	C	1101	UDP	PB-O2B	-2.01	1.47	1.54

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1101	UDP	C4-N3-C2	-4.85	120.60	126.61
3	A	1101	UDP	C4-N3-C2	-4.54	120.97	126.61
3	C	1101	UDP	C5-C4-N3	4.10	120.55	114.80
3	A	1101	UDP	C5-C4-N3	3.69	119.97	114.80
3	C	1101	UDP	N3-C2-N1	3.22	119.08	114.89
3	A	1101	UDP	N3-C2-N1	3.04	118.85	114.89
3	A	1101	UDP	O4-C4-C5	-2.45	120.93	125.16
3	C	1101	UDP	O4-C4-C5	-2.42	120.98	125.16
5	B	101	NAG	C1-O5-C5	2.14	115.06	112.19

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1101	UDP	O4'-C4'-C5'-O5'

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	C	1101	UDP	C3'-C4'-C5'-O5'
3	A	1101	UDP	C3'-C4'-C5'-O5'
3	C	1101	UDP	O4'-C4'-C5'-O5'
3	A	1101	UDP	PA-O3A-PB-O1B
3	C	1101	UDP	PA-O3A-PB-O1B
3	A	1101	UDP	PA-O3A-PB-O3B
3	C	1101	UDP	PA-O3A-PB-O3B
5	D	101	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	101	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	695/723 (96%)	0.37	70 (10%) 12 13	8, 20, 44, 91	8 (1%)
1	C	674/723 (93%)	0.50	81 (12%) 9 8	8, 19, 47, 119	7 (1%)
2	B	14/14 (100%)	0.84	2 (14%) 6 5	14, 19, 57, 64	0
2	D	14/14 (100%)	0.70	2 (14%) 6 5	13, 20, 53, 59	0
All	All	1397/1474 (94%)	0.44	155 (11%) 10 10	8, 19, 46, 119	15 (1%)

All (155) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	346	VAL	13.1
1	C	339	LEU	13.1
1	C	336	ALA	12.9
1	C	343	ALA	12.5
1	C	337	VAL	10.4
1	C	340	TYR	9.7
1	A	314	PRO	9.1
1	C	347	PHE	8.3
1	A	713	PHE	7.6
1	C	348	PRO	7.4
1	C	344	LEU	6.9
1	A	714	LYS	6.8
1	C	342	LYS	6.7
1	C	714	LYS	6.4
1	C	360	VAL	6.3
1	A	772	ILE	6.3
1	A	331	GLY	6.0
1	C	350	PHE	6.0
1	C	713	PHE	6.0
1	C	746	MET	5.9
1	C	718	HIS	5.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	26	MET	5.1
1	A	718	HIS	5.1
1	A	763	ASN	5.0
1	C	341	ARG	4.9
1	C	356	ASN	4.8
1	A	315	THR	4.8
1	C	1028	LYS	4.6
1	A	769	MET	4.6
1	C	383	THR	4.5
2	D	26	MET	4.5
1	C	860	ASN	4.1
1	A	860	ASN	4.1
2	B	25	MET	4.1
1	C	345	GLU	4.0
1	A	490	ASN	4.0
1	A	323	LEU	3.9
1	C	359	SER	3.9
2	D	25	MET	3.9
1	A	330	GLN	3.8
1	A	742	LYS	3.8
1	C	353	ALA	3.7
1	A	745	LYS	3.7
1	A	771	THR	3.7
1	A	744	VAL	3.7
1	C	352	ALA	3.7
1	A	431	ASP	3.7
1	A	433	GLY	3.6
1	C	772	ILE	3.6
1	A	491	ARG	3.6
1	A	719	ILE	3.6
1	C	771	THR	3.6
1	A	488	GLU	3.6
1	C	338	ARG	3.5
1	A	773	ALA	3.5
1	A	332	ASN	3.5
1	A	739	PRO	3.5
1	A	527	VAL	3.4
1	C	358	ALA	3.4
1	C	777	ILE	3.4
1	C	519	ASN	3.4
1	A	740	ASP	3.4
1	A	520	LEU	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	745	LYS	3.3
1	C	712	ASP	3.3
1	C	361	LEU	3.3
1	C	432	SER	3.3
1	C	351	ALA	3.3
1	A	743	ILE	3.3
1	A	720	TYR	3.2
1	C	769	MET	3.1
1	A	328	ARG	3.0
1	C	763	ASN	3.0
1	A	333	ILE	3.0
1	C	357	LEU	3.0
1	C	529	HIS	2.9
1	A	329	GLU	2.9
1	A	492	LEU	2.9
1	A	887	GLN	2.9
1	C	830	ASP	2.9
1	C	527	VAL	2.9
1	C	362	GLN	2.9
1	A	770	ASN	2.9
1	A	774	GLU	2.9
1	A	347	PHE	2.8
1	A	680	GLU	2.8
1	C	381	SER	2.8
1	C	528	LEU	2.8
1	C	677	GLU	2.8
1	A	537	LYS	2.8
1	C	678	VAL	2.8
1	C	887	GLN	2.8
1	C	740	ASP	2.8
1	C	491	ARG	2.7
1	C	431	ASP	2.7
1	A	432	SER	2.7
1	A	830	ASP	2.7
1	A	346	VAL	2.7
1	C	720	TYR	2.7
1	A	543	ASP	2.7
1	A	326	ILE	2.6
1	C	380	ILE	2.6
1	C	680	GLU	2.6
1	C	773	ALA	2.6
1	A	327	LYS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	401	VAL	2.5
1	C	365	GLY	2.5
1	C	774	GLU	2.5
1	C	543	ASP	2.5
1	A	777	ILE	2.5
1	C	719	ILE	2.5
1	C	739	PRO	2.5
1	A	363	GLN	2.5
1	C	884	GLY	2.5
1	C	363	GLN	2.4
1	A	609	PRO	2.4
1	C	382	PRO	2.4
1	A	319[A]	SER	2.4
1	A	489	LYS	2.4
1	C	355	SER	2.3
1	C	433	GLY	2.3
1	A	534	GLU	2.3
1	A	1015	GLU	2.3
1	A	712	ASP	2.3
1	A	859	PRO	2.3
1	C	676	ALA	2.3
1	C	384	PHE	2.3
1	A	526	ASN	2.3
1	C	490	ASN	2.3
1	C	770	ASN	2.3
1	A	335	GLU	2.3
1	A	768	PRO	2.3
1	A	384	PHE	2.3
1	C	447	LYS	2.2
1	A	678	VAL	2.2
1	C	744	VAL	2.2
1	A	775	ALA	2.2
1	A	507	HIS	2.2
1	C	742	LYS	2.2
1	A	1019	ALA	2.2
1	A	607	GLN	2.2
1	A	677	GLU	2.2
1	A	532	PRO	2.2
1	A	334	GLU	2.1
1	C	364	GLN	2.1
1	C	379	ARG	2.1
1	A	620	GLN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	519	ASN	2.1
1	C	662	LEU	2.1
1	C	609	PRO	2.1
1	C	378	ILE	2.1
1	C	743	ILE	2.1
1	A	541	LEU	2.0
1	C	399	GLN	2.0
1	C	620	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	SO4	A	1102	5/5	0.87	0.17	57,57,59,60	0
4	SO4	D	102	5/5	0.92	0.12	43,45,48,50	0
4	SO4	B	102	5/5	0.94	0.10	43,43,47,48	0
5	NAG	B	101	14/15	0.95	0.07	12,14,18,18	0
5	NAG	D	101	14/15	0.96	0.06	11,14,17,19	0
3	UDP	C	1101	25/25	0.99	0.04	9,11,13,15	0
3	UDP	A	1101	25/25	0.99	0.04	8,10,13,15	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around UDP C 1101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around UDP A 1101:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.