



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 10:49 PM UTC

PDB ID : 6GY2 / pdb_00006gy2
Title : Crystal structure of human Plk1-PBD in complex with WSSS-LATPPTLSSpTVLI phosphopeptide from BRCA2
Authors : Miron, S.; Ropars, V.; Zinn-Justin, S.
Deposited on : 2018-06-28
Resolution : 3.11 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1816 (3.14-3.10)
Clashscore	190562	1906 (3.14-3.10)
Ramachandran outliers	187476	1802 (3.14-3.10)
Sidechain outliers	187428	1802 (3.14-3.10)
RSRZ outliers	180081	1816 (3.14-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	245	
1	B	245	
2	C	17	
2	D	17	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3842 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase PLK1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	224	Total	C	N	O	S	1	0	0
			1818	1152	314	341	11			
1	B	224	Total	C	N	O	S	3	0	0
			1822	1153	317	341	11			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	359	ALA	-	expression tag	UNP P53350
A	360	MET	-	expression tag	UNP P53350
A	361	ASP	-	expression tag	UNP P53350
A	362	PRO	-	expression tag	UNP P53350
A	363	GLU	-	expression tag	UNP P53350
A	364	PHE	-	expression tag	UNP P53350
B	359	ALA	-	expression tag	UNP P53350
B	360	MET	-	expression tag	UNP P53350
B	361	ASP	-	expression tag	UNP P53350
B	362	PRO	-	expression tag	UNP P53350
B	363	GLU	-	expression tag	UNP P53350
B	364	PHE	-	expression tag	UNP P53350

- Molecule 2 is a protein called Phosphopeptide of BRCA2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	12	Total	C	N	O	P	0	0	0
			87	54	12	20	1			
2	D	12	Total	C	N	O	P	0	0	0
			87	54	12	20	1			

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	10	Total	O	0	0
			10	10		
4	B	9	Total	O	0	0
			9	9		
4	C	1	Total	O	0	0
			1	1		
4	D	2	Total	O	0	0
			2	2		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serine/threonine-protein kinase PLK1




- Molecule 1: Serine/threonine-protein kinase PLK1



- Molecule 2: Phosphopeptide of BRCA2



- Molecule 2: Phosphopeptide of BRCA2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	50.00Å 56.04Å 61.03Å 80.79° 79.23° 65.05°	Depositor
Resolution (Å)	22.57 – 3.11 22.57 – 3.11	Depositor EDS
% Data completeness (in resolution range)	93.7 (22.57-3.11) 93.4 (22.57-3.11)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.65 (at 3.13Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.187 , 0.215 0.199 , 0.226	Depositor DCC
R_{free} test set	474 reflections (4.47%)	wwPDB-VP
Wilson B-factor (Å ²)	73.3	Xtriage
Anisotropy	0.458	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 52.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3842	wwPDB-VP
Average B, all atoms (Å ²)	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/1856	1.18	6/2510 (0.2%)
1	B	0.67	0/1860	1.21	6/2514 (0.2%)
2	C	0.84	0/76	1.19	0/104
2	D	0.78	0/76	1.16	0/104
All	All	0.66	0/3868	1.20	12/5232 (0.2%)

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	500	ARG	CA-C-N	6.34	129.63	120.38
1	A	500	ARG	C-N-CA	6.34	129.63	120.38
1	B	395	VAL	N-CA-CB	5.99	117.06	110.53
1	B	503	ASP	CA-CB-CG	5.76	118.36	112.60
1	B	559	PHE	CA-CB-CG	5.65	119.45	113.80
1	A	559	PHE	CA-CB-CG	5.54	119.34	113.80
1	A	395	VAL	N-CA-CB	5.43	116.44	110.53
1	A	503	ASP	CA-CB-CG	5.31	117.91	112.60
1	B	396	ARG	CA-C-N	5.24	127.55	120.38
1	B	396	ARG	C-N-CA	5.24	127.55	120.38
1	B	447	ASP	CA-CB-CG	5.09	117.69	112.60
1	A	447	ASP	CA-CB-CG	5.03	117.63	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1818	0	1802	8	0
1	B	1822	0	1806	6	0
2	C	87	0	90	0	0
2	D	87	0	90	0	0
3	A	6	0	8	0	0
4	A	10	0	0	1	0
4	B	9	0	0	0	0
4	C	1	0	0	0	0
4	D	2	0	0	0	0
All	All	3842	0	3796	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:372:CYS:HB2	1:B:589:LYS:HE3	1.88	0.56
1:A:412:SER:HB2	1:A:492:LYS:HG3	1.97	0.45
1:B:412:SER:HB2	1:B:492:LYS:HG3	1.99	0.45
1:B:392:ARG:HB2	1:B:395:VAL:HG12	1.98	0.45
1:A:392:ARG:HB2	1:A:395:VAL:HG12	1.99	0.44
1:A:370:VAL:HG11	4:A:809:HOH:O	2.17	0.43
1:B:452:GLN:HE21	1:B:460:GLU:HG2	1.83	0.41
1:A:567:LEU:HD22	1:A:572:CYS:HB3	2.02	0.41
1:B:567:LEU:HD22	1:B:572:CYS:HB3	2.02	0.41
1:A:452:GLN:HE21	1:A:460:GLU:HG2	1.84	0.41
1:A:478:LEU:O	1:A:482:PHE:HD1	2.03	0.41
1:A:551:THR:HG21	1:A:559:PHE:CZ	2.56	0.41
1:B:478:LEU:O	1:B:482:PHE:HD1	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	222/245 (91%)	215 (97%)	7 (3%)	0	100	100
1	B	222/245 (91%)	216 (97%)	6 (3%)	0	100	100
2	C	9/17 (53%)	9 (100%)	0	0	100	100
2	D	9/17 (53%)	9 (100%)	0	0	100	100
All	All	462/524 (88%)	449 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	203/220 (92%)	194 (96%)	9 (4%)	25	54
1	B	203/220 (92%)	194 (96%)	9 (4%)	25	54
2	C	10/15 (67%)	9 (90%)	1 (10%)	7	27
2	D	10/15 (67%)	10 (100%)	0	100	100
All	All	426/470 (91%)	407 (96%)	19 (4%)	24	54

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	371	ASP
1	A	372	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	423	LEU
1	A	463	LEU
1	A	470	ASN
1	A	519	SER
1	A	578	SER
1	A	588	ASP
1	A	591	LEU
1	B	371	ASP
1	B	423	LEU
1	B	463	LEU
1	B	470	ASN
1	B	483	ARG
1	B	519	SER
1	B	578	SER
1	B	588	ASP
1	B	591	LEU
2	C	208	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	382	HIS
1	A	426	GLN
1	A	452	GLN
1	B	382	HIS
1	B	426	GLN
1	B	452	GLN
1	B	484	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	TPO	D	207	2	8,10,11	1.08	0	10,14,16	1.34	2 (20%)
2	TPO	C	207	2	8,10,11	1.22	1 (12%)	10,14,16	1.24	2 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TPO	D	207	2	-	1/9/11/13	-
2	TPO	C	207	2	-	1/9/11/13	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	207	TPO	P-OG1	-2.49	1.55	1.59

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	207	TPO	P-OG1-CB	-2.09	117.65	123.33
2	C	207	TPO	P-OG1-CB	-2.05	117.75	123.33
2	C	207	TPO	O3P-P-OG1	2.05	113.82	105.85
2	D	207	TPO	O3P-P-OG1	2.01	113.69	105.85

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	207	TPO	CB-OG1-P-O2P
2	D	207	TPO	CB-OG1-P-O2P

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	701	-	5,5,5	0.06	0	5,5,5	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	701	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	701	GOL	C1-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	224/245 (91%)	-0.03	0 100 100	47, 77, 107, 136	2 (0%)
1	B	224/245 (91%)	-0.08	2 (0%) 81 63	51, 78, 105, 131	4 (1%)
2	C	11/17 (64%)	0.12	0 100 100	67, 72, 94, 95	0
2	D	11/17 (64%)	0.01	0 100 100	70, 74, 85, 86	0
All	All	470/524 (89%)	-0.05	2 (0%) 88 76	47, 78, 106, 136	6 (1%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	544	CYS	2.6
1	B	503	ASP	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	TPO	C	207	11/12	0.95	0.06	69,71,73,74	0
2	TPO	D	207	11/12	0.96	0.07	70,72,75,76	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	A	701	6/6	0.62	0.18	83,83,88,88	0

6.5 Other polymers

There are no such residues in this entry.