



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:20 PM UTC

PDB ID : 3GZA / pdb_00003gza
Title : Crystal structure of putative alpha-L-fucosidase (NP_812709.1) from BACTEROIDES THETA IOTAOMICRON VPI-5482 at 1.60 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2009-04-06
Resolution : 1.60 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	443	16% 96% ..
1	B	443	12% 91% 6% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	UNL	B	461	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 8405 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called putative alpha-L-fucosidase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	438	Total	C	N	O	S	Se	0	15	0
			3576	2279	628	652	11	6			
1	B	431	Total	C	N	O	S	Se	0	18	0
			3519	2244	614	643	11	7			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP Q8A169
B	0	GLY	-	expression tag	UNP Q8A169

- Molecule 2 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	1
			15	8	2	4	1		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Na	0	0
			1	1		
4	B	1	Total	Na	0	0
			1	1		

- Molecule 5 is UNKNOWN LIGAND (CCD ID: UNL) (formula:).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total O 9 9	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	667	Total O 677 677	0	11
6	B	567	Total O 575 575	0	14


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: putative alpha-L-fucosidase



- Molecule 1: putative alpha-L-fucosidase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	117.05Å 81.53Å 111.57Å 90.00° 103.55° 90.00°	Depositor
Resolution (Å)	28.07 – 1.60 28.07 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.7 (28.07-1.60) 99.6 (28.07-1.60)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.54 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.2.0019, PHENIX	Depositor
R, R_{free}	0.139 , 0.168 (Not available) , 0.267	Depositor DCC
R_{free} test set	6726 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	14.8	Xtriage
Anisotropy	0.445	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8405	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, UNL, EDO, EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	0/3710	0.92	1/5024 (0.0%)
1	B	0.84	1/3656 (0.0%)	0.98	9/4950 (0.2%)
All	All	0.84	1/7366 (0.0%)	0.95	10/9974 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	62	ILE	N-CA	-8.62	1.35	1.46

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	61[A]	ARG	N-CA-C	8.36	120.48	111.36
1	B	61[B]	ARG	N-CA-C	8.36	120.48	111.36
1	B	61[A]	ARG	CA-C-N	5.89	132.58	121.97
1	B	61[A]	ARG	C-N-CA	5.89	132.58	121.97
1	B	61[B]	ARG	CA-C-N	5.89	132.58	121.97
1	B	61[B]	ARG	C-N-CA	5.89	132.58	121.97
1	B	342	ILE	N-CA-C	-5.54	103.52	108.95
1	B	299	GLU	N-CA-C	5.36	117.48	107.99
1	B	62	ILE	CA-C-O	-5.34	114.11	120.78
1	A	285	ALA	N-CA-C	5.19	117.97	110.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3576	0	3469	9	0
1	B	3519	0	3412	26	0
2	A	15	0	18	0	0
3	A	12	0	18	0	0
3	B	20	0	30	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	B	9	0	0	5	0
6	A	677	0	0	3	1
6	B	575	0	0	3	1
All	All	8405	0	6947	40	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:461:UNL:O4	5:B:461:UNL:O5	1.71	1.09
5:B:461:UNL:O7	5:B:461:UNL:O8	1.74	1.04
5:B:461:UNL:O2	5:B:461:UNL:O3	1.73	1.03
5:B:461:UNL:O5	5:B:461:UNL:O6	1.79	1.00
1:B:388[B]:CYS:SG	1:B:405:ILE:HD13	2.06	0.96
1:B:146:ILE:HD12	1:B:196[B]:ILE:HD11	1.53	0.90
5:B:461:UNL:O2	5:B:461:UNL:O1	1.90	0.88
1:B:388[B]:CYS:SG	1:B:405:ILE:CD1	2.68	0.82
1:A:388[B]:CYS:SG	1:A:405:ILE:HD13	2.24	0.78
1:B:270:GLU:O	1:B:273[A]:LYS:HG2	1.85	0.76
1:A:146:ILE:HD12	1:A:196[B]:ILE:HD11	1.69	0.74
1:A:388[B]:CYS:SG	1:A:405:ILE:CD1	2.81	0.69
1:B:270:GLU:C	1:B:270:GLU:CD	2.61	0.68
6:A:541:HOH:O	1:B:119[B]:LYS:CD	2.41	0.68
1:B:61[B]:ARG:HA	6:B:1033[B]:HOH:O	1.95	0.65
1:B:273[A]:LYS:HA	6:B:509:HOH:O	1.97	0.64
1:B:388[A]:CYS:SG	1:B:438:LEU:HD13	2.39	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:273[B]:LYS:HA	6:B:509:HOH:O	1.99	0.62
1:B:438:LEU:HD12	1:B:456:ALA:HB2	1.84	0.59
1:B:146:ILE:CD1	1:B:196[B]:ILE:HD11	2.27	0.59
1:B:59:ASN:C	1:B:61[B]:ARG:O	2.45	0.59
1:A:388[A]:CYS:SG	1:A:438:LEU:HD13	2.43	0.59
1:B:273[A]:LYS:HG3	1:B:274:HIS:CD2	2.40	0.56
1:B:270:GLU:HA	1:B:273[A]:LYS:HE3	1.87	0.55
1:A:390:ILE:HD11	1:A:405:ILE:HD11	1.91	0.53
1:A:314:ASN:CB	6:A:611:HOH:O	2.58	0.52
1:B:368:ILE:HG22	1:B:377:LEU:HD13	1.93	0.51
1:A:388[A]:CYS:SG	1:A:438:LEU:CD1	2.99	0.50
1:B:365:ILE:CD1	1:B:385[B]:VAL:HG12	2.41	0.50
1:A:162:GLU:HG2	6:A:699:HOH:O	2.12	0.49
1:B:39[A]:MSE:HE3	1:B:330:LEU:HD23	1.95	0.48
1:B:270:GLU:CD	1:B:271:LEU:N	2.72	0.47
1:B:390:ILE:HD11	1:B:405:ILE:HD11	1.97	0.47
1:B:270:GLU:OE1	1:B:271:LEU:N	2.48	0.46
1:B:337:ASP:HB2	1:B:338:PRO:CD	2.47	0.44
1:A:379:LEU:HD11	1:A:385[A]:VAL:HG21	1.99	0.44
1:B:62:ILE:N	1:B:62:ILE:HD13	2.33	0.44
1:B:59:ASN:O	1:B:61[B]:ARG:O	2.36	0.43
1:B:204:ASP:CG	1:B:229[B]:ILE:HD13	2.43	0.43
1:B:204:ASP:CG	1:B:229[B]:ILE:CD1	2.92	0.42

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:977:HOH:O	6:B:925:HOH:O[4_656]	2.14	0.06

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	452/443 (102%)	433 (96%)	19 (4%)	0	100	100
1	B	445/443 (100%)	427 (96%)	18 (4%)	0	100	100
All	All	897/886 (101%)	860 (96%)	37 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	375/370 (101%)	375 (100%)	0	100	100
1	B	369/370 (100%)	369 (100%)	0	100	100
All	All	744/740 (100%)	744 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	60	ASN
1	A	83	GLN
1	A	404	GLN
1	B	59	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 2 are monoatomic and 1 is unknown - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	B	464	-	3,3,3	0.35	0	2,2,2	1.12	0
3	EDO	A	470	-	3,3,3	0.44	0	2,2,2	0.26	0
2	EPE	A	462[A]	-	15,15,15	1.13	1 (6%)	19,20,20	1.96	5 (26%)
3	EDO	A	468	-	3,3,3	0.54	0	2,2,2	0.20	0
3	EDO	B	466	-	3,3,3	0.97	0	2,2,2	0.84	0
3	EDO	B	467	-	3,3,3	0.67	0	2,2,2	0.65	0
3	EDO	B	465	-	3,3,3	0.47	0	2,2,2	0.26	0
3	EDO	B	469	-	3,3,3	0.53	0	2,2,2	0.08	0
3	EDO	A	463	-	3,3,3	0.22	0	2,2,2	0.81	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	464	-	-	0/1/1/1	-
3	EDO	A	470	-	-	0/1/1/1	-
2	EPE	A	462[A]	-	-	0/9/19/19	0/1/1/1
3	EDO	A	468	-	-	1/1/1/1	-
3	EDO	B	466	-	-	1/1/1/1	-
3	EDO	B	467	-	-	0/1/1/1	-
3	EDO	B	465	-	-	0/1/1/1	-
3	EDO	B	469	-	-	0/1/1/1	-
3	EDO	A	463	-	-	0/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	462[A]	EPE	C10-S	3.52	1.82	1.77

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	462[A]	EPE	O1S-S-C10	-5.15	98.94	106.73
2	A	462[A]	EPE	C3-C2-N1	-3.51	103.58	110.65
2	A	462[A]	EPE	C9-N1-C6	3.06	119.39	111.24
2	A	462[A]	EPE	C7-N4-C5	2.29	117.33	111.24
2	A	462[A]	EPE	O3S-S-C10	2.03	109.97	106.00

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	468	EDO	O1-C1-C2-O2
3	B	466	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	432/443 (97%)	1.14	71 (16%)	4 4	9, 26, 33, 48	15 (3%)
1	B	425/443 (95%)	1.18	55 (12%)	7 7	13, 26, 37, 60	17 (4%)
All	All	857/886 (96%)	1.16	126 (14%)	6 5	9, 26, 35, 60	32 (3%)

All (126) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	269	LEU	9.3
1	B	55	TYR	6.8
1	B	58	GLY	6.2
1	B	272	LEU	5.3
1	B	62	ILE	4.8
1	B	268	GLN	4.7
1	B	380	GLY	4.4
1	B	273[A]	LYS	4.1
1	B	271	LEU	4.0
1	B	259	HIS	4.0
1	B	258	HIS	3.9
1	B	60	ASN	3.8
1	B	267	ASP	3.3
1	B	56	GLY	3.2
1	A	385[A]	VAL	3.2
1	A	105	LEU	3.2
1	A	411	GLY	3.2
1	B	270	GLU	3.1
1	A	380	GLY	3.0
1	B	207	ALA	3.0
1	A	120	TRP	3.0
1	B	206	ARG	3.0
1	A	103	PHE	3.0
1	A	459	VAL	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	266	ILE	2.9
1	A	106	TRP	2.9
1	A	168	ALA	2.8
1	A	114	CYS	2.8
1	B	53	ILE	2.8
1	A	110	VAL	2.8
1	A	68	TYR	2.8
1	B	57	GLN	2.8
1	B	65	ILE	2.8
1	A	125	GLY	2.8
1	A	413	TRP	2.7
1	A	379	LEU	2.7
1	A	127	ILE	2.7
1	B	70	ILE	2.7
1	B	61[A]	ARG	2.7
1	A	118	VAL	2.7
1	A	433	VAL	2.7
1	A	115	LEU	2.7
1	A	154	LEU	2.7
1	A	219	TYR	2.7
1	A	102	GLY	2.7
1	A	122[A]	ASP	2.7
1	A	261[A]	ARG	2.6
1	A	153	LEU	2.6
1	B	379	LEU	2.6
1	A	113	TYR	2.6
1	B	303	GLU	2.6
1	B	166	ALA	2.6
1	A	47	LEU	2.6
1	B	76	LEU	2.6
1	A	70	ILE	2.6
1	B	174	TRP	2.6
1	A	259	HIS	2.5
1	A	173	ALA	2.5
1	B	165	GLY	2.5
1	B	24	LEU	2.5
1	A	65	ILE	2.5
1	A	112	PRO	2.5
1	A	196[A]	ILE	2.5
1	A	262	ILE	2.5
1	A	415	THR	2.5
1	A	265	SER	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	150	TRP	2.5
1	B	302	TRP	2.5
1	B	47	LEU	2.5
1	A	165	GLY	2.5
1	B	59	ASN	2.4
1	B	257	SER	2.4
1	A	23	GLU	2.4
1	B	164	GLU	2.4
1	A	178	LEU	2.4
1	A	156	ILE	2.4
1	A	216	VAL	2.4
1	B	153	LEU	2.4
1	A	117	ALA	2.4
1	B	168	ALA	2.4
1	A	416	VAL	2.4
1	A	179	CYS	2.3
1	B	161	ALA	2.3
1	B	68	TYR	2.3
1	A	212	VAL	2.3
1	B	49	VAL	2.3
1	B	117	ALA	2.3
1	A	167	PHE	2.3
1	A	175	TYR	2.3
1	B	120	TRP	2.3
1	A	166	ALA	2.3
1	B	159	PHE	2.3
1	A	123	GLY	2.3
1	B	229[A]	ILE	2.3
1	A	132	VAL	2.3
1	A	410	ASN	2.2
1	B	381	LYS	2.2
1	B	163	GLY	2.2
1	B	171[A]	ARG	2.2
1	B	219	TYR	2.2
1	A	82	VAL	2.2
1	A	128	VAL	2.2
1	A	174	TRP	2.2
1	A	263	GLU	2.2
1	A	24	LEU	2.2
1	A	71	PHE	2.2
1	B	308[A]	ASN	2.2
1	B	304	PRO	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	183	VAL	2.1
1	A	95	LEU	2.1
1	B	23	GLU	2.1
1	A	146	ILE	2.1
1	A	190	TYR	2.1
1	A	97	ALA	2.1
1	B	438	LEU	2.1
1	A	101	THR	2.1
1	A	50	PHE	2.1
1	B	148	ILE	2.1
1	A	163	GLY	2.0
1	B	102	GLY	2.0
1	A	111	ASN	2.0
1	A	161	ALA	2.0
1	A	409	VAL	2.0
1	B	114	CYS	2.0
1	A	184	THR	2.0
1	A	384[A]	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	A	468	4/4	0.83	0.17	42,42,43,44	0
3	EDO	A	470	4/4	0.86	0.17	34,38,38,38	0
3	EDO	B	464	4/4	0.86	0.18	26,30,30,35	0
3	EDO	B	466	4/4	0.90	0.16	17,22,23,25	0
3	EDO	B	469	4/4	0.90	0.19	23,32,32,33	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	UNL	B	461	9/-	0.91	0.13	20,28,43,47	0
3	EDO	A	463	4/4	0.95	0.18	18,25,25,26	0
2	EPE	A	462[A]	15/15	0.96	0.13	11,20,23,24	15
3	EDO	B	467	4/4	0.96	0.14	14,14,14,15	0
3	EDO	B	465	4/4	0.97	0.12	14,14,17,17	0
4	NA	A	471	1/1	0.98	0.18	12,12,12,12	0
4	NA	B	472	1/1	0.99	0.20	17,17,17,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.