



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 02:19 PM UTC

PDB ID : 3GZC / pdb_00003gzc
Title : Structure of human selenocysteine lyase
Authors : Collins, R.; Hogbom, M.; Arrowsmith, C.; Berglund, H.; Edwards, A.; Ehn, M.; Flodin, S.; Flores, A.; Graslund, S.; Hallberg, B.M.; Hammarstrom, M.; Karlberg, T.; Kotenyova, T.; Nilsson-Ehle, P.; Nordlund, P.; Nyman, T.; Ogg, D.; Persson, C.; Sagemark, J.; Stenmark, P.; Sundstrom, M.; Thorsell, A.G.; Uppenberg, J.; Van Den Berg, S.; Weigelt, J.; Holmberg-Schiavone, L.; Schuler, H.; Structural Genomics Consortium (SGC)
Deposited on : 2009-04-07
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	440	 2% 77% 12% • 8%
1	B	440	 4% 75% 13% • 9%

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6684 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Selenocysteine lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	403	Total	C	N	O	S	0	0	0
			3103	1948	558	583	14			
1	B	402	Total	C	N	O	S	0	0	0
			3094	1943	556	581	14			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	6	SER	-	expression tag	UNP Q96I15
A	7	MET	-	expression tag	UNP Q96I15
A	175	THR	ALA	SEE REMARK 999	UNP Q96I15
B	6	SER	-	expression tag	UNP Q96I15
B	7	MET	-	expression tag	UNP Q96I15
B	175	THR	ALA	SEE REMARK 999	UNP Q96I15

- Molecule 2 is (5-HYDROXY-4,6-DIMETHYLPYRIDIN-3-YL)METHYL DIHYDROGEN PHOSPHATE (CCD ID: PLR) (formula: C₈H₁₂NO₅P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	269	Total	O	0	0
			269	269		
3	B	188	Total	O	0	0
			188	188		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Selenocysteine lyase



• Molecule 1: Selenocysteine lyase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.22Å 85.81Å 188.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.73 – 2.10 19.73 – 2.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.73-2.10) 99.8 (19.73-2.10)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.86 (at 2.09Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.179 , 0.235 0.184 , 0.239	Depositor DCC
R_{free} test set	2844 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	29.4	Xtriage
Anisotropy	0.116	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 45.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6684	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PLR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.14	2/3164 (0.1%)	1.09	6/4295 (0.1%)
1	B	1.02	1/3155 (0.0%)	1.05	3/4283 (0.1%)
All	All	1.08	3/6319 (0.0%)	1.07	9/8578 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	103	ASN	C-O	-5.92	1.17	1.24
1	B	156	VAL	CA-CB	5.27	1.61	1.54
1	A	73	ASP	C-O	-5.04	1.18	1.24

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	324	ARG	NE-CZ-NH2	6.15	124.74	119.20
1	A	166	VAL	CA-C-N	-5.91	114.53	120.21
1	A	166	VAL	C-N-CA	-5.91	114.53	120.21
1	B	367	ARG	CG-CD-NE	5.64	124.40	112.00
1	B	348	GLN	N-CA-C	5.23	116.83	108.41
1	A	324	ARG	NE-CZ-NH1	-5.21	116.29	121.50
1	B	383	SER	N-CA-C	5.13	116.79	109.14
1	A	43	GLU	CA-C-N	5.05	126.15	119.84
1	A	43	GLU	C-N-CA	5.05	126.15	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3103	0	3099	45	0
1	B	3094	0	3090	50	0
2	A	15	0	7	0	0
2	B	15	0	8	0	0
3	A	269	0	0	15	0
3	B	188	0	0	8	0
All	All	6684	0	6204	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (91) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:50:MET:HE1	1:A:301:MET:CE	1.77	1.15
1:A:50:MET:CE	1:A:301:MET:HE2	1.78	1.13
1:B:50:MET:CE	1:B:301:MET:HE2	1.92	1.00
1:A:50:MET:HE1	1:A:301:MET:HE2	1.35	0.93
1:B:50:MET:HE2	1:B:301:MET:HE2	1.55	0.87
1:A:301:MET:HE3	1:B:301:MET:HE3	1.56	0.87
1:A:50:MET:HE1	1:A:301:MET:HE1	1.55	0.87
1:A:50:MET:HE3	1:A:301:MET:HE2	1.60	0.82
1:B:290:ARG:HH11	1:B:290:ARG:HG2	1.45	0.82
1:B:50:MET:HE1	1:B:301:MET:CE	2.14	0.78
1:B:50:MET:HE1	1:B:301:MET:HE2	1.65	0.78
1:B:415:ARG:HD2	3:B:809:HOH:O	1.85	0.75
1:B:367:ARG:HH11	1:B:367:ARG:HG3	1.55	0.72
1:B:240:LYS:HA	1:B:323:MET:HE1	1.72	0.69
1:B:47:ILE:O	1:B:51:THR:HG23	1.93	0.69
1:B:169:SER:OG	1:B:171:VAL:HG13	1.96	0.66
1:B:50:MET:CE	1:B:301:MET:CE	2.70	0.65
1:A:74:ILE:HD13	3:A:605:HOH:O	1.96	0.64
1:B:337:GLU:HG3	1:B:431:VAL:HG11	1.80	0.62
1:A:319:TYR:HB3	1:A:323:MET:CE	2.31	0.60
1:A:344:HIS:NE2	1:A:364:ARG:HD2	2.17	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:50:MET:CE	1:A:301:MET:CE	2.51	0.59
1:A:115:HIS:HE1	3:A:639:HOH:O	1.87	0.58
1:B:348:GLN:HE21	1:B:354:ARG:NH2	2.03	0.56
1:B:197:ASN:HD22	1:B:200:THR:H	1.53	0.56
1:B:50:MET:HE3	1:B:301:MET:HA	1.88	0.55
1:B:242:ARG:HD3	3:B:786:HOH:O	2.06	0.55
1:A:70:LYS:HE2	3:A:842:HOH:O	2.05	0.55
1:A:319:TYR:C	1:A:323:MET:HE3	2.32	0.55
1:A:366:PRO:O	1:A:369:GLN:NE2	2.39	0.54
1:B:202:ILE:HD11	1:B:347:SER:O	2.07	0.53
1:A:91:GLN:HB2	3:A:457:HOH:O	2.09	0.53
1:A:242:ARG:NE	3:A:744:HOH:O	2.43	0.52
1:B:112:LYS:NZ	3:B:832:HOH:O	2.43	0.51
1:B:369:GLN:HG2	1:B:394:ASP:OD2	2.11	0.51
1:B:299:THR:CG2	3:B:3:HOH:O	2.59	0.51
1:A:153:GLU:O	1:A:156:VAL:HG12	2.11	0.51
1:B:89:LYS:NZ	3:B:572:HOH:O	2.44	0.50
1:B:320:GLU:HA	1:B:323:MET:HE3	1.94	0.49
1:A:319:TYR:O	1:A:323:MET:HE3	2.12	0.49
1:B:78:ALA:O	1:B:82:LEU:HD23	2.12	0.49
1:A:347:SER:HB2	3:A:628:HOH:O	2.13	0.48
1:B:299:THR:HG23	3:B:3:HOH:O	2.12	0.48
1:A:415:ARG:HD2	3:A:801:HOH:O	2.13	0.48
1:A:30:ARG:HD3	3:A:767:HOH:O	2.13	0.47
1:B:434:LEU:O	1:B:438:VAL:HG13	2.14	0.47
1:B:108:HIS:CD2	1:B:282:MET:HG2	2.50	0.47
1:B:396:PRO:HG2	1:B:401:LEU:HD11	1.97	0.47
1:A:216:LEU:HD22	1:A:220:ARG:HG3	1.97	0.47
1:B:337:GLU:HG3	1:B:431:VAL:CG1	2.43	0.47
1:A:43:GLU:HG2	1:A:261:TYR:HB3	1.97	0.46
1:B:176:GLU:HG3	3:B:621:HOH:O	2.14	0.46
1:A:54:MET:HE1	1:B:54:MET:CE	2.46	0.46
1:A:186:PRO:O	1:A:189:ARG:NH2	2.48	0.46
1:B:322:HIS:O	1:B:326:VAL:HG13	2.14	0.46
1:A:91:GLN:CB	3:A:457:HOH:O	2.63	0.46
1:B:319:TYR:HB3	1:B:323:MET:HE2	1.97	0.46
1:A:112:LYS:HA	1:A:112:LYS:HD2	1.88	0.45
1:B:118:GLN:O	1:B:119:THR:OG1	2.27	0.45
1:B:391:ASP:OD2	1:B:391:ASP:N	2.50	0.45
1:A:185:ARG:O	1:A:220:ARG:NH2	2.49	0.45
1:A:273:LEU:HD23	1:A:290:ARG:NH2	2.32	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:65:TYR:O	1:A:69:ARG:HG2	2.17	0.45
1:B:383:SER:HB2	3:B:599:HOH:O	2.17	0.44
1:A:169:SER:OG	1:A:171:VAL:HG12	2.17	0.44
1:B:99:GLY:HA3	1:B:256:VAL:HG22	1.98	0.44
1:A:432:GLN:NE2	1:A:435:LYS:HD3	2.31	0.44
1:B:329:TYR:HA	1:B:332:GLU:HG2	1.99	0.44
1:A:258:HIS:HD2	3:A:453:HOH:O	2.00	0.44
1:A:156:VAL:HG11	3:A:586:HOH:O	2.17	0.44
1:A:290:ARG:HG2	3:A:758:HOH:O	2.18	0.44
1:A:54:MET:CE	1:B:54:MET:HE1	2.49	0.43
1:A:342:ARG:NH2	3:A:754:HOH:O	2.51	0.43
1:B:185:ARG:O	1:B:220:ARG:NH2	2.51	0.43
1:B:108:HIS:CG	1:B:282:MET:HG2	2.54	0.43
1:B:338:PHE:HB2	1:B:343:ILE:HD11	1.99	0.43
1:A:434:LEU:O	1:A:438:VAL:HG13	2.19	0.42
1:B:50:MET:CE	1:B:301:MET:HA	2.49	0.42
1:B:275:GLU:HG3	1:B:276:PHE:CD1	2.55	0.42
1:A:236:GLN:O	1:A:240:LYS:HD2	2.19	0.42
1:A:376:GLN:HG2	3:A:601:HOH:O	2.19	0.41
1:B:206:VAL:N	1:B:207:PRO:CD	2.83	0.41
1:B:89:LYS:HB3	1:B:91:GLN:OE1	2.20	0.41
1:B:290:ARG:HH11	1:B:290:ARG:CG	2.22	0.41
1:B:91:GLN:H	1:B:91:GLN:CD	2.28	0.41
1:B:105:LEU:HD13	1:B:282:MET:HG3	2.02	0.41
1:A:31:LYS:HB3	1:A:381:MET:HE2	2.01	0.41
1:A:65:TYR:CZ	1:B:381:MET:HA	2.56	0.41
1:A:326:VAL:CG2	1:A:418:VAL:HG21	2.51	0.41
1:A:433:ASP:O	1:A:436:GLN:HG3	2.21	0.40
1:A:177:VAL:HG13	3:A:464:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	399/440 (91%)	390 (98%)	9 (2%)	0	100	100
1	B	398/440 (90%)	388 (98%)	9 (2%)	1 (0%)	36	36
All	All	797/880 (91%)	778 (98%)	18 (2%)	1 (0%)	48	50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	65	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	334/360 (93%)	317 (95%)	17 (5%)	21	21
1	B	333/360 (92%)	314 (94%)	19 (6%)	18	17
All	All	667/720 (93%)	631 (95%)	36 (5%)	20	18

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48	GLN
1	A	82	LEU
1	A	89	LYS
1	A	112	LYS
1	A	119	THR
1	A	176	GLU
1	A	177	VAL
1	A	185	ARG
1	A	199	GLU
1	A	216	LEU
1	A	225	LEU
1	A	283	LEU
1	A	290	ARG
1	A	307	LYS
1	A	326	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	340	GLN
1	A	432	GLN
1	B	48	GLN
1	B	69	ARG
1	B	73	ASP
1	B	89	LYS
1	B	118	GLN
1	B	156	VAL
1	B	171	VAL
1	B	185	ARG
1	B	211	GLN
1	B	216	LEU
1	B	225	LEU
1	B	246	GLU
1	B	283	LEU
1	B	290	ARG
1	B	299	THR
1	B	341	LYS
1	B	342	ARG
1	B	367	ARG
1	B	391	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	108	HIS
1	A	218	GLN
1	A	241	GLN
1	A	432	GLN
1	B	113	HIS
1	B	117	ASN
1	B	197	ASN
1	B	211	GLN
1	B	344	HIS
1	B	348	GLN
1	B	392	HIS
1	B	440	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PLR	A	500	1	15,15,15	0.78	1 (6%)	21,22,22	1.66	5 (23%)
2	PLR	B	500	1	15,15,15	1.03	0	21,22,22	2.29	8 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLR	A	500	1	-	0/6/6/6	0/1/1/1
2	PLR	B	500	1	-	1/6/6/6	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	PLR	C6-N1	-2.05	1.30	1.34

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	500	PLR	C4A-C4-C5	6.38	127.51	120.94

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	500	PLR	C4A-C4-C3	-3.98	113.89	120.52
2	A	500	PLR	C2A-C2-C3	3.64	125.06	120.80
2	B	500	PLR	C6-N1-C2	3.13	124.87	119.20
2	A	500	PLR	O3P-P-O4P	-2.68	99.69	106.67
2	B	500	PLR	C5A-C5-C6	-2.67	115.01	119.36
2	A	500	PLR	C6-N1-C2	2.62	123.95	119.20
2	B	500	PLR	C5-C6-N1	-2.62	119.58	123.83
2	A	500	PLR	C3-C2-N1	-2.51	117.80	120.96
2	B	500	PLR	C3-C2-N1	-2.31	118.05	120.96
2	A	500	PLR	O4P-C5A-C5	2.21	113.50	109.36
2	B	500	PLR	O4P-C5A-C5	2.21	113.49	109.36
2	B	500	PLR	O3-C3-C2	2.14	122.01	117.58

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	500	PLR	C4-C5-C5A-O4P

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	403/440 (91%)	-0.02	7 (1%) 69 71	25, 31, 44, 58	0
1	B	402/440 (91%)	0.25	16 (3%) 42 44	23, 32, 44, 61	0
All	All	805/880 (91%)	0.12	23 (2%) 53 56	23, 31, 44, 61	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	119	THR	4.1
1	B	347	SER	3.7
1	B	394	ASP	3.6
1	B	133	GLY	3.5
1	B	393	GLY	3.4
1	B	408	ASP	3.2
1	B	73	ASP	3.1
1	B	318	ALA	3.0
1	A	393	GLY	2.9
1	B	69	ARG	2.6
1	B	443	ASP	2.5
1	A	443	ASP	2.4
1	B	313	THR	2.4
1	A	390	SER	2.4
1	A	276	PHE	2.4
1	B	311	LEU	2.3
1	A	29	GLU	2.2
1	B	389	HIS	2.1
1	A	394	ASP	2.1
1	B	64	PRO	2.1
1	B	310	GLU	2.0
1	B	118	GLN	2.0
1	B	391	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PLR	A	500	15/15	0.95	0.08	22,23,26,29	0
2	PLR	B	500	15/15	0.97	0.08	23,31,33,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.