



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 04:39 PM UTC

PDB ID : 4GZR / pdb_00004gzs
Title : Crystal structure of the Mycobacterium tuberculosis H37Rv EsxOP (Rv2346c-Rv2347c) complex in space group C2221
Authors : Arbing, M.A.; Chan, S.; He, Q.; Harris, L.; Zhou, T.T.; Kuo, E.; Ahn, C.J.; Eisenberg, D.; TB Structural Genomics Consortium (TBSGC); Integrated Center for Structure and Function Innovation (ISFI)
Deposited on : 2012-09-06
Resolution : 2.55 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1853 (2.58-2.54)
Clashscore	190562	1897 (2.58-2.54)
Ramachandran outliers	187476	1875 (2.58-2.54)
Sidechain outliers	187428	1875 (2.58-2.54)
RSRZ outliers	180081	1853 (2.58-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	101	55% 43%
1	C	101	54% 6% 40%
2	B	99	2% 73% 22%
2	D	99	3% 75% 6% 19%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4189 atoms, of which 1982 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ESAT-6-like protein 6.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	58	Total	C	H	N	O	S	0	0	0
			840	273	404	80	81	2			
1	C	61	Total	C	H	N	O	S	0	0	0
			870	286	411	83	88	2			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	95	THR	-	expression tag	UNP P95242
A	96	HIS	-	expression tag	UNP P95242
A	97	HIS	-	expression tag	UNP P95242
A	98	HIS	-	expression tag	UNP P95242
A	99	HIS	-	expression tag	UNP P95242
A	100	HIS	-	expression tag	UNP P95242
A	101	HIS	-	expression tag	UNP P95242
C	95	THR	-	expression tag	UNP P95242
C	96	HIS	-	expression tag	UNP P95242
C	97	HIS	-	expression tag	UNP P95242
C	98	HIS	-	expression tag	UNP P95242
C	99	HIS	-	expression tag	UNP P95242
C	100	HIS	-	expression tag	UNP P95242
C	101	HIS	-	expression tag	UNP P95242


- Molecule 2 is a protein called ESAT-6-like protein 7.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	77	Total	C	H	N	O	S	0	0	0
			1200	376	576	122	119	7			
2	D	80	Total	C	H	N	O	S	0	1	0
			1243	394	591	126	125	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	SER	-	expression tag	UNP P95243
D	0	SER	-	expression tag	UNP P95243

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	6	Total	O	0	0
			6	6		
4	B	6	Total	O	0	0
			6	6		
4	C	9	Total	O	0	0
			9	9		
4	D	10	Total	O	0	0
			10	10		


- Molecule 1: ESAT-6-like protein 6





- Chain C:  54% 6% 40%



- Chain B: 2% 73% .. 22%



- Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	76.10Å 112.07Å 95.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.05 – 2.55 38.05 – 2.55	Depositor EDS
% Data completeness (in resolution range)	98.0 (38.05-2.55) 98.1 (38.05-2.55)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.03 (at 2.55Å)	Xtriage
Refinement program	PHENIX 1.8_1069	Depositor
R, R_{free}	0.199 , 0.233 (Not available) , 0.263	Depositor DCC
R_{free} test set	1334 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	65.3	Xtriage
Anisotropy	0.484	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 27.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4189	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.88	0/443	0.86	0/598
1	C	1.02	0/466	0.94	0/630
2	B	0.82	0/632	0.88	0/847
2	D	0.83	0/665	0.86	0/894
All	All	0.88	0/2206	0.89	0/2969

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	436	404	416	0	0
1	C	459	411	433	3	0
2	B	624	576	585	4	0
2	D	652	591	608	8	0
3	D	5	0	0	0	0
4	A	6	0	0	0	0
4	B	6	0	0	0	0
4	C	9	0	0	0	0
4	D	10	0	0	0	0
All	All	2207	1982	2042	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 14 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:54:LEU:HD12	2:D:54:LEU:H	1.53	0.74
2:D:34[B]:TRP:CZ3	2:D:57:MET:HB3	2.42	0.54
2:D:89:GLU:HA	2:D:89:GLU:OE1	2.08	0.53
2:D:54:LEU:HD12	2:D:54:LEU:N	2.22	0.53
2:B:94:GLN:O	2:B:95:ILE:HG22	2.10	0.52

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	56/101 (55%)	56 (100%)	0	0	100	100
1	C	59/101 (58%)	58 (98%)	1 (2%)	0	100	100
2	B	73/99 (74%)	71 (97%)	2 (3%)	0	100	100
2	D	77/99 (78%)	77 (100%)	0	0	100	100
All	All	265/400 (66%)	262 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	40/73 (55%)	38 (95%)	2 (5%)	22	33
1	C	43/73 (59%)	42 (98%)	1 (2%)	44	60
2	B	64/79 (81%)	62 (97%)	2 (3%)	35	50
2	D	67/79 (85%)	67 (100%)	0	100	100
All	All	214/304 (70%)	209 (98%)	5 (2%)	44	60

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48	VAL
1	A	54	ILE
2	B	90	GLN
2	B	95	ILE
1	C	11	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	56	GLN
1	A	62	GLN
1	C	60	ASN
2	D	10	HIS
2	D	66	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	D	101	-	4,4,4	0.26	0	6,6,6	0.10	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	58/101 (57%)	-0.16	1 (1%) 69 70	21, 43, 66, 76	0
1	C	61/101 (60%)	-0.18	0 100 100	20, 30, 75, 87	0
2	B	77/99 (77%)	-0.13	2 (2%) 57 59	22, 43, 89, 104	0
2	D	80/99 (80%)	-0.19	3 (3%) 44 45	21, 37, 84, 98	1 (1%)
All	All	276/400 (69%)	-0.16	6 (2%) 62 63	20, 38, 82, 104	1 (0%)

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	51	ALA	5.0
2	B	95	ILE	3.5
1	A	48	VAL	2.4
2	D	54	LEU	2.1
2	B	54	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	D	101	5/5	0.64	0.09	107,116,117,121	0

6.5 Other polymers [i](#)

There are no such residues in this entry.