



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 10:44 AM UTC

PDB ID : 6GZY / pdb_00006gzy
Title : HOIP-fragment5 complex
Authors : Johansson, H.; Tsai, Y.C.I.; Fantom, K.; Chung, C.W.; Martino, L.; House, D.; Rittinger, K.
Deposited on : 2018-07-05
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	223	12% 84% 5% 11%
1	B	223	14% 84% • 13%

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 6412 atoms, of which 2903 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called E3 ubiquitin-protein ligase RNF31.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	199	Total	C	H	N	O	S	49	6	0
			3077	1014	1455	298	293	17			
1	B	194	Total	C	H	N	O	S	28	3	0
			2986	986	1417	286	280	17			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	850	GLY	-	expression tag	UNP Q96EP0
A	851	PRO	-	expression tag	UNP Q96EP0
A	852	GLY	-	expression tag	UNP Q96EP0
B	850	GLY	-	expression tag	UNP Q96EP0
B	851	PRO	-	expression tag	UNP Q96EP0
B	852	GLY	-	expression tag	UNP Q96EP0

- Molecule 2 is methyl 4-[(2-oxidanylidene-1,5,6,7-tetrahydrocyclopenta[b]pyridin-3-yl)carbon ylamino]butanoate (CCD ID: FHH) (formula: C₁₄H₁₈N₂O₄).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	1	0
			20	14	2	4		
2	B	1	Total	C	N	O	0	0
			20	14	2	4		

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Zn	0	0
			4	4		
3	B	4	Total	Zn	0	0
			4	4		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	0	0
			7	2	3	2		
5	A	1	Total	C	H	O	0	0
			10	2	6	2		
5	B	1	Total	C	H	O	0	0
			7	2	3	2		
5	B	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 6 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	H	O	0	0
			23	6	13	4		

- Molecule 7 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Na	0	0
			1	1		
7	B	1	Total	Na	0	0
			1	1		

- Molecule 8 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	116	Total	O	0	0
			116	116		
8	B	91	Total	O	0	0
			91	91		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: E3 ubiquitin-protein ligase RNF31

Chain A: 



- Molecule 1: E3 ubiquitin-protein ligase RNF31

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	111.20Å 117.25Å 37.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	80.69 – 2.15 80.69 – 2.15	Depositor EDS
% Data completeness (in resolution range)	79.0 (80.69-2.15) 78.9 (80.69-2.15)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.95 (at 2.14Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.199 , 0.246 0.199 , 0.245	Depositor DCC
R_{free} test set	1307 reflections (2.79%)	wwPDB-VP
Wilson B-factor (Å ²)	25.1	Xtriage
Anisotropy	0.092	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 59.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6412	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 43.52 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.7435e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, NA, PGE, FHH, EDO, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.12	0/1691	0.32	0/2282
1	B	0.12	0/1629	0.34	0/2196
All	All	0.12	0/3320	0.33	0/4478

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1622	1455	1543	7	0
1	B	1569	1417	1507	4	0
2	A	20	0	0	0	0
2	B	20	0	0	0	0
3	A	4	0	0	0	0
3	B	4	0	0	0	0
4	A	20	0	0	0	0
4	B	15	0	0	0	0
5	A	8	9	12	0	0
5	B	8	9	12	0	0
6	A	10	13	14	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	1	0	0	0	0
7	B	1	0	0	0	0
8	A	116	0	0	0	0
8	B	91	0	0	2	0
All	All	3509	2903	3088	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:867:ASN:HB2	1:A:1070:ARG:NH2	2.12	0.64
1:A:865:GLN:HE22	1:A:972:ILE:HD12	1.67	0.58
1:B:1032:ARG:NH1	8:B:4101:HOH:O	2.43	0.51
1:A:866:GLU:HA	8:B:4181:HOH:O	2.12	0.49
1:A:867:ASN:HB2	1:A:1070:ARG:HH21	1.78	0.48
1:A:886[A]:MET:HA	1:A:886[A]:MET:HE2	1.98	0.46
1:A:913:GLU:HB3	1:A:916:CYS:HB2	2.00	0.43
1:A:940:LEU:C	1:A:940:LEU:HD23	2.44	0.43
1:B:886[B]:MET:HE1	1:B:931:LEU:HD23	2.03	0.41
1:B:919:LYS:HE3	1:B:919:LYS:HB3	1.88	0.41
1:B:908:LYS:HB2	1:B:920:LYS:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	199/223 (89%)	193 (97%)	6 (3%)	0	100	100
1	B	189/223 (85%)	182 (96%)	7 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	388/446 (87%)	375 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	177/185 (96%)	173 (98%)	4 (2%)	44	49
1	B	171/185 (92%)	170 (99%)	1 (1%)	78	84
All	All	348/370 (94%)	343 (99%)	5 (1%)	57	66

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	866	GLU
1	A	999	GLN
1	A	1036	VAL
1	A	1058	THR
1	B	1058	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	947	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 10 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	A	4007	-	4,4,4	0.24	0	6,6,6	0.10	0
4	SO4	A	4008	-	4,4,4	0.23	0	6,6,6	0.16	0
4	SO4	B	4007	-	4,4,4	0.23	0	6,6,6	0.11	0
5	EDO	A	4011	-	3,3,3	0.37	0	2,2,2	0.45	0
5	EDO	B	4010	-	3,3,3	0.39	0	2,2,2	0.41	0
6	PGE	A	4012	-	9,9,9	0.52	0	8,8,8	0.30	0
5	EDO	A	4010	-	3,3,3	0.44	0	2,2,2	0.36	0
4	SO4	A	4009	-	4,4,4	0.25	0	6,6,6	0.07	0
5	EDO	B	4009	-	3,3,3	0.44	0	2,2,2	0.34	0
4	SO4	B	4006	-	4,4,4	0.23	0	6,6,6	0.07	0
4	SO4	B	4008	-	4,4,4	0.24	0	6,6,6	0.11	0
4	SO4	A	4006	-	4,4,4	0.23	0	6,6,6	0.10	0
2	FHH	B	4001	1	21,21,21	0.57	0	25,28,28	0.96	1 (4%)
2	FHH	A	4001	1	21,21,21	0.54	0	25,28,28	0.94	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	4011	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	B	4010	-	-	1/1/1/1	-
6	PGE	A	4012	-	-	2/7/7/7	-
5	EDO	A	4010	-	-	0/1/1/1	-
5	EDO	B	4009	-	-	1/1/1/1	-
2	FHH	B	4001	1	-	7/13/19/19	0/2/2/2
2	FHH	A	4001	1	-	8/13/19/19	0/2/2/2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	4001	FHH	C-C7-C12	-2.00	121.04	122.64

There are no chirality outliers.

All (19) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	4001	FHH	O-C-C7-C12
2	A	4001	FHH	NA-C-C7-C12
2	A	4001	FHH	C1-C2-C3-C4
2	B	4001	FHH	C1-C2-C3-C4
2	A	4001	FHH	C3-C4-O4-C6
2	B	4001	FHH	C3-C4-O4-C6
2	B	4001	FHH	NA-C1-C2-C3
2	A	4001	FHH	NA-C1-C2-C3
2	A	4001	FHH	O2-C4-O4-C6
6	A	4012	PGE	O1-C1-C2-O2
2	B	4001	FHH	O2-C4-O4-C6
5	B	4009	EDO	O1-C1-C2-O2
5	B	4010	EDO	O1-C1-C2-O2
2	B	4001	FHH	O-C-C7-C12
6	A	4012	PGE	O2-C3-C4-O3
2	A	4001	FHH	O-C-C7-C8
2	B	4001	FHH	O-C-C7-C8
2	B	4001	FHH	C2-C3-C4-O4
2	A	4001	FHH	C2-C3-C4-O4

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	199/223 (89%)	0.59	27 (13%) 7 7	11, 34, 72, 101	14 (7%)
1	B	194/223 (86%)	0.69	31 (15%) 5 5	10, 34, 79, 100	8 (4%)
All	All	393/446 (88%)	0.64	58 (14%) 5 6	10, 34, 77, 101	22 (5%)

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1047	PRO	8.0
1	B	918	VAL	6.4
1	B	1046	PRO	6.3
1	A	867	ASN	6.1
1	A	982	ARG	6.1
1	B	862	MET	4.7
1	A	918	VAL	4.7
1	B	968	GLY	4.3
1	B	867	ASN	4.3
1	A	915	ASN	4.2
1	A	920	LYS	4.0
1	A	968	GLY	3.8
1	B	958	PRO	3.8
1	B	974	GLN	3.7
1	B	1041	LEU	3.6
1	B	917	ARG	3.5
1	B	859	GLY	3.5
1	B	982	ARG	3.4
1	A	974	GLN	3.4
1	B	1037	ARG	3.2
1	A	862	MET	3.2
1	A	983	ASP	3.1
1	A	919	LYS	3.1
1	A	859	GLY	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	987	GLY	3.0
1	B	1048	ALA	2.9
1	B	921	SER	2.9
1	B	914	PRO	2.8
1	B	957	PRO	2.8
1	A	858	GLN	2.8
1	B	970	ARG	2.8
1	B	1040	PRO	2.8
1	B	919	LYS	2.8
1	B	909	ASN	2.8
1	A	910	LYS	2.8
1	A	865	GLN	2.7
1	A	866	GLU	2.7
1	A	868	GLY	2.7
1	A	992	ALA	2.6
1	A	914	PRO	2.6
1	B	990	THR	2.5
1	A	989	GLU	2.5
1	B	956	GLU	2.4
1	A	913	GLU	2.4
1	B	983	ASP	2.4
1	B	986	CYS	2.3
1	A	1039	GLN	2.3
1	B	997	LEU	2.3
1	A	987	GLY	2.3
1	B	915	ASN	2.2
1	B	868	GLY	2.2
1	B	920	LYS	2.2
1	A	956	GLU	2.1
1	A	861	ALA	2.1
1	A	909	ASN	2.1
1	B	992	ALA	2.1
1	A	917	ARG	2.0
1	A	1045	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	SO4	A	4009	5/5	0.76	0.16	83,84,88,92	0
6	PGE	A	4012	10/10	0.76	0.26	69,83,89,89	0
5	EDO	B	4010	4/4	0.80	0.25	31,38,45,52	0
4	SO4	A	4008	5/5	0.82	0.18	69,71,75,80	0
5	EDO	A	4011	4/4	0.83	0.18	37,44,53,59	0
5	EDO	B	4009	4/4	0.86	0.18	38,45,54,63	0
5	EDO	A	4010	4/4	0.88	0.15	36,43,51,59	0
2	FHH	B	4001	20/20	0.89	0.12	18,24,33,42	2
4	SO4	B	4008	5/5	0.91	0.13	54,62,66,67	0
2	FHH	A	4001	20/20	0.91	0.12	22,25,36,36	4
7	NA	B	4011	1/1	0.93	0.15	44,44,44,44	0
7	NA	A	4013	1/1	0.94	0.18	48,48,48,48	0
4	SO4	A	4007	5/5	0.96	0.07	36,44,52,53	0
4	SO4	B	4007	5/5	0.96	0.07	35,36,45,51	0
4	SO4	B	4006	5/5	0.98	0.06	29,31,34,38	0
4	SO4	A	4006	5/5	0.98	0.06	28,30,31,39	0
3	ZN	A	4004	1/1	0.99	0.02	35,35,35,35	0
3	ZN	B	4003	1/1	0.99	0.02	14,14,14,14	0
3	ZN	B	4004	1/1	0.99	0.03	35,35,35,35	0
3	ZN	B	4005	1/1	0.99	0.04	40,40,40,40	0
3	ZN	A	4002	1/1	0.99	0.02	17,17,17,17	0
3	ZN	A	4003	1/1	0.99	0.02	16,16,16,16	0
3	ZN	B	4002	1/1	1.00	0.03	22,22,22,22	0
3	ZN	A	4005	1/1	1.00	0.03	36,36,36,36	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FHH B 4001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FHH A 4001:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.