



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 15, 2026 – 01:43 AM UTC

PDB ID : 3H0M / pdb_00003h0m
Title : Structure of trna-dependent amidotransferase gatcab from aquifex aeolicus
Authors : Wu, J.; Bu, W.; Sheppard, K.; Kitabatake, M.; Soll, D.; Smith, J.L.
Deposited on : 2009-04-09
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	478	 71% 25% .
1	D	478	 71% 26% .
1	G	478	 75% 23% .
1	J	478	 72% 24% .
1	M	478	 75% 22% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	P	478	
1	S	478	
1	V	478	
2	B	478	
2	E	478	
2	H	478	
2	K	478	
2	N	478	
2	Q	478	
2	T	478	
2	W	478	
3	C	94	
3	F	94	
3	I	94	
3	L	94	
3	O	94	
3	R	94	
3	U	94	
3	X	94	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GLN	A	901	-	-	X	-
4	GLN	G	903	-	-	X	-
4	GLN	J	904	-	-	X	-
4	GLN	M	905	-	-	X	-
4	GLN	P	906	-	-	X	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GLN	S	907	-	-	X	-
4	GLN	V	908	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 62935 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamyl-tRNA(Gln) amidotransferase subunit A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	478	Total	C	N	O	S	0	0	0
			3784	2450	615	712	7			
1	D	478	Total	C	N	O	S	0	0	0
			3784	2450	615	712	7			
1	G	478	Total	C	N	O	S	0	0	0
			3784	2450	615	712	7			
1	J	478	Total	C	N	O	S	0	0	0
			3784	2450	615	712	7			
1	M	478	Total	C	N	O	S	0	0	0
			3784	2450	615	712	7			
1	P	478	Total	C	N	O	S	0	0	0
			3784	2450	615	712	7			
1	S	478	Total	C	N	O	S	0	0	0
			3784	2450	615	712	7			
1	V	478	Total	C	N	O	S	0	0	0
			3784	2450	615	712	7			

- Molecule 2 is a protein called Aspartyl/glutamyl-tRNA(Asn/Gln) amidotransferase subunit B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	410	Total	C	N	O	S	0	0	0
			3308	2104	567	622	15			
2	E	410	Total	C	N	O	S	0	0	0
			3308	2104	567	622	15			
2	H	410	Total	C	N	O	S	0	0	0
			3308	2104	567	622	15			
2	K	410	Total	C	N	O	S	0	0	0
			3308	2104	567	622	15			
2	N	410	Total	C	N	O	S	0	0	0
			3308	2104	567	622	15			
2	Q	410	Total	C	N	O	S	0	0	0
			3308	2104	567	622	15			

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	T	410	Total	C	N	O	S	0	0	0
			3308	2104	567	622	15			
2	W	410	Total	C	N	O	S	0	0	0
			3308	2104	567	622	15			

- Molecule 3 is a protein called Glutamyl-tRNA(Gln) amidotransferase subunit C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	91	Total	C	N	O	S	0	0	0
			764	487	125	150	2			
3	F	91	Total	C	N	O	S	0	0	0
			764	487	125	150	2			
3	I	91	Total	C	N	O	S	0	0	0
			764	487	125	150	2			
3	L	91	Total	C	N	O	S	0	0	0
			764	487	125	150	2			
3	O	91	Total	C	N	O	S	0	0	0
			764	487	125	150	2			
3	R	91	Total	C	N	O	S	0	0	0
			764	487	125	150	2			
3	U	91	Total	C	N	O	S	0	0	0
			764	487	125	150	2			
3	X	91	Total	C	N	O	S	0	0	0
			764	487	125	150	2			

- Molecule 4 is GLUTAMINE (CCD ID: GLN) (formula: C₅H₁₀N₂O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O 9 5 1 3	0	0
4	D	1	Total C N O 9 5 1 3	0	0
4	G	1	Total C N O 9 5 1 3	0	0
4	J	1	Total C N O 9 5 1 3	0	0
4	M	1	Total C N O 9 5 1 3	0	0
4	P	1	Total C N O 9 5 1 3	0	0
4	S	1	Total C N O 9 5 1 3	0	0
4	V	1	Total C N O 9 5 1 3	0	0

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Mg 1 1	0	0
5	H	1	Total Mg 1 1	0	0
5	K	1	Total Mg 1 1	0	0
5	N	1	Total Mg 1 1	0	0
5	Q	1	Total Mg 1 1	0	0
5	T	1	Total Mg 1 1	0	0
5	W	1	Total Mg 1 1	0	0

- Molecule 6 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total Zn 1 1	0	0
6	E	1	Total Zn 1 1	0	0
6	H	1	Total Zn 1 1	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	K	1	Total 1	Zn 1	0	0
6	N	1	Total 1	Zn 1	0	0
6	Q	1	Total 1	Zn 1	0	0
6	T	1	Total 1	Zn 1	0	0
6	W	1	Total 1	Zn 1	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glutamyl-tRNA(Gln) amidotransferase subunit A




• Molecule 1: Glutamyl-tRNA(Gln) amidotransferase subunit A




• Molecule 1: Glutamyl-tRNA(Gln) amidotransferase subunit A






• Molecule 1: Glutamyl-tRNA(Gln) amidotransferase subunit A

Chain J:  72% 24%




• Molecule 1: Glutamyl-tRNA(Gln) amidotransferase subunit A

Chain M:  75% 22%




• Molecule 1: Glutamyl-tRNA(Gln) amidotransferase subunit A

Chain P:  70% 26%




• Molecule 1: Glutamyl-tRNA(Gln) amidotransferase subunit A

Chain S:  71% 25%




• Molecule 1: Glutamyl-tRNA(Gln) amidotransferase subunit A

Chain V:  71% 25%




• Molecule 2: Aspartyl/glutamyl-tRNA(Asn/Gln) amidotransferase subunit B

Chain B:  64% 18% 14%




• Molecule 2: Aspartyl/glutamyl-tRNA(Asn/Gln) amidotransferase subunit B

Chain E: 62% 22% 14%





• Molecule 2: Aspartyl/glutamyl-tRNA(Asn/Gln) amidotransferase subunit B

Chain H: 62% 21% 14%




Chain K: 63% 21% • 14%






• Molecule 2: Aspartyl/glutamyl-tRNA(Asn/Gln) amidotransferase subunit B

Chain T: 63% 19% 14%



• Molecule 2: Aspartyl/glutamyl-tRNA(Asn/Gln) amidotransferase subunit B

Chain W: 63% 20% 14%



• Molecule 3: Glutamyl-tRNA(Gln) amidotransferase subunit C

Chain C: 68% 28% 4%



- Molecule 3: Glutamyl-tRNA(Gln) amidotransferase subunit C



- Molecule 3: Glutamyl-tRNA(Gln) amidotransferase subunit C




- Molecule 3: Glutamyl-tRNA(Gln) amidotransferase subunit C



- Molecule 3: Glutamyl-tRNA(Gln) amidotransferase subunit C



- Molecule 3: Glutamyl-tRNA(Gln) amidotransferase subunit C



- Molecule 3: Glutamyl-tRNA(Gln) amidotransferase subunit C



- Molecule 3: Glutamyl-tRNA(Gln) amidotransferase subunit C





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	128.25Å 129.86Å 155.07Å 90.01° 89.96° 90.11°	Depositor
Resolution (Å)	39.37 – 2.80 39.37 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.0 (39.37-2.80) 97.6 (39.37-2.80)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.89 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.254 , 0.305 0.260 , 0.263	Depositor DCC
R_{free} test set	12256 reflections (2.76%)	wwPDB-VP
Wilson B-factor (Å ²)	62.9	Xtriage
Anisotropy	0.753	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 33.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.000 for -k,h,l 0.000 for k,-h,l 0.185 for h,-k,-l 0.437 for -h,k,-l 0.184 for -h,-k,l 0.000 for -k,-h,-l 0.000 for k,h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	62935	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 37.49 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.2523e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	1/3874 (0.0%)	0.98	6/5244 (0.1%)
1	D	0.74	0/3874	0.97	6/5244 (0.1%)
1	G	0.77	2/3874 (0.1%)	0.96	3/5244 (0.1%)
1	J	0.75	0/3874	0.99	7/5244 (0.1%)
1	M	0.74	0/3874	0.95	5/5244 (0.1%)
1	P	0.73	0/3874	0.96	4/5244 (0.1%)
1	S	0.75	0/3874	0.98	4/5244 (0.1%)
1	V	0.76	1/3874 (0.0%)	1.00	13/5244 (0.2%)
2	B	0.69	0/3371	0.96	8/4541 (0.2%)
2	E	0.71	0/3371	0.96	2/4541 (0.0%)
2	H	0.72	0/3371	0.94	5/4541 (0.1%)
2	K	0.73	0/3371	0.97	5/4541 (0.1%)
2	N	0.72	0/3371	0.97	7/4541 (0.2%)
2	Q	0.73	0/3371	0.98	10/4541 (0.2%)
2	T	0.74	1/3371 (0.0%)	0.99	7/4541 (0.2%)
2	W	0.70	0/3371	0.96	7/4541 (0.2%)
3	C	0.68	0/778	0.96	2/1050 (0.2%)
3	F	0.67	0/778	0.89	0/1050
3	I	0.69	0/778	0.94	0/1050
3	L	0.67	0/778	0.98	2/1050 (0.2%)
3	O	0.70	0/778	0.92	1/1050 (0.1%)
3	R	0.69	0/778	1.01	3/1050 (0.3%)
3	U	0.72	0/778	0.96	0/1050
3	X	0.71	0/778	0.99	2/1050 (0.2%)
All	All	0.73	5/64184 (0.0%)	0.97	109/86680 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	M	0	1
2	B	0	2
2	E	0	2
2	T	0	1
3	F	0	1
All	All	0	7

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	299	ILE	CA-CB	9.11	1.59	1.54
1	G	299	ILE	CA-CB	8.71	1.59	1.54
1	V	299	ILE	CA-CB	7.46	1.58	1.54
2	T	190	GLY	C-O	5.30	1.30	1.24
1	G	35	THR	C-O	5.18	1.30	1.23

The worst 5 of 109 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	88	LEU	CA-C-N	8.55	128.13	119.24
2	B	88	LEU	C-N-CA	8.55	128.13	119.24
2	B	85	TYR	CA-C-N	8.41	128.91	119.32
2	B	85	TYR	C-N-CA	8.41	128.91	119.32
1	A	295	VAL	CB-CA-C	-7.87	101.73	112.04

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	217	VAL	Peptide
2	B	411	GLY	Peptide
2	E	217	VAL	Peptide
2	E	97	TYR	Peptide
3	F	45	THR	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3784	0	3817	123	0
1	D	3784	0	3816	110	0
1	G	3784	0	3817	105	0
1	J	3784	0	3816	112	0
1	M	3784	0	3817	112	0
1	P	3784	0	3816	122	0
1	S	3784	0	3817	122	0
1	V	3784	0	3817	121	0
2	B	3308	0	3353	96	0
2	E	3308	0	3353	97	0
2	H	3308	0	3353	85	0
2	K	3308	0	3353	80	0
2	N	3308	0	3353	74	0
2	Q	3308	0	3353	69	0
2	T	3308	0	3353	87	0
2	W	3308	0	3353	91	0
3	C	764	0	755	22	0
3	F	764	0	755	20	0
3	I	764	0	755	19	0
3	L	764	0	755	17	0
3	O	764	0	755	21	0
3	R	764	0	755	19	0
3	U	764	0	755	18	0
3	X	764	0	755	22	0
4	A	9	0	5	11	0
4	D	9	0	5	3	0
4	G	9	0	5	13	0
4	J	9	0	5	6	0
4	M	9	0	5	14	0
4	P	9	0	5	7	0
4	S	9	0	5	6	0
4	V	9	0	5	12	0
5	B	1	0	0	0	0
5	H	1	0	0	0	0
5	K	1	0	0	0	0
5	N	1	0	0	0	0
5	Q	1	0	0	0	0
5	T	1	0	0	0	0
5	W	1	0	0	0	0
6	B	1	0	0	0	0
6	E	1	0	0	0	0
6	H	1	0	0	0	0
6	K	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	N	1	0	0	0	0
6	Q	1	0	0	0	0
6	T	1	0	0	0	0
6	W	1	0	0	0	0
All	All	62935	0	63437	1638	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 1638 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:S:171:SER:OG	4:S:907:GLN:CD	1.75	1.27
2:B:412:LEU:HD22	2:B:412:LEU:O	1.39	1.22
1:P:464:LEU:HD12	1:P:464:LEU:O	1.40	1.20
1:G:171:SER:OG	4:G:903:GLN:CD	1.85	1.17
1:J:122:MET:HE2	1:J:199:PHE:CE1	1.80	1.17

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	476/478 (100%)	455 (96%)	20 (4%)	1 (0%)	43	72
1	D	476/478 (100%)	456 (96%)	18 (4%)	2 (0%)	30	60
1	G	476/478 (100%)	452 (95%)	23 (5%)	1 (0%)	43	72
1	J	476/478 (100%)	453 (95%)	23 (5%)	0	100	100
1	M	476/478 (100%)	453 (95%)	23 (5%)	0	100	100
1	P	476/478 (100%)	451 (95%)	24 (5%)	1 (0%)	43	72

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	S	476/478 (100%)	456 (96%)	20 (4%)	0	100	100
1	V	476/478 (100%)	456 (96%)	18 (4%)	2 (0%)	30	60
2	B	408/478 (85%)	396 (97%)	12 (3%)	0	100	100
2	E	408/478 (85%)	390 (96%)	18 (4%)	0	100	100
2	H	408/478 (85%)	395 (97%)	13 (3%)	0	100	100
2	K	408/478 (85%)	392 (96%)	16 (4%)	0	100	100
2	N	408/478 (85%)	395 (97%)	13 (3%)	0	100	100
2	Q	408/478 (85%)	393 (96%)	15 (4%)	0	100	100
2	T	408/478 (85%)	388 (95%)	20 (5%)	0	100	100
2	W	408/478 (85%)	396 (97%)	12 (3%)	0	100	100
3	C	89/94 (95%)	86 (97%)	3 (3%)	0	100	100
3	F	89/94 (95%)	83 (93%)	6 (7%)	0	100	100
3	I	89/94 (95%)	85 (96%)	4 (4%)	0	100	100
3	L	89/94 (95%)	86 (97%)	3 (3%)	0	100	100
3	O	89/94 (95%)	82 (92%)	7 (8%)	0	100	100
3	R	89/94 (95%)	83 (93%)	6 (7%)	0	100	100
3	U	89/94 (95%)	85 (96%)	4 (4%)	0	100	100
3	X	89/94 (95%)	84 (94%)	4 (4%)	1 (1%)	11	36
All	All	7784/8400 (93%)	7451 (96%)	325 (4%)	8 (0%)	48	77

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	P	148	SER
1	A	148	SER
1	V	148	SER
1	D	2	LEU
1	D	148	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	406/406 (100%)	382 (94%)	24 (6%)	18	48
1	D	406/406 (100%)	373 (92%)	33 (8%)	11	33
1	G	406/406 (100%)	387 (95%)	19 (5%)	23	57
1	J	406/406 (100%)	375 (92%)	31 (8%)	12	36
1	M	406/406 (100%)	382 (94%)	24 (6%)	18	48
1	P	406/406 (100%)	374 (92%)	32 (8%)	11	35
1	S	406/406 (100%)	379 (93%)	27 (7%)	15	42
1	V	406/406 (100%)	381 (94%)	25 (6%)	16	45
2	B	364/427 (85%)	340 (93%)	24 (7%)	15	43
2	E	364/427 (85%)	344 (94%)	20 (6%)	19	51
2	H	364/427 (85%)	341 (94%)	23 (6%)	16	45
2	K	364/427 (85%)	345 (95%)	19 (5%)	21	53
2	N	364/427 (85%)	337 (93%)	27 (7%)	13	37
2	Q	364/427 (85%)	342 (94%)	22 (6%)	17	47
2	T	364/427 (85%)	342 (94%)	22 (6%)	17	47
2	W	364/427 (85%)	342 (94%)	22 (6%)	17	47
3	C	86/89 (97%)	81 (94%)	5 (6%)	18	49
3	F	86/89 (97%)	81 (94%)	5 (6%)	18	49
3	I	86/89 (97%)	79 (92%)	7 (8%)	11	33
3	L	86/89 (97%)	81 (94%)	5 (6%)	18	49
3	O	86/89 (97%)	80 (93%)	6 (7%)	14	40
3	R	86/89 (97%)	82 (95%)	4 (5%)	23	57
3	U	86/89 (97%)	79 (92%)	7 (8%)	11	33
3	X	86/89 (97%)	81 (94%)	5 (6%)	18	49
All	All	6848/7376 (93%)	6410 (94%)	438 (6%)	16	44

5 of 438 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	M	245	VAL
1	P	162	VAL
1	V	181	VAL
1	M	478	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	N	390	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 43 such sidechains are listed below:

Mol	Chain	Res	Type
3	O	53	GLN
2	T	405	GLN
2	Q	218	ASN
2	T	110	ASN
1	V	53	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 23 ligands modelled in this entry, 15 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GLN	D	902	-	6,8,9	1.04	1 (16%)	6,9,11	0.94	1 (16%)
4	GLN	J	904	-	6,8,9	0.97	1 (16%)	6,9,11	1.03	1 (16%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GLN	G	903	-	6,8,9	1.17	0	6,9,11	1.06	1 (16%)
4	GLN	P	906	-	6,8,9	1.16	1 (16%)	6,9,11	1.29	1 (16%)
4	GLN	S	907	-	6,8,9	0.96	1 (16%)	6,9,11	1.19	1 (16%)
4	GLN	V	908	-	6,8,9	1.24	1 (16%)	6,9,11	0.95	0
4	GLN	A	901	-	6,8,9	1.17	1 (16%)	6,9,11	1.20	1 (16%)
4	GLN	M	905	-	6,8,9	1.00	1 (16%)	6,9,11	1.54	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GLN	D	902	-	-	1/8/8/9	-
4	GLN	J	904	-	-	2/8/8/9	-
4	GLN	G	903	-	-	1/8/8/9	-
4	GLN	P	906	-	-	3/8/8/9	-
4	GLN	S	907	-	-	1/8/8/9	-
4	GLN	V	908	-	-	1/8/8/9	-
4	GLN	A	901	-	-	1/8/8/9	-
4	GLN	M	905	-	-	1/8/8/9	-

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	P	906	GLN	OXT-C	-2.47	1.22	1.30
4	V	908	GLN	OXT-C	-2.34	1.23	1.30
4	A	901	GLN	OXT-C	-2.33	1.23	1.30
4	J	904	GLN	OXT-C	-2.13	1.23	1.30
4	S	907	GLN	OXT-C	-2.07	1.24	1.30

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	M	905	GLN	OXT-C-O	-3.61	115.89	124.08
4	P	906	GLN	OXT-C-O	-2.91	117.47	124.08
4	S	907	GLN	OXT-C-O	-2.79	117.76	124.08
4	A	901	GLN	OXT-C-O	-2.67	118.03	124.08
4	J	904	GLN	OXT-C-O	-2.43	118.56	124.08

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	901	GLN	OE1-CD-CG-CB
4	D	902	GLN	OE1-CD-CG-CB
4	G	903	GLN	OE1-CD-CG-CB
4	J	904	GLN	OE1-CD-CG-CB
4	M	905	GLN	OE1-CD-CG-CB

There are no ring outliers.

8 monomers are involved in 72 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	902	GLN	3	0
4	J	904	GLN	6	0
4	G	903	GLN	13	0
4	P	906	GLN	7	0
4	S	907	GLN	6	0
4	V	908	GLN	12	0
4	A	901	GLN	11	0
4	M	905	GLN	14	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	478/478 (100%)	-1.52	0 100 100	23, 52, 78, 87	0
1	D	478/478 (100%)	-1.57	0 100 100	23, 52, 78, 87	0
1	G	478/478 (100%)	-1.56	0 100 100	23, 51, 77, 87	0
1	J	478/478 (100%)	-1.51	0 100 100	23, 51, 77, 87	0
1	M	478/478 (100%)	-1.56	0 100 100	23, 51, 77, 87	0
1	P	478/478 (100%)	-1.51	0 100 100	23, 51, 77, 87	0
1	S	478/478 (100%)	-1.57	0 100 100	23, 52, 78, 87	0
1	V	478/478 (100%)	-1.52	0 100 100	23, 52, 78, 87	0
2	B	410/478 (85%)	-1.53	0 100 100	31, 61, 92, 111	0
2	E	410/478 (85%)	-1.55	0 100 100	31, 62, 92, 111	0
2	H	410/478 (85%)	-1.53	0 100 100	31, 61, 92, 111	0
2	K	410/478 (85%)	-1.53	0 100 100	31, 61, 92, 111	0
2	N	410/478 (85%)	-1.56	0 100 100	31, 62, 92, 111	0
2	Q	410/478 (85%)	-1.53	0 100 100	31, 61, 92, 111	0
2	T	410/478 (85%)	-1.55	0 100 100	31, 61, 92, 111	0
2	W	410/478 (85%)	-1.54	0 100 100	31, 61, 92, 111	0
3	C	91/94 (96%)	-1.60	0 100 100	25, 58, 71, 75	0
3	F	91/94 (96%)	-1.57	0 100 100	25, 58, 70, 75	0
3	I	91/94 (96%)	-1.66	0 100 100	25, 58, 71, 75	0
3	L	91/94 (96%)	-1.58	0 100 100	25, 58, 71, 75	0
3	O	91/94 (96%)	-1.58	0 100 100	25, 58, 71, 75	0
3	R	91/94 (96%)	-1.55	0 100 100	25, 58, 70, 75	0
3	U	91/94 (96%)	-1.60	0 100 100	25, 58, 71, 75	0
3	X	91/94 (96%)	-1.62	0 100 100	25, 58, 71, 75	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
All	All	7832/8400 (93%)	-1.55	0 100 100	23, 55, 87, 111	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MG	Q	479	1/1	0.68	0.18	61,61,61,61	1
5	MG	K	479	1/1	0.83	0.13	53,53,53,53	1
5	MG	N	479	1/1	0.86	0.15	86,86,86,86	1
5	MG	B	479	1/1	0.89	0.15	57,57,57,57	1
5	MG	H	479	1/1	0.91	0.10	61,61,61,61	1
5	MG	W	479	1/1	0.94	0.19	75,75,75,75	1
5	MG	T	479	1/1	0.95	0.04	60,60,60,60	1
4	GLN	V	908	9/10	0.99	0.04	49,51,54,59	0
4	GLN	A	901	9/10	0.99	0.05	48,49,55,59	0
4	GLN	D	902	9/10	0.99	0.06	55,56,57,61	0
4	GLN	G	903	9/10	0.99	0.05	51,52,60,65	0
4	GLN	J	904	9/10	0.99	0.05	59,59,60,63	0
4	GLN	M	905	9/10	0.99	0.06	47,48,53,60	0
4	GLN	P	906	9/10	0.99	0.06	52,53,54,56	0
4	GLN	S	907	9/10	0.99	0.05	55,56,57,62	0
6	ZN	B	901	1/1	1.00	0.01	49,49,49,49	0
6	ZN	E	902	1/1	1.00	0.01	59,59,59,59	0
6	ZN	H	903	1/1	1.00	0.01	51,51,51,51	0
6	ZN	K	904	1/1	1.00	0.00	46,46,46,46	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	ZN	N	905	1/1	1.00	0.01	47,47,47,47	0
6	ZN	Q	906	1/1	1.00	0.01	44,44,44,44	0
6	ZN	T	907	1/1	1.00	0.01	49,49,49,49	0
6	ZN	W	908	1/1	1.00	0.01	50,50,50,50	0

6.5 Other polymers [i](#)

There are no such residues in this entry.