



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 11:19 PM UTC

PDB ID : 3H18 / pdb_00003h18
Title : Crystal structure of EstE5-PMSF (II)
Authors : Hwang, K.Y.; Nam, K.H.
Deposited on : 2009-04-11
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	322	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PMS	A	310	-	X	X	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2251 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Esterase/lipase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	293	Total	C	N	O	S	0	0	0
			2214	1408	389	403	14			

There are 25 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-12	MET	-	expression tag	UNP Q0GMU2
A	-11	ALA	-	expression tag	UNP Q0GMU2
A	-10	SER	-	expression tag	UNP Q0GMU2
A	-9	MET	-	expression tag	UNP Q0GMU2
A	-8	THR	-	expression tag	UNP Q0GMU2
A	-7	GLY	-	expression tag	UNP Q0GMU2
A	-6	GLY	-	expression tag	UNP Q0GMU2
A	-5	GLN	-	expression tag	UNP Q0GMU2
A	-4	GLN	-	expression tag	UNP Q0GMU2
A	-3	MET	-	expression tag	UNP Q0GMU2
A	-2	GLY	-	expression tag	UNP Q0GMU2
A	-1	ARG	-	expression tag	UNP Q0GMU2
A	0	GLY	-	expression tag	UNP Q0GMU2
A	298	LEU	-	expression tag	UNP Q0GMU2
A	299	ALA	-	expression tag	UNP Q0GMU2
A	300	ALA	-	expression tag	UNP Q0GMU2
A	301	ALA	-	expression tag	UNP Q0GMU2
A	302	LEU	-	expression tag	UNP Q0GMU2
A	303	GLU	-	expression tag	UNP Q0GMU2
A	304	HIS	-	expression tag	UNP Q0GMU2
A	305	HIS	-	expression tag	UNP Q0GMU2
A	306	HIS	-	expression tag	UNP Q0GMU2
A	307	HIS	-	expression tag	UNP Q0GMU2
A	308	HIS	-	expression tag	UNP Q0GMU2
A	309	HIS	-	expression tag	UNP Q0GMU2

- Molecule 2 is phenylmethanesulfonic acid (CCD ID: PMS) (formula: C₇H₈O₃S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	S	0	0
			10	7	2	1		
2	A	1	Total	C	O	S	0	0
			10	7	2	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	17	Total	O	0	0
			17	17		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Esterase/lipase



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	61.28Å 61.28Å 149.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.36 – 2.40 28.36 – 2.40	Depositor EDS
% Data completeness (in resolution range)	91.9 (28.36-2.40) 91.8 (28.36-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.01 (at 2.39Å)	Xtriage
Refinement program	CNS, REFMAC 5.2.0019	Depositor
R, R_{free}	0.184 , 0.260 0.204 , 0.200	Depositor DCC
R_{free} test set	525 reflections (4.43%)	wwPDB-VP
Wilson B-factor (Å ²)	39.8	Xtriage
Anisotropy	0.376	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 28.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2251	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.65	18/2266 (0.8%)	1.59	31/3074 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	79	VAL	CA-CB	9.12	1.66	1.54
1	A	186	THR	CA-CB	6.89	1.64	1.53
1	A	203	ALA	CA-CB	-6.57	1.43	1.53
1	A	245	ILE	CA-CB	6.17	1.63	1.54
1	A	141	SER	C-O	-6.12	1.16	1.23
1	A	98	GLN	CA-C	-5.87	1.45	1.53
1	A	290	PHE	CA-C	5.72	1.60	1.52
1	A	114	PHE	C-O	5.65	1.31	1.24
1	A	107	ARG	CA-C	-5.53	1.45	1.52
1	A	61	ALA	CA-CB	-5.36	1.46	1.53
1	A	175	ALA	C-O	-5.28	1.16	1.23
1	A	285	VAL	CA-CB	5.21	1.62	1.54
1	A	167	SER	N-CA	5.20	1.52	1.46
1	A	6	ILE	CA-CB	5.15	1.61	1.54
1	A	152	ALA	C-O	-5.14	1.18	1.24
1	A	172	SER	CA-C	5.12	1.59	1.52
1	A	258	SER	CA-C	-5.09	1.46	1.52
1	A	272	ALA	CA-C	-5.07	1.45	1.52

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	19	PRO	CA-C-N	-9.70	110.06	119.76
1	A	19	PRO	C-N-CA	-9.70	110.06	119.76
1	A	96	ALA	N-CA-C	-7.26	103.39	112.90
1	A	49	VAL	CB-CA-C	-6.82	100.72	110.82
1	A	292	ARG	CA-C-N	6.73	130.92	120.82
1	A	292	ARG	C-N-CA	6.73	130.92	120.82
1	A	274	HIS	CA-C-N	-6.71	111.90	119.28
1	A	274	HIS	C-N-CA	-6.71	111.90	119.28
1	A	140	ILE	CB-CA-C	6.60	119.96	110.33
1	A	292	ARG	N-CA-C	6.55	120.49	112.23
1	A	88	SER	N-CA-C	6.46	118.91	111.02
1	A	287	VAL	N-CA-C	-6.43	104.22	110.72
1	A	61	ALA	CA-C-N	-5.94	114.49	120.31
1	A	61	ALA	C-N-CA	-5.94	114.49	120.31
1	A	271	HIS	N-CA-C	-5.83	104.28	111.75
1	A	77	GLY	N-CA-C	-5.72	107.35	115.30
1	A	218	SER	CA-C-N	5.66	125.75	119.87
1	A	218	SER	C-N-CA	5.66	125.75	119.87
1	A	214	HIS	N-CA-C	-5.61	101.12	109.42
1	A	112	HIS	CA-C-N	-5.54	114.89	120.21
1	A	112	HIS	C-N-CA	-5.54	114.89	120.21
1	A	281	LYS	N-CA-C	-5.52	105.34	111.36
1	A	120	ASP	N-CA-C	5.42	116.87	111.07
1	A	221	PHE	CB-CA-C	-5.27	102.53	111.02
1	A	105	ASP	CB-CA-C	-5.22	104.12	112.05
1	A	293	GLU	N-CA-C	5.18	119.92	111.37
1	A	85	THR	CA-C-N	-5.12	114.30	122.29
1	A	85	THR	C-N-CA	-5.12	114.30	122.29
1	A	6	ILE	N-CA-CB	5.08	118.20	110.58
1	A	86	HIS	CA-C-N	5.02	127.31	120.54
1	A	86	HIS	C-N-CA	5.02	127.31	120.54

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	192	PRO	Peptide
1	A	292	ARG	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2214	0	2219	61	0
2	A	20	0	14	7	0
3	A	17	0	0	2	0
All	All	2251	0	2233	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:80:MET:CE	2:A:310:PMS:H4	1.99	0.93
1:A:283:ALA:O	1:A:287:VAL:HG23	1.77	0.84
1:A:187:ARG:HD3	1:A:236:ARG:O	1.82	0.80
2:A:310:PMS:H2	2:A:311:PMS:S	2.25	0.76
1:A:223:ASN:HD21	1:A:225:LYS:HB2	1.51	0.76
1:A:233:HIS:HD2	1:A:271:HIS:NE2	1.85	0.74
1:A:80:MET:HE3	2:A:310:PMS:H4	1.71	0.70
1:A:168:ALA:C	1:A:169:ILE:HD12	2.17	0.69
1:A:104:LEU:HD13	1:A:106:TYR:HB3	1.76	0.67
1:A:64:CYS:HB2	3:A:407:HOH:O	1.93	0.67
1:A:19:PRO:O	1:A:22:THR:HG23	1.98	0.64
1:A:26:LEU:HD23	1:A:29:MET:HE3	1.80	0.63
1:A:169:ILE:HD12	1:A:169:ILE:N	2.15	0.62
1:A:169:ILE:N	1:A:169:ILE:CD1	2.65	0.59
1:A:279:GLU:OE1	1:A:279:GLU:N	2.31	0.57
1:A:77:GLY:O	1:A:78:TYR:HB2	2.05	0.57
1:A:140:ILE:HG12	1:A:150:VAL:HG13	1.86	0.56
1:A:159:ASP:OD2	1:A:216:TYR:OH	2.22	0.56
1:A:80:MET:HE2	2:A:310:PMS:H4	1.87	0.55
1:A:13:LEU:HD13	1:A:193:MET:HE1	1.89	0.52
1:A:233:HIS:CD2	1:A:271:HIS:NE2	2.73	0.52
1:A:181:ASN:ND2	1:A:246:LYS:NZ	2.58	0.52
1:A:134:LYS:O	1:A:137:HIS:N	2.38	0.52
1:A:134:LYS:O	1:A:135:PRO:C	2.51	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:287:VAL:HG12	1:A:291:MET:HE2	1.91	0.51
1:A:97:SER:HB3	1:A:288:GLY:HA2	1.93	0.51
1:A:271:HIS:H	1:A:271:HIS:HD1	1.59	0.51
1:A:60:ARG:HG2	1:A:100:ALA:CB	2.42	0.50
1:A:92:GLU:HG3	1:A:274:HIS:CE1	2.46	0.50
1:A:181:ASN:HB2	1:A:184:PHE:CE1	2.48	0.49
1:A:181:ASN:HB3	1:A:242:ASP:HB2	1.93	0.49
1:A:80:MET:HE3	2:A:310:PMS:C4	2.41	0.48
1:A:46:VAL:HG21	1:A:87:ARG:HD3	1.95	0.48
1:A:79:VAL:CG2	1:A:202:MET:HG2	2.44	0.48
1:A:114:PHE:HB2	1:A:206:TYR:CZ	2.48	0.48
1:A:268:HIS:NE2	2:A:310:PMS:H2A	2.28	0.48
1:A:140:ILE:HG13	1:A:150:VAL:HG22	1.94	0.47
1:A:169:ILE:HG21	1:A:287:VAL:HG13	1.96	0.47
1:A:278:PRO:O	1:A:282:GLN:HG2	2.15	0.47
1:A:26:LEU:HA	1:A:29:MET:HE3	1.95	0.47
1:A:136:GLN:H	1:A:136:GLN:HE21	1.62	0.47
1:A:182:ASP:N	1:A:242:ASP:OD2	2.49	0.46
1:A:60:ARG:HG2	1:A:100:ALA:HB2	1.97	0.46
1:A:190:ALA:O	1:A:192:PRO:HD3	2.17	0.45
1:A:223:ASN:ND2	1:A:225:LYS:H	2.15	0.44
1:A:73:LEU:HA	3:A:412:HOH:O	2.18	0.44
1:A:279:GLU:HA	1:A:282:GLN:CG	2.48	0.44
1:A:181:ASN:HB2	1:A:184:PHE:HE1	1.82	0.43
1:A:193:MET:HB2	1:A:193:MET:HE3	1.38	0.43
1:A:118:VAL:O	1:A:122:VAL:HG23	2.19	0.43
1:A:167:SER:HB3	1:A:294:GLN:HG3	2.00	0.43
1:A:109:ALA:HB1	1:A:110:PRO:HA	2.01	0.42
2:A:310:PMS:H2	2:A:311:PMS:O2S	2.19	0.42
1:A:151:LEU:O	1:A:155:VAL:HG23	2.20	0.42
1:A:144:SER:HA	1:A:172:SER:O	2.20	0.42
1:A:181:ASN:ND2	1:A:246:LYS:HZ2	2.18	0.42
1:A:248:ASP:O	1:A:252:LYS:HG2	2.19	0.42
1:A:184:PHE:O	1:A:188:ALA:HB2	2.19	0.42
1:A:79:VAL:HA	1:A:109:ALA:O	2.19	0.41
1:A:73:LEU:HD13	1:A:149:LEU:HD13	2.02	0.41
1:A:26:LEU:HD23	1:A:26:LEU:HA	1.94	0.41
1:A:70:ILE:O	1:A:102:LEU:N	2.42	0.41
1:A:24:VAL:CG1	1:A:28:VAL:HG12	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	289/322 (90%)	271 (94%)	18 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/246 (92%)	194 (86%)	32 (14%)	3	4

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	ILE
1	A	11	LYS
1	A	18	VAL
1	A	22	THR
1	A	23	GLU
1	A	26	LEU
1	A	27	ASP
1	A	33	MET
1	A	35	LYS
1	A	46	VAL
1	A	49	VAL
1	A	50	THR
1	A	65	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	102	LEU
1	A	104	LEU
1	A	136	GLN
1	A	140	ILE
1	A	149	LEU
1	A	169	ILE
1	A	193	MET
1	A	225	LYS
1	A	227	LEU
1	A	230	LEU
1	A	231	LEU
1	A	233	HIS
1	A	241	LEU
1	A	250	LYS
1	A	258	SER
1	A	260	LEU
1	A	261	GLU
1	A	281	LYS
1	A	293	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	98	GLN
1	A	136	GLN
1	A	181	ASN
1	A	223	ASN
1	A	233	HIS
1	A	274	HIS
1	A	282	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PMS	A	310	1	7,10,11	2.56	4 (57%)	11,12,15	3.91	9 (81%)
2	PMS	A	311	-	7,10,11	1.22	1 (14%)	11,12,15	4.20	5 (45%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PMS	A	310	1	-	1/4/4/5	0/1/1/1
2	PMS	A	311	-	-	4/4/4/5	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	310	PMS	C4-C3	4.04	1.47	1.38
2	A	310	PMS	C5-C4	3.23	1.45	1.38
2	A	310	PMS	C2-C1	2.69	1.44	1.38
2	A	310	PMS	C6-C1	2.51	1.43	1.38
2	A	311	PMS	C5-C4	2.23	1.43	1.38

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	311	PMS	O1S-S-C	8.38	120.02	105.55

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	311	PMS	C1-C-S	8.13	131.26	110.71
2	A	310	PMS	O2S-S-C	-5.98	95.24	105.55
2	A	311	PMS	C-C1-C6	-5.36	114.72	120.56
2	A	310	PMS	C4-C3-C2	-5.34	113.65	120.24
2	A	310	PMS	C5-C4-C3	4.94	126.64	119.87
2	A	310	PMS	C6-C1-C2	4.93	125.56	118.23
2	A	310	PMS	O1S-S-C	4.65	113.58	105.55
2	A	310	PMS	C-C1-C2	-3.79	116.43	120.56
2	A	311	PMS	O2S-S-C	-3.64	99.26	105.55
2	A	311	PMS	C6-C1-C2	2.84	122.45	118.23
2	A	310	PMS	C5-C6-C1	-2.79	116.69	120.61
2	A	310	PMS	C-C1-C6	-2.49	117.84	120.56
2	A	310	PMS	C4-C5-C6	-2.12	117.62	120.24

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	310	PMS	C1-C-S-O1S
2	A	311	PMS	C1-C-S-O1S
2	A	311	PMS	S-C-C1-C6
2	A	311	PMS	S-C-C1-C2
2	A	311	PMS	C1-C-S-O2S

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	310	PMS	7	0
2	A	311	PMS	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	293/322 (90%)	-0.49	2 (0%) 84 81	26, 37, 55, 63	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	ALA	4.8
1	A	297	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PMS	A	311	10/11	0.94	0.10	45,47,53,54	0
2	PMS	A	310	10/11	0.96	0.13	33,40,44,44	0

6.5 Other polymers [i](#)

There are no such residues in this entry.