



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 12:45 PM UTC

PDB ID : 3H1C / pdb_00003h1c
Title : Crystal structure of Polynucleotide Phosphorylase (PNPase) core bound to RNase E and Tungstate
Authors : Nurmohamed, S.
Deposited on : 2009-04-11
Resolution : 3.57 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1521 (3.66-3.50)
Clashscore	190562	1595 (3.66-3.50)
Ramachandran outliers	187476	1551 (3.66-3.50)
Sidechain outliers	187428	1551 (3.66-3.50)
RSRZ outliers	180081	1520 (3.66-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	549	3% 76% 22% ..
1	B	549	3% 75% 22% ..
1	C	549	3% 73% 24% ..
1	G	549	2% 76% 22% ..
1	I	549	% 76% 22% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	549	
1	M	549	
1	O	549	
1	R	549	
1	T	549	
1	V	549	
1	X	549	
2	D	41	
2	E	41	
2	F	41	
2	H	41	
2	J	41	
2	L	41	
2	N	41	
2	P	41	
2	S	41	
2	U	41	
2	W	41	
2	Y	41	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 51109 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polyribonucleotide nucleotidyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	544	Total	C	N	O	S	0	0	0
			4075	2559	704	792	20			
1	B	544	Total	C	N	O	S	0	0	0
			4107	2575	717	797	18			
1	C	544	Total	C	N	O	S	0	0	0
			4121	2583	720	798	20			
1	G	543	Total	C	N	O	S	0	0	0
			4095	2571	715	789	20			
1	I	543	Total	C	N	O	S	0	0	0
			4109	2576	716	797	20			
1	K	544	Total	C	N	O	S	0	0	0
			4127	2588	721	798	20			
1	M	544	Total	C	N	O	S	0	0	0
			4116	2581	718	797	20			
1	O	544	Total	C	N	O	S	0	0	0
			4124	2584	723	798	19			
1	R	544	Total	C	N	O	S	0	0	0
			4134	2590	724	800	20			
1	T	543	Total	C	N	O	S	0	0	0
			4116	2582	720	795	19			
1	V	544	Total	C	N	O	S	0	0	0
			4118	2580	719	799	20			
1	X	544	Total	C	N	O	S	0	0	0
			4113	2581	717	795	20			

- Molecule 2 is a protein called Ribonuclease E.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	21	Total	C	N	O	0	0	0
			138	82	31	25			
2	E	21	Total	C	N	O	0	0	0
			138	82	31	25			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	F	21	Total	C	N	O	0	0	0
			138	82	31	25			
2	H	21	Total	C	N	O	0	0	0
			138	82	31	25			
2	J	21	Total	C	N	O	0	0	0
			138	82	31	25			
2	L	21	Total	C	N	O	0	0	0
			138	82	31	25			
2	N	21	Total	C	N	O	0	0	0
			138	82	31	25			
2	P	21	Total	C	N	O	0	0	0
			138	82	31	25			
2	S	21	Total	C	N	O	0	0	0
			138	82	31	25			
2	U	21	Total	C	N	O	0	0	0
			138	82	31	25			
2	W	21	Total	C	N	O	0	0	0
			131	78	29	24			
2	Y	21	Total	C	N	O	0	0	0
			138	82	31	25			

- Molecule 3 is TUNGSTATE(VI)ION (CCD ID: WO4) (formula: O₄W).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	O W	0	0
			5 4 1			

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 5	O 4	W 1	0	0
3	B	1	Total 5	O 4	W 1	0	0
3	B	1	Total 5	O 4	W 1	0	0
3	C	1	Total 5	O 4	W 1	0	0
3	G	1	Total 5	O 4	W 1	0	0
3	I	1	Total 5	O 4	W 1	0	0
3	K	1	Total 5	O 4	W 1	0	0
3	K	1	Total 5	O 4	W 1	0	0
3	M	1	Total 5	O 4	W 1	0	0
3	M	1	Total 5	O 4	W 1	0	0
3	O	1	Total 5	O 4	W 1	0	0
3	O	1	Total 5	O 4	W 1	0	0
3	R	1	Total 5	O 4	W 1	0	0
3	R	1	Total 5	O 4	W 1	0	0
3	T	1	Total 5	O 4	W 1	0	0
3	T	1	Total 5	O 4	W 1	0	0
3	V	1	Total 5	O 4	W 1	0	0
3	V	1	Total 5	O 4	W 1	0	0
3	X	1	Total 5	O 4	W 1	0	0
3	X	1	Total 5	O 4	W 1	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Polyribonucleotide nucleotidyltransferase




• Molecule 1: Polyribonucleotide nucleotidyltransferase




ASP
ILE
SER


• Molecule 1: Polyribonucleotide nucleotidyltransferase




• Molecule 1: Polyribonucleotide nucleotidyltransferase





• Molecule 1: Polyribonucleotide nucleotidyltransferase








● Molecule 1: Polyribonucleotide nucleotidyltransferase




● Molecule 1: Polyribonucleotide nucleotidyltransferase




● Molecule 1: Polyribonucleotide nucleotidyltransferase







- Molecule 1: Polyribonucleotide nucleotidyltransferase




- Molecule 1: Polyrribonucleotide nucleotidyltransferase





- Molecule 1: Polyrribonucleotide nucleotidyltransferase









• Molecule 1: Polyribonucleotide nucleotidyltransferase




• Molecule 2: Ribonuclease E




• Molecule 2: Ribonuclease E




• Molecule 2: Ribonuclease E









• Molecule 2: Ribonuclease E




• Molecule 2: Ribonuclease E




• Molecule 2: Ribonuclease E





• Molecule 2: Ribonuclease E




• Molecule 2: Ribonuclease E




• Molecule 2: Ribonuclease E




• Molecule 2: Ribonuclease E



● Molecule 2: Ribonuclease E



● Molecule 2: Ribonuclease E



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	167.74Å 262.89Å 264.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 3.57 25.00 – 3.57	Depositor EDS
% Data completeness (in resolution range)	94.2 (25.00-3.57) 93.6 (25.00-3.57)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.57 (at 3.62Å)	Xtriage
Refinement program	REFMAC 5.0	Depositor
R, R_{free}	0.270 , 0.304 0.267 , 0.291	Depositor DCC
R_{free} test set	6514 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	79.4	Xtriage
Anisotropy	0.028	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 26.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	0.038 for -h,l,k	Xtriage
F_o, F_c correlation	0.85	EDS
Total number of atoms	51109	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: WO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	1/4138 (0.0%)	0.77	2/5621 (0.0%)
1	B	0.43	0/4170	0.78	2/5660 (0.0%)
1	C	0.50	1/4184 (0.0%)	0.78	2/5675 (0.0%)
1	G	0.58	2/4158 (0.0%)	0.84	7/5643 (0.1%)
1	I	0.51	0/4172	0.80	5/5660 (0.1%)
1	K	0.52	2/4190 (0.0%)	0.85	5/5681 (0.1%)
1	M	0.45	1/4179 (0.0%)	0.87	8/5668 (0.1%)
1	O	0.48	2/4187 (0.0%)	0.78	4/5679 (0.1%)
1	R	0.42	0/4197	0.77	2/5690 (0.0%)
1	T	0.46	0/4179	0.79	4/5667 (0.1%)
1	V	0.49	1/4180 (0.0%)	0.78	3/5671 (0.1%)
1	X	0.43	0/4175	0.79	5/5661 (0.1%)
2	D	0.55	0/143	0.77	0/196
2	E	0.47	0/143	0.62	0/196
2	F	0.46	0/143	0.72	0/196
2	H	0.48	0/143	0.65	0/196
2	J	0.49	0/143	0.83	0/196
2	L	0.53	0/143	0.71	0/196
2	N	0.49	0/143	0.66	0/196
2	P	0.47	0/143	0.71	0/196
2	S	0.49	0/143	0.69	0/196
2	U	0.45	0/143	0.67	0/196
2	W	0.76	0/135	1.21	2/185 (1.1%)
2	Y	0.46	0/143	0.76	0/196
All	All	0.48	10/51817 (0.0%)	0.80	51/70317 (0.1%)

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	514	ILE	CA-CB	-11.30	1.47	1.55
1	C	120	VAL	CA-CB	-7.03	1.45	1.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	V	333	VAL	CA-CB	-6.77	1.45	1.54
1	K	64	GLN	C-N	6.54	1.43	1.33
1	G	186	VAL	CA-CB	-6.14	1.46	1.54

The worst 5 of 51 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	M	434	SER	CB-CA-C	-17.91	84.47	110.62
1	K	37	ASP	CB-CA-C	-13.11	84.33	110.42
1	M	438	SER	CB-CA-C	13.06	136.42	110.42
1	M	435	ASN	N-CA-CB	-11.18	91.60	110.49
1	M	434	SER	N-CA-C	11.03	126.89	107.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4075	0	4044	156	0
1	B	4107	0	4096	133	0
1	C	4121	0	4126	192	0
1	G	4095	0	4087	168	0
1	I	4109	0	4108	141	0
1	K	4127	0	4141	154	0
1	M	4116	0	4116	101	0
1	O	4124	0	4130	124	0
1	R	4134	0	4150	107	0
1	T	4116	0	4127	157	0
1	V	4118	0	4115	176	0
1	X	4113	0	4117	185	0
2	D	138	0	125	5	0
2	E	138	0	125	8	0
2	F	138	0	125	16	0
2	H	138	0	125	2	0
2	J	138	0	125	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	L	138	0	125	8	0
2	N	138	0	125	0	0
2	P	138	0	125	1	0
2	S	138	0	125	10	0
2	U	138	0	125	4	0
2	W	131	0	115	21	0
2	Y	138	0	125	5	0
3	A	10	0	0	0	0
3	B	10	0	0	0	0
3	C	5	0	0	0	0
3	G	5	0	0	1	0
3	I	5	0	0	0	0
3	K	10	0	0	0	0
3	M	10	0	0	0	0
3	O	10	0	0	0	0
3	R	10	0	0	0	0
3	T	10	0	0	0	0
3	V	10	0	0	0	0
3	X	10	0	0	1	0
All	All	51109	0	50847	1696	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 1696 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:310:LEU:HD21	1:I:469:VAL:CG1	1.47	1.44
1:T:491:MET:HE1	1:T:493:PHE:CD1	1.52	1.44
1:G:26:GLN:NE2	1:K:387:VAL:HA	1.36	1.40
1:R:51:LYS:HB3	1:R:54:GLN:CG	1.52	1.36
1:O:239:ASN:ND2	1:O:242:LEU:HB2	1.40	1.35

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	542/549 (99%)	496 (92%)	40 (7%)	6 (1%)	11	43
1	B	542/549 (99%)	508 (94%)	30 (6%)	4 (1%)	18	51
1	C	542/549 (99%)	497 (92%)	39 (7%)	6 (1%)	11	43
1	G	541/549 (98%)	500 (92%)	38 (7%)	3 (1%)	21	54
1	I	541/549 (98%)	502 (93%)	36 (7%)	3 (1%)	21	54
1	K	542/549 (99%)	502 (93%)	34 (6%)	6 (1%)	11	43
1	M	542/549 (99%)	501 (92%)	37 (7%)	4 (1%)	18	51
1	O	542/549 (99%)	507 (94%)	32 (6%)	3 (1%)	21	54
1	R	542/549 (99%)	505 (93%)	33 (6%)	4 (1%)	18	51
1	T	541/549 (98%)	499 (92%)	36 (7%)	6 (1%)	11	43
1	V	542/549 (99%)	506 (93%)	33 (6%)	3 (1%)	21	54
1	X	542/549 (99%)	510 (94%)	29 (5%)	3 (1%)	21	54
2	D	19/41 (46%)	19 (100%)	0	0	100	100
2	E	19/41 (46%)	19 (100%)	0	0	100	100
2	F	19/41 (46%)	17 (90%)	2 (10%)	0	100	100
2	H	19/41 (46%)	19 (100%)	0	0	100	100
2	J	19/41 (46%)	18 (95%)	1 (5%)	0	100	100
2	L	19/41 (46%)	19 (100%)	0	0	100	100
2	N	19/41 (46%)	17 (90%)	2 (10%)	0	100	100
2	P	19/41 (46%)	18 (95%)	1 (5%)	0	100	100
2	S	19/41 (46%)	19 (100%)	0	0	100	100
2	U	19/41 (46%)	18 (95%)	1 (5%)	0	100	100
2	W	19/41 (46%)	13 (68%)	4 (21%)	2 (10%)	0	5
2	Y	19/41 (46%)	19 (100%)	0	0	100	100
All	All	6729/7080 (95%)	6248 (93%)	428 (6%)	53 (1%)	16	49

5 of 53 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	38	THR
1	G	187	LEU
1	G	189	VAL
1	O	37	ASP
1	O	38	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	426/446 (96%)	420 (99%)	6 (1%)	59	71
1	B	432/446 (97%)	425 (98%)	7 (2%)	55	69
1	C	436/446 (98%)	428 (98%)	8 (2%)	51	68
1	G	429/446 (96%)	423 (99%)	6 (1%)	59	71
1	I	435/446 (98%)	428 (98%)	7 (2%)	55	69
1	K	437/446 (98%)	433 (99%)	4 (1%)	70	76
1	M	434/446 (97%)	423 (98%)	11 (2%)	42	63
1	O	436/446 (98%)	423 (97%)	13 (3%)	36	60
1	R	439/446 (98%)	434 (99%)	5 (1%)	65	74
1	T	435/446 (98%)	430 (99%)	5 (1%)	65	74
1	V	435/446 (98%)	428 (98%)	7 (2%)	55	69
1	X	433/446 (97%)	430 (99%)	3 (1%)	76	78
2	D	11/28 (39%)	11 (100%)	0	100	100
2	E	11/28 (39%)	11 (100%)	0	100	100
2	F	11/28 (39%)	11 (100%)	0	100	100
2	H	11/28 (39%)	11 (100%)	0	100	100
2	J	11/28 (39%)	11 (100%)	0	100	100
2	L	11/28 (39%)	11 (100%)	0	100	100
2	N	11/28 (39%)	11 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	P	11/28 (39%)	11 (100%)	0	100	100
2	S	11/28 (39%)	11 (100%)	0	100	100
2	U	11/28 (39%)	11 (100%)	0	100	100
2	W	9/28 (32%)	8 (89%)	1 (11%)	6	27
2	Y	11/28 (39%)	11 (100%)	0	100	100
All	All	5337/5688 (94%)	5254 (98%)	83 (2%)	55	69

5 of 83 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	O	344	THR
1	T	518	ILE
1	O	509	ILE
1	R	411	VAL
1	V	310	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 129 such sidechains are listed below:

Mol	Chain	Res	Type
1	V	239	ASN
1	V	532	HIS
1	I	239	ASN
1	I	218	ASN
1	X	26	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

21 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	WO4	X	551	-	2,4,4	12.55	2 (100%)	-		
3	WO4	B	550	-	2,4,4	12.56	2 (100%)	-		
3	WO4	A	550	-	2,4,4	12.54	2 (100%)	-		
3	WO4	I	550	-	2,4,4	12.51	2 (100%)	-		
3	WO4	O	550	-	2,4,4	12.55	2 (100%)	-		
3	WO4	M	551	-	2,4,4	12.53	2 (100%)	-		
3	WO4	A	551	-	2,4,4	12.53	2 (100%)	-		
3	WO4	V	550	-	2,4,4	12.54	2 (100%)	-		
3	WO4	O	551	-	2,4,4	12.54	2 (100%)	-		
3	WO4	M	550	-	2,4,4	11.42	2 (100%)	-		
3	WO4	T	551	-	2,4,4	12.55	2 (100%)	-		
3	WO4	C	550	-	2,4,4	12.52	2 (100%)	-		
3	WO4	T	550	-	2,4,4	12.52	2 (100%)	-		
3	WO4	K	550	-	2,4,4	12.50	2 (100%)	-		
3	WO4	V	551	-	2,4,4	12.52	2 (100%)	-		
3	WO4	R	551	-	2,4,4	12.54	2 (100%)	-		
3	WO4	G	550	-	2,4,4	12.52	2 (100%)	-		
3	WO4	K	551	-	2,4,4	12.53	2 (100%)	-		
3	WO4	R	550	-	2,4,4	12.53	2 (100%)	-		
3	WO4	X	550	-	2,4,4	12.52	2 (100%)	-		
3	WO4	B	551	-	2,4,4	12.53	2 (100%)	-		

The worst 5 of 42 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	550	WO4	W-O2	12.57	2.04	1.77

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	X	551	WO4	W-O2	12.56	2.04	1.77
3	A	550	WO4	W-O2	12.56	2.04	1.77
3	T	551	WO4	W-O2	12.56	2.04	1.77
3	O	550	WO4	W-O2	12.56	2.04	1.77

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	G	550	WO4	1	0
3	X	550	WO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	544/549 (99%)	0.42	14 (2%)	57	32	15, 50, 83, 103	1 (0%)
1	B	544/549 (99%)	0.39	14 (2%)	57	32	9, 39, 76, 101	1 (0%)
1	C	544/549 (99%)	0.44	18 (3%)	49	27	14, 48, 83, 106	1 (0%)
1	G	543/549 (98%)	0.39	13 (2%)	59	34	20, 51, 83, 111	1 (0%)
1	I	543/549 (98%)	0.37	8 (1%)	72	44	14, 40, 76, 98	1 (0%)
1	K	544/549 (99%)	0.47	16 (2%)	53	30	13, 43, 82, 115	1 (0%)
1	M	544/549 (99%)	0.53	26 (4%)	35	19	20, 54, 89, 110	1 (0%)
1	O	544/549 (99%)	0.43	11 (2%)	65	37	8, 41, 79, 97	1 (0%)
1	R	544/549 (99%)	0.43	17 (3%)	51	28	13, 46, 83, 101	1 (0%)
1	T	543/549 (98%)	0.46	15 (2%)	55	30	20, 53, 84, 108	1 (0%)
1	V	544/549 (99%)	0.48	19 (3%)	47	26	19, 62, 92, 105	1 (0%)
1	X	544/549 (99%)	0.49	15 (2%)	55	30	20, 52, 85, 100	1 (0%)
2	D	21/41 (51%)	1.44	5 (23%)	2	2	44, 80, 95, 96	0
2	E	21/41 (51%)	1.48	7 (33%)	1	1	34, 69, 98, 104	0
2	F	21/41 (51%)	1.84	10 (47%)	0	1	70, 92, 111, 117	0
2	H	21/41 (51%)	1.65	6 (28%)	1	1	62, 95, 107, 109	0
2	J	21/41 (51%)	1.23	5 (23%)	2	2	43, 70, 85, 91	0
2	L	21/41 (51%)	1.42	5 (23%)	2	2	49, 78, 103, 107	0
2	N	21/41 (51%)	1.01	2 (9%)	14	9	64, 84, 97, 97	0
2	P	21/41 (51%)	1.24	4 (19%)	3	3	33, 73, 93, 98	0
2	S	21/41 (51%)	1.47	5 (23%)	2	2	64, 96, 117, 120	0
2	U	21/41 (51%)	1.00	2 (9%)	14	9	49, 91, 107, 111	0
2	W	21/41 (51%)	1.60	6 (28%)	1	1	78, 111, 130, 134	0
2	Y	21/41 (51%)	1.09	2 (9%)	14	9	62, 90, 101, 103	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
All	All	6777/7080 (95%)	0.48	245 (3%) 46 25	8, 49, 88, 134	12 (0%)

The worst 5 of 245 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	159	ASP	6.5
1	K	159	ASP	6.4
1	T	434	SER	6.2
1	A	159	ASP	5.9
1	C	37	ASP	5.3

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	WO4	M	550	5/5	0.68	0.26	112,113,118,118	5
3	WO4	T	551	5/5	0.68	0.31	113,114,117,119	5
3	WO4	K	550	5/5	0.72	0.26	78,79,82,88	5
3	WO4	M	551	5/5	0.75	0.25	131,133,133,137	5
3	WO4	V	550	5/5	0.76	0.20	107,108,112,113	5
3	WO4	K	551	5/5	0.81	0.23	96,97,101,108	5
3	WO4	C	550	5/5	0.82	0.19	80,84,89,90	5
3	WO4	G	550	5/5	0.82	0.22	79,79,88,93	5
3	WO4	R	550	5/5	0.84	0.18	106,107,111,116	5
3	WO4	X	550	5/5	0.84	0.20	94,99,99,100	5
3	WO4	T	550	5/5	0.87	0.19	103,104,113,115	5
3	WO4	O	550	5/5	0.89	0.15	100,100,105,109	5
3	WO4	A	550	5/5	0.89	0.19	84,86,88,93	5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	WO4	V	551	5/5	0.89	0.18	102,102,105,109	5
3	WO4	A	551	5/5	0.89	0.16	103,105,108,113	5
3	WO4	R	551	5/5	0.90	0.22	86,88,95,98	5
3	WO4	X	551	5/5	0.90	0.24	88,90,91,93	5
3	WO4	B	551	5/5	0.91	0.17	88,89,93,97	5
3	WO4	B	550	5/5	0.92	0.15	72,73,77,82	5
3	WO4	O	551	5/5	0.92	0.18	71,75,80,87	5
3	WO4	I	550	5/5	0.93	0.15	89,90,91,91	5

6.5 Other polymers [i](#)

There are no such residues in this entry.