



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 04:13 AM UTC

PDB ID : 6H1O / pdb_00006h1o
Title : Structure of the BM3 heme domain in complex with voriconazole
Authors : Jeffreys, L.N.; Munro, A.W.M.; Leys, D.
Deposited on : 2018-07-12
Resolution : 1.73 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1187 (1.74-1.74)
Clashscore	190562	1207 (1.74-1.74)
Ramachandran outliers	187476	1200 (1.74-1.74)
Sidechain outliers	187428	1200 (1.74-1.74)
RSRZ outliers	180081	1188 (1.74-1.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	457	5% 91% 8%
1	B	457	2% 94% 6%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 8338 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Bifunctional cytochrome P450/NADPH-P450 reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	456	Total	C	N	O	S	0	5	0
			3716	2373	635	691	17			
1	B	457	Total	C	N	O	S	0	4	0
			3712	2369	631	695	17			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	82	PHE	ALA	engineered mutation	UNP P14779
A	87	VAL	PHE	engineered mutation	UNP P14779
B	82	PHE	ALA	engineered mutation	UNP P14779
B	87	VAL	PHE	engineered mutation	UNP P14779

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 3 is Voriconazole (CCD ID: VOR) (formula: C₁₆H₁₄F₃N₅O) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	0	0
			25	16	3	5	1		
3	B	1	Total	C	F	N	O	0	0
			25	16	3	5	1		

- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		
4	A	1	Total	O	P	0	0
			5	4	1		
4	A	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	1
			12	6	6		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	326	Total	O	0	0
			326	326		
7	B	389	Total	O	0	0
			389	389		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bifunctional cytochrome P450/NADPH-P450 reductase



- Molecule 1: Bifunctional cytochrome P450/NADPH-P450 reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	59.85Å 150.01Å 61.07Å 90.00° 96.85° 90.00°	Depositor
Resolution (Å)	60.63 – 1.73 60.63 – 1.73	Depositor EDS
% Data completeness (in resolution range)	90.6 (60.63-1.73) 90.6 (60.63-1.73)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 1.73Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.161 , 0.200 0.162 , 0.201	Depositor DCC
R_{free} test set	4831 reflections (4.38%)	wwPDB-VP
Wilson B-factor (Å ²)	21.2	Xtriage
Anisotropy	0.295	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 34.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.023 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8338	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, VOR, GOL, EDO, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/3806	0.60	0/5140
1	B	0.41	0/3802	0.58	0/5136
All	All	0.40	0/7608	0.59	0/10276

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	189	GLN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3716	0	3694	37	0
1	B	3712	0	3683	18	0
2	A	43	0	30	1	0
2	B	43	0	30	2	0
3	A	25	0	14	0	0
3	B	25	0	14	1	0
4	A	15	0	0	0	0
4	B	10	0	0	0	0
5	A	24	0	31	1	0
5	B	6	0	8	0	0
6	A	4	0	6	0	0
7	A	326	0	0	7	1
7	B	389	0	0	1	0
All	All	8338	0	7510	55	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:199:ASP:HB2	1:A:202:LYS:HA	1.50	0.94
1:A:199:ASP:OD1	1:A:205:PHE:N	2.11	0.83
1:A:177:MET:HE3	1:A:181:LEU:HD11	1.62	0.82
1:A:195:ASP:O	1:A:201:ASN:ND2	2.16	0.79
1:B:296[B]:ARG:NH1	7:B:601:HOH:O	2.17	0.78
1:A:169:GLN:NE2	7:A:601:HOH:O	2.19	0.75
1:A:195:ASP:HB2	1:A:196:PRO:HD3	1.75	0.69
1:A:388:HIS:HA	1:A:391:LYS:HD3	1.74	0.69
1:A:3:LYS:HE3	1:A:344:GLU:HB3	1.75	0.68
1:A:189:GLN:O	1:A:198:TYR:OH	2.14	0.66
1:B:118:MET:HE2	1:B:155:LEU:HG	1.78	0.64
1:A:199:ASP:HB2	1:A:202:LYS:CA	2.29	0.57
1:A:118:MET:HG2	7:A:871:HOH:O	2.06	0.56
1:A:199:ASP:CB	1:A:202:LYS:HA	2.31	0.56
1:B:267:GLU:OE1	3:B:502:VOR:H21	2.05	0.56
1:A:190:ARG:NH2	7:A:605:HOH:O	2.39	0.55
1:A:342:GLY:O	1:A:344:GLU:HG3	2.05	0.55
1:B:388:HIS:HA	1:B:391:LYS:HD3	1.89	0.54
1:B:44:ALA:HB3	1:B:47:ARG:HG2	1.90	0.54
1:A:183:GLU:OE1	1:A:198:TYR:HB3	2.08	0.53
1:A:34:ASP:OD1	1:A:359:GLN:NE2	2.40	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:118:MET:HE2	1:B:155:LEU:CD1	2.41	0.51
1:A:194:ASP:OD1	1:A:195:ASP:N	2.42	0.50
1:A:188:LEU:C	1:A:189:GLN:HG2	2.36	0.50
1:A:75:LEU:HD21	5:A:508:GOL:H11	1.93	0.50
1:A:170:PRO:HB3	1:A:174:ILE:HD11	1.94	0.50
1:A:128:GLN:NE2	7:A:607:HOH:O	2.42	0.49
1:A:170:PRO:CB	1:A:174:ILE:HD11	2.43	0.48
1:A:190:ARG:NE	7:A:608:HOH:O	2.45	0.48
1:A:192:ASN:HB2	1:A:193:PRO:HD3	1.96	0.48
1:B:43:GLU:HG2	1:B:48:VAL:HG22	1.94	0.48
1:A:3:LYS:NZ	1:A:344:GLU:OE2	2.39	0.48
1:A:2:ILE:HG12	1:A:344:GLU:HA	1.96	0.47
1:A:199:ASP:OD1	1:A:205:PHE:CB	2.63	0.47
1:A:5:MET:HE3	1:A:5:MET:HB2	1.70	0.46
1:B:1:THR:O	1:B:344:GLU:HG2	2.16	0.46
1:B:118:MET:HE2	1:B:155:LEU:CG	2.45	0.45
1:B:87:VAL:HG22	2:B:501:HEM:C3D	2.52	0.45
1:B:354:MET:HB2	1:B:354:MET:HE2	1.81	0.45
1:A:196:PRO:C	1:A:201:ASN:HD22	2.25	0.45
1:A:199:ASP:OD1	1:A:205:PHE:HB2	2.16	0.44
1:B:272:LEU:HD13	1:B:322:LEU:HG	1.99	0.44
1:B:241:LYS:HD3	1:B:248:PRO:HB3	2.00	0.44
1:B:58:ILE:HD12	1:B:360:LEU:HD13	1.99	0.43
1:A:56:ARG:NH1	1:A:344:GLU:OE1	2.51	0.43
1:A:167:ARG:NH1	7:A:601:HOH:O	2.24	0.43
1:A:187:LYS:C	1:A:189:GLN:H	2.26	0.43
1:A:382:PRO:HA	1:A:385:ILE:HD12	1.99	0.43
1:B:400:CYS:HB2	2:B:501:HEM:NA	2.34	0.43
1:B:75:LEU:HA	1:B:78:VAL:HG22	2.00	0.43
1:B:60:GLU:HG2	1:B:341:LEU:HD12	2.01	0.42
1:A:58:ILE:HD13	1:A:355:VAL:HG13	2.02	0.41
1:A:400:CYS:HA	2:A:501:HEM:CHA	2.49	0.41
1:A:343:GLY:N	7:A:614:HOH:O	2.50	0.41
1:B:135:ALA:O	1:B:136:ASP:HB2	2.20	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:608:HOH:O	7:A:855:HOH:O[1_455]	1.99	0.21

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	457/457 (100%)	435 (95%)	19 (4%)	3 (1%)	18	6
1	B	457/457 (100%)	447 (98%)	10 (2%)	0	100	100
All	All	914/914 (100%)	882 (96%)	29 (3%)	3 (0%)	36	23

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	190	ARG
1	A	197	ALA
1	A	195	ASP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	404/400 (101%)	404 (100%)	0	100	100
1	B	404/400 (101%)	404 (100%)	0	100	100
All	All	808/800 (101%)	808 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	21	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	70	ASN
1	A	201	ASN
1	B	70	ASN
1	B	169	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GOL	A	506[A]	-	5,5,5	0.45	0	5,5,5	0.28	0
4	PO4	B	504	-	4,4,4	0.68	0	6,6,6	0.66	0
5	GOL	A	507	-	5,5,5	0.31	0	5,5,5	0.49	0
2	HEM	B	501	3,1	50,50,50	1.38	7 (14%)	67,82,82	1.43	10 (14%)
6	EDO	A	509	-	3,3,3	0.43	0	2,2,2	0.37	0
5	GOL	A	506[B]	-	5,5,5	0.41	0	5,5,5	0.71	0
4	PO4	B	503	-	4,4,4	0.94	0	6,6,6	0.43	0
5	GOL	A	508	-	5,5,5	0.36	0	5,5,5	0.61	0
4	PO4	A	504	-	4,4,4	0.90	0	6,6,6	0.58	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	VOR	B	502	2	27,27,27	0.99	1 (3%)	38,39,39	4.40	21 (55%)
2	HEM	A	501	3,1	50,50,50	1.34	7 (14%)	67,82,82	1.23	3 (4%)
4	PO4	A	503	-	4,4,4	0.94	0	6,6,6	0.44	0
5	GOL	B	505	-	5,5,5	0.25	0	5,5,5	0.51	0
3	VOR	A	502	2	27,27,27	0.99	1 (3%)	38,39,39	4.98	20 (52%)
4	PO4	A	505	-	4,4,4	0.90	0	6,6,6	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	506[A]	-	-	2/4/4/4	-
5	GOL	A	507	-	-	2/4/4/4	-
2	HEM	B	501	3,1	-	2/14/54/54	-
6	EDO	A	509	-	-	0/1/1/1	-
5	GOL	A	506[B]	-	-	4/4/4/4	-
5	GOL	A	508	-	-	2/4/4/4	-
3	VOR	B	502	2	-	6/21/21/21	0/3/3/3
2	HEM	A	501	3,1	-	2/14/54/54	-
5	GOL	B	505	-	-	0/4/4/4	-
3	VOR	A	502	2	-	7/21/21/21	0/3/3/3

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	HEM	FE-NB	4.28	2.08	1.94
2	A	501	HEM	CAC-C3C	3.50	1.56	1.47
3	B	502	VOR	C18-C15	-3.49	1.32	1.38
2	A	501	HEM	FE-NB	3.20	2.04	1.94
2	B	501	HEM	CAB-C3B	3.20	1.55	1.47
2	A	501	HEM	CAB-C3B	2.92	1.55	1.47
2	B	501	HEM	FE-ND	2.83	2.03	1.94
2	A	501	HEM	FE-NA	2.78	2.04	1.95
2	B	501	HEM	CAC-C3C	2.72	1.54	1.47
2	A	501	HEM	FE-ND	2.49	2.02	1.94
3	A	502	VOR	C18-C15	-2.39	1.34	1.38
2	A	501	HEM	CMB-C2B	2.33	1.55	1.50
2	B	501	HEM	CMC-C2C	2.18	1.55	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	HEM	FE-NC	2.12	2.02	1.95
2	A	501	HEM	CMC-C2C	2.07	1.55	1.50
2	B	501	HEM	CMB-C2B	2.05	1.55	1.50

All (54) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	VOR	F2-C18-C15	23.29	138.72	120.11
3	B	502	VOR	F2-C18-C15	19.51	135.69	120.11
3	A	502	VOR	F2-C18-C23	-10.61	102.01	118.60
3	B	502	VOR	F2-C18-C23	-8.55	105.22	118.60
3	B	502	VOR	C18-C15-C11	-5.75	114.02	123.14
3	A	502	VOR	C17-C13-C10	-5.32	114.60	120.79
3	B	502	VOR	C17-C13-C10	-5.14	114.80	120.79
3	A	502	VOR	N9-C24-N6	-5.05	118.74	127.33
3	A	502	VOR	C18-C15-C11	-4.96	115.27	123.14
3	A	502	VOR	C18-C15-N6	4.93	121.99	118.09
3	B	502	VOR	C18-C15-N6	4.92	121.99	118.09
3	A	502	VOR	C10-C11-C15	4.80	120.64	112.67
3	B	502	VOR	C12-N5-N7	4.68	128.07	121.25
3	A	502	VOR	C23-N9-C24	4.54	121.15	115.80
3	A	502	VOR	C19-C16-C13	-4.53	118.96	123.88
3	A	502	VOR	C12-N5-N7	4.51	127.82	121.25
3	B	502	VOR	C13-C10-C11	4.29	116.92	110.88
3	B	502	VOR	C10-C11-C15	4.10	119.48	112.67
3	B	502	VOR	C14-C11-C15	-3.95	103.49	108.68
2	A	501	HEM	CHC-C1C-NC	3.91	128.71	124.45
3	B	502	VOR	N9-C24-N6	-3.75	120.97	127.33
3	B	502	VOR	C19-C16-C13	-3.56	120.01	123.88
3	B	502	VOR	C11-C15-N6	3.55	123.88	118.67
2	B	501	HEM	CHD-C1D-ND	3.53	128.23	124.42
2	B	501	HEM	C4D-ND-C1D	3.48	109.33	105.21
3	A	502	VOR	C16-C19-C22	3.34	120.23	116.67
3	B	502	VOR	C16-C19-C22	3.23	120.11	116.67
3	B	502	VOR	F1-C16-C13	3.18	122.51	119.01
2	B	501	HEM	C1B-NB-C4B	3.07	108.85	105.21
3	A	502	VOR	C13-C10-C11	3.07	115.19	110.88
3	B	502	VOR	F3-C22-C20	2.96	123.30	118.55
2	B	501	HEM	C3D-C4D-ND	-2.90	106.99	110.17
3	A	502	VOR	F1-C16-C13	2.90	122.20	119.01
2	B	501	HEM	C3B-C4B-NB	-2.83	107.43	109.47
3	A	502	VOR	C17-C13-C16	2.79	120.20	116.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	VOR	C14-C11-C15	-2.77	105.03	108.68
3	A	502	VOR	C11-C15-N6	2.73	122.67	118.67
2	B	501	HEM	C2D-C1D-ND	-2.58	106.92	109.90
3	B	502	VOR	C12-N5-C21	-2.53	123.88	128.82
2	A	501	HEM	C2A-C1A-NA	-2.48	107.40	110.15
3	A	502	VOR	N7-C25-N8	-2.48	107.83	114.90
3	A	502	VOR	C12-N5-C21	-2.45	124.04	128.82
3	A	502	VOR	C25-N7-N5	2.41	108.16	102.60
2	B	501	HEM	CHA-C4D-ND	2.36	127.29	124.37
2	A	501	HEM	CHD-C4C-NC	2.31	126.97	124.45
3	B	502	VOR	C10-C13-C16	2.31	125.76	123.17
3	B	502	VOR	N7-C25-N8	-2.28	108.39	114.90
2	B	501	HEM	CHB-C4A-NA	2.27	127.98	123.86
2	B	501	HEM	C3B-C2B-C1B	2.25	108.10	106.41
3	B	502	VOR	C25-N7-N5	2.25	107.80	102.60
3	B	502	VOR	C17-C13-C16	2.21	119.39	116.28
3	B	502	VOR	C20-C22-C19	-2.16	120.38	123.23
2	B	501	HEM	O2A-CGA-CBA	2.12	120.70	114.00
3	A	502	VOR	C20-C22-C19	-2.02	120.57	123.23

There are no chirality outliers.

All (27) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502	VOR	O4-C10-C11-C14
3	A	502	VOR	C12-C10-C11-C14
3	A	502	VOR	C12-C10-C11-C15
3	A	502	VOR	C13-C10-C11-C15
3	A	502	VOR	C10-C11-C15-N6
3	B	502	VOR	O4-C10-C11-C14
3	B	502	VOR	C12-C10-C11-C14
3	B	502	VOR	C12-C10-C11-C15
3	B	502	VOR	C13-C10-C11-C15
5	A	506[B]	GOL	O1-C1-C2-O2
5	A	506[B]	GOL	O1-C1-C2-C3
5	A	506[B]	GOL	C1-C2-C3-O3
5	A	507	GOL	O1-C1-C2-C3
5	A	508	GOL	O1-C1-C2-O2
5	A	508	GOL	O1-C1-C2-C3
5	A	506[A]	GOL	O1-C1-C2-C3
5	A	506[B]	GOL	O2-C2-C3-O3
5	A	507	GOL	O1-C1-C2-O2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	A	506[A]	GOL	O1-C1-C2-O2
3	A	502	VOR	C10-C11-C15-C18
3	A	502	VOR	C14-C11-C15-N6
3	B	502	VOR	C10-C11-C15-N6
3	B	502	VOR	C14-C11-C15-N6
2	B	501	HEM	CAD-CBD-CGD-O1D
2	B	501	HEM	CAD-CBD-CGD-O2D
2	A	501	HEM	CAD-CBD-CGD-O2D
2	A	501	HEM	CAD-CBD-CGD-O1D


There are no ring outliers.


4 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	HEM	2	0
5	A	508	GOL	1	0
3	B	502	VOR	1	0
2	A	501	HEM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	1
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	455:LEU	C	456:GLY	N	6.26
1	A	455:LEU	C	456:GLY	N	5.31

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	456/457 (99%)	0.18	25 (5%)	30 37	10, 24, 53, 90	5 (1%)
1	B	457/457 (100%)	-0.13	10 (2%)	62 70	13, 22, 41, 78	4 (0%)
All	All	913/914 (99%)	0.03	35 (3%)	44 53	10, 23, 47, 90	9 (0%)

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	457	GLY	7.3
1	A	198	TYR	5.6
1	A	202	LYS	5.0
1	A	2	ILE	5.0
1	A	191	ALA	4.7
1	A	456	GLY	4.5
1	A	196	PRO	4.1
1	A	197	ALA	4.1
1	A	190	ARG	4.0
1	A	188	LEU	3.9
1	B	1	THR	3.8
1	B	46	GLY	3.8
1	B	457	GLY	3.7
1	A	199	ASP	3.6
1	A	193	PRO	3.5
1	A	200	GLU	3.2
1	A	195	ASP	3.2
1	B	456	GLY	3.1
1	B	227	GLY	2.9
1	A	455	LEU	2.9
1	A	174	ILE	2.8
1	A	285	HIS	2.8
1	A	201	ASN	2.7
1	A	169	GLN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	168	ASP	2.6
1	A	194	ASP	2.5
1	B	2	ILE	2.5
1	A	189	GLN	2.4
1	B	437	LEU	2.4
1	B	455	LEU	2.4
1	A	225	ALA	2.3
1	B	228	GLU	2.2
1	B	136	ASP	2.1
1	A	454	PRO	2.1
1	A	186	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PO4	B	504	5/5	0.68	0.14	44,45,77,82	0
5	GOL	A	508	6/6	0.73	0.22	44,55,55,68	0
5	GOL	A	507	6/6	0.78	0.18	26,50,52,54	0
4	PO4	A	504	5/5	0.78	0.18	55,65,89,99	0
6	EDO	A	509	4/4	0.81	0.17	45,59,65,66	0
4	PO4	B	503	5/5	0.82	0.14	65,65,82,90	0
4	PO4	A	505	5/5	0.83	0.14	34,69,79,88	0
5	GOL	B	505	6/6	0.85	0.13	35,40,43,52	0
4	PO4	A	503	5/5	0.86	0.11	47,62,74,81	0
5	GOL	A	506[A]	6/6	0.88	0.13	19,23,31,41	6
5	GOL	A	506[B]	6/6	0.88	0.13	24,25,30,35	6
3	VOR	A	502	25/25	0.92	0.08	17,27,33,35	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	VOR	B	502	25/25	0.95	0.07	14,22,27,28	0
2	HEM	A	501	43/43	0.98	0.05	12,16,19,19	0
2	HEM	B	501	43/43	0.99	0.05	11,13,16,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around VOR A 502:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around VOR B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.