



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:21 PM UTC

PDB ID : 1H22 / pdb_00001h22
Title : Structure of acetylcholinesterase (E.C. 3.1.1.7) complexed with (S,S)-(-)-bis(10)-hupyrindone at 2.15Å resolution
Authors : Wong, D.M.; Greenblatt, H.M.; Carlier, P.R.; Han, Y.-F.; Pang, Y.-P.; Silman, I.; Sussman, J.L.
Deposited on : 2002-07-30
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	543	

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 4475 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACETYLCHOLINESTERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	529	4191	2692	710	767	22	0	2	1

- Molecule 2 is (S,S)-(-)-N,N'-DI-5'-[5',6',7',8'-TETRAHYDRO- 2'(1'H)-QUINOLYNYL]-1,10-DIAMINODECANE DIHYDROCHLORIDE (CCD ID: E10) (formula: C₂₈H₄₂N₄O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	34	28	4	2	0	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	222	Total	O	0	0
			222	222		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ACETYLCHOLINESTERASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	111.46Å 111.46Å 137.34Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	26.40 – 2.15 26.40 – 2.15	Depositor EDS
% Data completeness (in resolution range)	95.0 (26.40-2.15) 95.3 (26.40-2.15)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.91 (at 2.15Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.190 , 0.221 0.178 , 0.207	Depositor DCC
R_{free} test set	2549 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	32.3	Xtriage
Anisotropy	0.596	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 33.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.022 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4475	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: E10, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.15	9/4312 (0.2%)	1.23	24/5861 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	175	MET	SD-CE	-17.70	1.35	1.79
1	A	16	MET	SD-CE	-7.76	1.60	1.79
1	A	379	MET	SD-CE	7.45	1.98	1.79
1	A	43	MET	SD-CE	6.59	1.96	1.79
1	A	436	MET	SD-CE	6.15	1.95	1.79
1	A	354	SER	CA-C	5.86	1.60	1.52
1	A	535	THR	C-N	-5.67	1.25	1.33
1	A	356	VAL	CA-CB	5.12	1.61	1.54
1	A	347	ILE	CA-CB	5.08	1.60	1.54

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	518	VAL	CB-CA-C	-12.19	96.36	111.97
1	A	401	ILE	N-CA-C	7.70	117.82	110.42
1	A	491	LYS	N-CA-C	7.22	120.43	108.96
1	A	291	SER	N-CA-C	7.20	118.81	110.97

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	527	PHE	N-CA-C	7.07	118.78	111.14
1	A	518	VAL	N-CA-CB	6.96	118.69	110.55
1	A	285	ASP	N-CA-C	-6.92	99.41	109.59
1	A	297	ASP	N-CA-C	6.70	121.62	113.38
1	A	329	SER	N-CA-C	6.48	119.17	111.33
1	A	439	ILE	N-CA-C	6.43	118.36	109.80
1	A	216	ARG	N-CA-C	6.41	119.09	111.33
1	A	346	LYS	N-CA-C	-6.39	100.82	110.28
1	A	442	TYR	N-CA-C	6.12	120.44	113.15
1	A	117	GLY	N-CA-C	6.00	127.41	113.18
1	A	504	ASP	N-CA-C	-5.90	100.94	110.32
1	A	257	ASN	N-CA-C	5.70	118.27	111.71
1	A	255	ASN	N-CA-C	-5.68	101.44	109.96
1	A	183	ASN	N-CA-C	5.62	121.56	114.31
1	A	57	VAL	N-CA-C	5.51	115.79	107.80
1	A	363	ALA	N-CA-C	5.37	118.57	109.76
1	A	248	LEU	N-CA-C	-5.33	105.36	111.07
1	A	287	ILE	CB-CG1-CD1	-5.16	102.96	113.80
1	A	282	LEU	CA-C-N	-5.00	114.59	119.64
1	A	282	LEU	C-N-CA	-5.00	114.59	119.64

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	442	TYR	Sidechain
1	A	472	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4191	0	4005	76	0
2	A	34	0	42	0	0
3	A	28	0	26	0	0
4	A	222	0	0	5	0
All	All	4475	0	4073	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:VAL:HG12	4:A:2092:HOH:O	1.65	0.96
1:A:221:ARG:HD3	1:A:318:GLN:OE1	1.70	0.91
1:A:9:ASN:HD21	1:A:14:LYS:HE2	1.40	0.87
1:A:455:GLU:CD	1:A:455:GLU:H	1.89	0.81
1:A:9:ASN:ND2	1:A:14:LYS:HE2	1.98	0.78
1:A:6:LEU:O	1:A:16:MET:HE3	1.89	0.73
1:A:132:GLY:HA3	1:A:143[B]:LEU:HD23	1.71	0.70
1:A:6:LEU:HB2	1:A:16:MET:CE	2.22	0.70
1:A:453:VAL:CG2	1:A:456:LEU:HG	2.22	0.68
1:A:346:LYS:HG2	1:A:385:ILE:HD13	1.76	0.68
1:A:19:ARG:NH2	1:A:26:HIS:HB2	2.08	0.67
1:A:287:ILE:HD11	1:A:335:GLY:C	2.19	0.67
1:A:194:VAL:HG13	1:A:219:PHE:HA	1.76	0.67
1:A:287:ILE:HD11	1:A:335:GLY:O	1.94	0.66
1:A:383:ASN:C	1:A:383:ASN:HD22	2.05	0.65
1:A:508:GLU:HB2	1:A:509:PRO:HD2	1.79	0.64
1:A:36:ALA:HB2	1:A:175:MET:CE	2.28	0.64
1:A:6:LEU:C	1:A:16:MET:HE3	2.24	0.62
1:A:508:GLU:HB2	1:A:509:PRO:CD	2.30	0.61
1:A:27:ILE:HD12	1:A:27:ILE:C	2.27	0.59
1:A:74:GLN:HG2	4:A:2033:HOH:O	2.02	0.58
1:A:6:LEU:CB	1:A:16:MET:CE	2.82	0.58
1:A:314:PHE:O	1:A:316:LYS:HD2	2.04	0.57
1:A:20:VAL:HG22	1:A:27:ILE:HD12	1.87	0.56
1:A:36:ALA:HB2	1:A:175:MET:HE2	1.88	0.56
1:A:461:GLU:H	1:A:461:GLU:CD	2.12	0.56
1:A:47:ARG:NE	4:A:2011:HOH:O	2.38	0.56
1:A:197:PHE:CB	1:A:223:ILE:HB	2.36	0.56
1:A:78:PHE:O	1:A:82:GLU:HG3	2.06	0.56
1:A:449:GLY:HA2	1:A:466:SER:OG	2.07	0.55
1:A:5:GLU:OE2	1:A:104:PRO:HA	2.06	0.55
1:A:426:ARG:CZ	1:A:430:LEU:HD12	2.37	0.55
1:A:46:ARG:NH1	1:A:163:GLU:OE2	2.38	0.55
1:A:19:ARG:CZ	1:A:26:HIS:HB2	2.37	0.55
1:A:531:LEU:C	1:A:531:LEU:HD23	2.32	0.55
1:A:194:VAL:CG1	1:A:219:PHE:HA	2.39	0.53
1:A:181:HIS:HD2	4:A:2093:HOH:O	1.93	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:146:LEU:C	1:A:146:LEU:HD12	2.35	0.52
1:A:452:LEU:HD13	1:A:467:ARG:NH2	2.25	0.51
1:A:20:VAL:HG22	1:A:27:ILE:CD1	2.39	0.51
1:A:36:ALA:CB	1:A:175:MET:CE	2.88	0.51
1:A:506:ASN:OD1	1:A:508:GLU:HG2	2.11	0.50
1:A:223:ILE:HA	1:A:320:LEU:O	2.11	0.50
1:A:228:SER:HB2	1:A:229:PRO:HD2	1.93	0.50
1:A:220:ARG:HG3	1:A:221:ARG:HG3	1.94	0.50
1:A:453:VAL:CG2	1:A:453:VAL:O	2.60	0.50
1:A:515:ARG:HB3	1:A:518:VAL:HG22	1.94	0.49
1:A:455:GLU:CD	1:A:455:GLU:N	2.63	0.48
1:A:252:LEU:HB3	1:A:269:LYS:HE3	1.96	0.47
1:A:451:PRO:HA	1:A:458:TYR:CD1	2.48	0.47
1:A:455:GLU:N	1:A:455:GLU:OE2	2.46	0.47
1:A:312:GLY:O	1:A:316:LYS:HE3	2.16	0.46
1:A:47:ARG:HG3	4:A:2078:HOH:O	2.15	0.46
1:A:197:PHE:HB3	1:A:223:ILE:HB	1.97	0.46
1:A:20:VAL:CG2	1:A:27:ILE:HD11	2.46	0.45
1:A:456:LEU:O	1:A:457:ASN:HB2	2.17	0.45
1:A:408:VAL:CG2	1:A:418:THR:HG21	2.47	0.45
1:A:405:MET:HA	1:A:408:VAL:HG12	1.99	0.45
1:A:185:GLN:HA	1:A:189:GLY:O	2.17	0.45
1:A:349:ARG:HA	1:A:349:ARG:HD2	1.79	0.45
1:A:197:PHE:HB2	1:A:223:ILE:HB	1.98	0.45
1:A:453:VAL:HG22	1:A:456:LEU:HG	1.99	0.45
1:A:206:VAL:CG1	1:A:222:ALA:HB1	2.47	0.44
1:A:426:ARG:NH2	1:A:430:LEU:HD12	2.32	0.44
1:A:314:PHE:O	1:A:316:LYS:CD	2.66	0.43
1:A:199:GLU:HG3	1:A:443:GLU:OE2	2.19	0.42
1:A:528:LEU:HB3	1:A:529:PRO:HD3	2.00	0.41
1:A:212:SER:HA	1:A:213:PRO:HD3	1.91	0.41
1:A:332:LEU:HD11	1:A:392:ASP:HA	2.03	0.41
1:A:421:TYR:CD1	1:A:421:TYR:C	2.98	0.41
1:A:20:VAL:CG2	1:A:27:ILE:CD1	2.99	0.40
1:A:383:ASN:C	1:A:383:ASN:ND2	2.77	0.40
1:A:450:LEU:N	1:A:451:PRO:CD	2.84	0.40
1:A:377:ASP:OD1	1:A:377:ASP:C	2.64	0.40
1:A:379:MET:HE2	1:A:379:MET:HB3	2.01	0.40
1:A:491:LYS:O	1:A:493:PRO:HD3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	527/543 (97%)	502 (95%)	25 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	451/474 (95%)	434 (96%)	17 (4%)	29	29

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	74	GLN
1	A	129	VAL
1	A	140	GLU
1	A	194	VAL
1	A	287	ILE
1	A	294	PRO
1	A	354	SER
1	A	356	VAL
1	A	358	LEU
1	A	383	ASN
1	A	430	LEU
1	A	434	GLU
1	A	452	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	453	VAL
1	A	455	GLU
1	A	461	GLU
1	A	518	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	ASN
1	A	68	GLN
1	A	74	GLN
1	A	181	HIS
1	A	257	ASN
1	A	374	GLN
1	A	383	ASN
1	A	457	ASN
1	A	526	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	E10	A	1536	-	37,37,37	2.11	14 (37%)	40,48,48	1.26	7 (17%)
3	NAG	A	1537	1	14,14,15	1.19	3 (21%)	17,19,21	1.07	2 (11%)
3	NAG	A	1538	1	14,14,15	0.78	0	17,19,21	0.98	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	E10	A	1536	-	-	4/15/35/35	0/4/4/4
3	NAG	A	1537	1	-	2/6/23/26	0/1/1/1
3	NAG	A	1538	1	-	2/6/23/26	0/1/1/1

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1536	E10	C37-C32	4.54	1.57	1.51
2	A	1536	E10	C34-C29	4.40	1.53	1.43
2	A	1536	E10	C6-C1	3.85	1.52	1.43
2	A	1536	E10	C41-N40	3.64	1.54	1.47
2	A	1536	E10	C10-N17	3.50	1.54	1.46
2	A	1536	E10	C9-C3	3.44	1.55	1.49
2	A	1536	E10	C36-C31	2.93	1.54	1.49
2	A	1536	E10	C24-N17	2.77	1.52	1.47
2	A	1536	E10	C37-N40	2.59	1.52	1.46
2	A	1536	E10	C10-C4	2.59	1.54	1.51
3	A	1537	NAG	C1-C2	-2.52	1.48	1.52
2	A	1536	E10	C13-C10	2.52	1.57	1.53
2	A	1536	E10	C34-C33	2.40	1.41	1.35
2	A	1536	E10	C38-C37	2.35	1.57	1.53
3	A	1537	NAG	C4-C3	2.26	1.58	1.52
2	A	1536	E10	C31-C32	2.26	1.40	1.37
3	A	1537	NAG	O5-C5	2.08	1.47	1.43

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1536	E10	C29-N30-C31	3.11	127.60	124.83
2	A	1536	E10	C38-C37-N40	-2.86	103.00	111.82
2	A	1536	E10	C1-N2-C3	2.83	127.34	124.83
2	A	1536	E10	C41-N40-C37	2.52	118.27	113.76
2	A	1536	E10	C24-N17-C10	2.39	118.05	113.76
3	A	1537	NAG	C3-C4-C5	2.26	114.33	110.23
2	A	1536	E10	C33-C34-C29	-2.16	118.59	120.66
2	A	1536	E10	C14-C13-C10	2.15	113.81	110.97
3	A	1538	NAG	C2-N2-C7	-2.04	120.17	122.90
3	A	1537	NAG	O5-C1-C2	-2.01	108.17	111.29
3	A	1538	NAG	C3-C4-C5	-2.00	106.60	110.23

There are no chirality outliers.


All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1538	NAG	O5-C5-C6-O6
3	A	1538	NAG	C4-C5-C6-O6
3	A	1537	NAG	O5-C5-C6-O6
2	A	1536	E10	C67-C68-C69-C70
2	A	1536	E10	C64-C65-C66-C67
2	A	1536	E10	C68-C69-C70-C41
2	A	1536	E10	C65-C66-C67-C68
3	A	1537	NAG	C1-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	529/543 (97%)	-0.28	8 (1%) 72 76	15, 32, 46, 70	2 (0%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	485	PRO	3.6
1	A	535	THR	3.3
1	A	490	SER	2.9
1	A	379	MET	2.7
1	A	536	ALA	2.7
1	A	534	ALA	2.5
1	A	347	ILE	2.1
1	A	26	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	A	1537	14/15	0.69	0.18	57,62,64,65	0
3	NAG	A	1538	14/15	0.89	0.10	44,47,51,51	0
2	E10	A	1536	34/34	0.96	0.07	29,32,43,45	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.