



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 12:44 AM UTC

PDB ID : 1H2Z / pdb_00001h2z
Title : PROLYL OLIGOPEPTIDASE FROM PORCINE BRAIN, S554A MUTANT
WITH BOUND PEPTIDE LIGAND SUC-GLY-PRO
Authors : Rea, D.; Fulop, V.
Deposited on : 2002-08-20
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2662 (1.66-1.66)
Ramachandran outliers	187476	2621 (1.66-1.66)
Sidechain outliers	187428	2621 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	710	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6792 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PROLYL ENDOPEPTIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	710	Total	C	N	O	S	0	0	0
			5700	3659	944	1069	28			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	554	ALA	SER	engineered mutation	UNP P23687

- Molecule 2 is GLYCINE (CCD ID: GLY) (formula: C₂H₅NO₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			4	2	1	1		

- Molecule 3 is PROLINE (CCD ID: PRO) (formula: C₅H₉NO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			8	5	1	2		

- Molecule 4 is SUCCINIC ACID (CCD ID: SIN) (formula: $C_4H_6O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1055	Total	O	0	0
			1055	1055		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: PROLYL ENDOPEPTIDASE



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	71.40 Å 100.20 Å 111.40 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	52.00 – 1.65	Depositor
% Data completeness (in resolution range)	98.8 (52.00-1.65)	Depositor
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.202 , 0.228	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6792	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SIN, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/5854	0.94	24/7938 (0.3%)

There are no bond length outliers.

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	406	ILE	N-CA-C	-9.79	94.02	108.12
1	A	320	ASP	CA-C-N	8.35	128.00	119.82
1	A	320	ASP	C-N-CA	8.35	128.00	119.82
1	A	321	PRO	N-CA-C	7.46	123.61	113.98
1	A	260	ARG	N-CA-C	-7.05	101.03	110.55
1	A	405	GLY	N-CA-C	6.50	120.86	112.54
1	A	506	GLY	N-CA-C	-6.00	107.55	114.69
1	A	174	SER	N-CA-C	5.82	119.53	110.17
1	A	175	CYS	N-CA-C	-5.81	102.16	110.59
1	A	566	GLN	N-CA-C	5.75	117.63	111.36
1	A	599	TYR	N-CA-C	5.74	120.41	113.41
1	A	303	LYS	N-CA-C	-5.74	99.94	109.46
1	A	157	LYS	N-CA-C	-5.52	100.30	109.46
1	A	579	GLY	N-CA-C	5.48	120.04	112.37
1	A	179	THR	N-CA-C	-5.39	102.90	110.50
1	A	659	ILE	N-CA-C	5.35	115.56	110.53
1	A	404	PRO	N-CA-C	-5.34	101.48	112.47
1	A	77	SER	N-CA-C	-5.28	102.38	110.14
1	A	231	GLU	CA-C-N	5.20	125.50	119.47
1	A	231	GLU	C-N-CA	5.20	125.50	119.47
1	A	643	ARG	N-CA-C	-5.17	104.58	111.24
1	A	600	GLY	N-CA-C	-5.16	104.06	112.83
1	A	79	HIS	N-CA-C	-5.05	103.38	110.50
1	A	611	LEU	N-CA-C	5.00	116.42	111.07

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5700	0	5516	23	0
2	A	4	0	2	0	0
3	A	8	0	7	2	0
4	A	7	0	4	0	0
5	A	18	0	24	3	0
6	A	1055	0	0	5	0
All	All	6792	0	5553	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:345:ARG:HH21	1:A:347:ASN:HD21	1.25	0.84
1:A:418:GLU:OE2	6:A:2771:HOH:O	2.10	0.69
1:A:461:LYS:HD2	1:A:463:ASP:HB3	1.75	0.68
1:A:347:ASN:H	1:A:347:ASN:HD22	1.43	0.64
1:A:645:VAL:HG23	1:A:647:LEU:HG	1.84	0.59
1:A:25:CYS:HB3	6:A:2059:HOH:O	2.07	0.55
1:A:554:ALA:HB1	3:A:726:PRO:C	2.33	0.53
1:A:595:TRP:CD1	3:A:726:PRO:HD3	2.45	0.52
1:A:362:GLN:HG2	1:A:374:ILE:HD13	1.93	0.50
1:A:341:VAL:HG13	1:A:351:LEU:HD22	1.94	0.48
1:A:440:ILE:C	1:A:440:ILE:HD12	2.39	0.48
1:A:257:PRO:O	1:A:289:GLU:HB2	2.14	0.47
1:A:398:PHE:O	1:A:405:GLY:HA2	2.14	0.46
1:A:323:GLU:HA	1:A:326:TRP:CE2	2.51	0.46
1:A:546:LYS:O	1:A:704:LEU:HD22	2.14	0.46
1:A:353:TYR:HB2	1:A:360:THR:OG1	2.16	0.46
5:A:792:GOL:H11	6:A:3051:HOH:O	2.15	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:794:GOL:H31	6:A:2553:HOH:O	2.17	0.45
1:A:28:TYR:HB3	1:A:31:LEU:HD12	1.99	0.45
1:A:128:ARG:HD2	1:A:151:VAL:CG2	2.50	0.42
1:A:2:LEU:HA	6:A:2003:HOH:O	2.19	0.42
1:A:6:TYR:HA	1:A:7:PRO:HD3	1.93	0.41
1:A:481:THR:HB	1:A:482:PRO:CD	2.51	0.40
1:A:305:ASN:HB3	1:A:311:TYR:CE2	2.56	0.40
1:A:121:ASP:OD1	5:A:792:GOL:H2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	708/710 (100%)	686 (97%)	21 (3%)	1 (0%)	48	30

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	357	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	616/616 (100%)	608 (99%)	8 (1%)	61 42

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48	LYS
1	A	59	ILE
1	A	218	ASP
1	A	347	ASN
1	A	351	LEU
1	A	361	LEU
1	A	426	THR
1	A	598	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	192	GLN
1	A	315	ASN
1	A	347	ASN
1	A	477	ASN
1	A	522	ASN
1	A	551	ASN
1	A	577	GLN
1	A	667	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PRO	A	726	-	8,8,8	0.75	0	10,10,10	0.93	0
5	GOL	A	794	-	5,5,5	0.20	0	5,5,5	0.12	0
5	GOL	A	792	-	5,5,5	0.33	0	5,5,5	0.24	0
4	SIN	A	724	2	5,6,7	0.98	0	6,6,8	0.84	0
5	GOL	A	791	-	5,5,5	0.25	0	5,5,5	0.27	0
2	GLY	A	725	4	3,3,4	0.51	0	1,2,4	0.85	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PRO	A	726	-	-	4/4/11/11	0/1/1/1
5	GOL	A	794	-	-	2/4/4/4	-
5	GOL	A	792	-	-	2/4/4/4	-
4	SIN	A	724	2	-	2/4/4/5	-
5	GOL	A	791	-	-	0/4/4/4	-
2	GLY	A	725	4	-	0/0/1/2	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	726	PRO	O-C-CA-N
3	A	726	PRO	OXT-C-CA-N
5	A	794	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	A	792	GOL	O1-C1-C2-C3
5	A	794	GOL	O1-C1-C2-O2
3	A	726	PRO	OXT-C-CA-CB
3	A	726	PRO	O-C-CA-CB
5	A	792	GOL	O1-C1-C2-O2
4	A	724	SIN	O1-C1-C2-C3
4	A	724	SIN	O2-C1-C2-C3

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	726	PRO	2	0
5	A	794	GOL	1	0
5	A	792	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.