



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 01:18 AM UTC

PDB ID : 3H24 / pdb_00003h24
Title : Structural Studies of Pterin-Based Inhibitors of Dihydropteroate Synthase
Authors : Yun, M.-K.; White, S.W.
Deposited on : 2009-04-14
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	297	6% 66% 19% 5% 10%
1	B	297	6% 62% 24% • 12%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4219 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dihydropteroate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	267	Total	C	N	O	S	0	0	0
			2060	1294	354	396	16			
1	B	260	Total	C	N	O	S	0	0	0
			2012	1265	347	384	16			

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP B1UXN2
A	-18	GLY	-	expression tag	UNP B1UXN2
A	-17	SER	-	expression tag	UNP B1UXN2
A	-16	SER	-	expression tag	UNP B1UXN2
A	-15	HIS	-	expression tag	UNP B1UXN2
A	-14	HIS	-	expression tag	UNP B1UXN2
A	-13	HIS	-	expression tag	UNP B1UXN2
A	-12	HIS	-	expression tag	UNP B1UXN2
A	-11	HIS	-	expression tag	UNP B1UXN2
A	-10	HIS	-	expression tag	UNP B1UXN2
A	-9	SER	-	expression tag	UNP B1UXN2
A	-8	SER	-	expression tag	UNP B1UXN2
A	-7	GLY	-	expression tag	UNP B1UXN2
A	-6	LEU	-	expression tag	UNP B1UXN2
A	-5	VAL	-	expression tag	UNP B1UXN2
A	-4	PRO	-	expression tag	UNP B1UXN2
A	-3	ARG	-	expression tag	UNP B1UXN2
A	-2	GLY	-	expression tag	UNP B1UXN2
A	-1	SER	-	expression tag	UNP B1UXN2
A	0	HIS	-	expression tag	UNP B1UXN2
B	-19	MET	-	expression tag	UNP B1UXN2
B	-18	GLY	-	expression tag	UNP B1UXN2
B	-17	SER	-	expression tag	UNP B1UXN2
B	-16	SER	-	expression tag	UNP B1UXN2
B	-15	HIS	-	expression tag	UNP B1UXN2

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	expression tag	UNP B1UXN2
B	-13	HIS	-	expression tag	UNP B1UXN2
B	-12	HIS	-	expression tag	UNP B1UXN2
B	-11	HIS	-	expression tag	UNP B1UXN2
B	-10	HIS	-	expression tag	UNP B1UXN2
B	-9	SER	-	expression tag	UNP B1UXN2
B	-8	SER	-	expression tag	UNP B1UXN2
B	-7	GLY	-	expression tag	UNP B1UXN2
B	-6	LEU	-	expression tag	UNP B1UXN2
B	-5	VAL	-	expression tag	UNP B1UXN2
B	-4	PRO	-	expression tag	UNP B1UXN2
B	-3	ARG	-	expression tag	UNP B1UXN2
B	-2	GLY	-	expression tag	UNP B1UXN2
B	-1	SER	-	expression tag	UNP B1UXN2
B	0	HIS	-	expression tag	UNP B1UXN2

- Molecule 2 is 2-amino-8-sulfanyl-1,9-dihydro-6H-purin-6-one (CCD ID: B55) (formula: C₅H₅N₅OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			12	5	5	1	1		
2	B	1	Total	C	N	O	S	0	0
			12	5	5	1	1		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	42	Total 42	O 42	0	0
4	B	21	Total 21	O 21	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Dihydropteroate synthase



• Molecule 1: Dihydropteroate synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, α , β , γ	97.46Å 97.46Å 263.96Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.50 50.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-2.50) 99.9 (50.00-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.69 (at 2.40Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.224 , 0.269 0.222 , 0.267	Depositor DCC
R_{free} test set	1518 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	46.9	Xtriage
Anisotropy	0.922	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 61.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4219	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 45.69 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.2736e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: B55, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.39	14/2090 (0.7%)	1.37	20/2819 (0.7%)
1	B	1.18	6/2039 (0.3%)	1.25	6/2747 (0.2%)
All	All	1.29	20/4129 (0.5%)	1.31	26/5566 (0.5%)

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	113	GLU	C-N	11.20	1.49	1.33
1	A	113	GLU	C-O	10.06	1.36	1.24
1	B	43	ASP	CG-OD2	8.89	1.42	1.25
1	A	84	VAL	CA-CB	8.77	1.59	1.54
1	A	43	ASP	CG-OD1	7.88	1.40	1.25
1	A	114	ALA	C-N	7.00	1.44	1.33
1	A	93	GLU	CD-OE1	6.74	1.38	1.25
1	A	92	LYS	C-N	6.48	1.43	1.33
1	A	43	ASP	CG-OD2	5.86	1.36	1.25
1	A	240	ALA	CA-CB	5.83	1.62	1.53
1	B	43	ASP	CG-OD1	5.72	1.36	1.25
1	A	93	GLU	CD-OE2	5.66	1.36	1.25
1	B	65	GLU	CA-CB	5.64	1.61	1.53
1	A	176	VAL	CA-CB	5.61	1.60	1.54
1	B	43	ASP	CA-C	5.38	1.60	1.52
1	B	256	HIS	N-CA	-5.32	1.39	1.46
1	B	43	ASP	N-CA	5.30	1.52	1.46
1	A	265	ALA	CA-CB	-5.12	1.45	1.53
1	A	176	VAL	CA-C	5.11	1.58	1.52
1	A	84	VAL	CA-C	5.05	1.57	1.52

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	250	CYS	CA-CB-SG	-7.61	96.90	114.40
1	A	250	CYS	CA-CB-SG	-7.12	98.02	114.40
1	A	114	ALA	CA-C-N	-6.87	113.80	123.08
1	A	114	ALA	C-N-CA	-6.87	113.80	123.08
1	B	86	MET	N-CA-C	6.81	118.78	111.36
1	A	164	LEU	N-CA-C	-6.75	103.85	111.14
1	A	76	VAL	N-CA-C	-6.39	105.60	111.67
1	A	221	SER	N-CA-C	6.32	118.98	111.33
1	A	118	ILE	CA-C-N	-6.28	113.92	122.91
1	A	118	ILE	C-N-CA	-6.28	113.92	122.91
1	B	40	ASN	N-CA-C	6.21	117.72	111.07
1	A	154	ARG	N-CA-C	-6.13	104.51	111.07
1	A	86	MET	N-CA-C	5.90	117.79	111.36
1	A	92	LYS	CA-C-N	-5.90	111.72	121.92
1	A	92	LYS	C-N-CA	-5.90	111.72	121.92
1	B	75	SER	N-CA-C	5.89	117.50	110.19
1	B	84	VAL	CA-C-N	-5.79	112.91	119.28
1	B	84	VAL	C-N-CA	-5.79	112.91	119.28
1	A	107	VAL	N-CA-C	-5.71	104.94	110.42
1	A	112	ILE	CB-CA-C	-5.64	104.53	112.14
1	A	196	ASN	N-CA-C	5.61	117.84	111.11
1	A	152	ASN	N-CA-C	-5.38	99.74	108.41
1	A	64	GLY	N-CA-C	5.38	125.93	113.18
1	A	113	GLU	CA-C-N	-5.22	113.58	122.56
1	A	113	GLU	C-N-CA	-5.22	113.58	122.56
1	A	255	VAL	N-CA-C	5.12	117.40	108.95

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2060	0	2057	38	1
1	B	2012	0	2024	39	0
2	A	12	0	5	0	0
2	B	12	0	5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	25	0	0	0	0
3	B	35	0	0	1	0
4	A	42	0	0	1	0
4	B	21	0	0	0	0
All	All	4219	0	4091	77	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:108:ALA:O	1:A:112:ILE:HD12	1.44	1.14
1:B:104:LYS:HA	1:B:128:GLU:OE1	1.56	1.02
1:B:155:ASN:HD22	1:B:155:ASN:C	1.67	0.97
1:A:87:ILE:O	1:A:91:SER:HB2	1.76	0.86
1:A:157:MET:SD	1:A:160:MET:HE3	2.22	0.80
1:A:8:ARG:NH1	1:A:13:THR:OG1	2.20	0.74
1:B:104:LYS:CA	1:B:128:GLU:OE1	2.36	0.72
1:B:155:ASN:C	1:B:155:ASN:ND2	2.42	0.70
1:A:108:ALA:O	1:A:112:ILE:CD1	2.32	0.70
1:A:123:TRP:HB2	1:A:127:ALA:HB2	1.76	0.68
1:A:156:LEU:HG	1:A:160:MET:HE2	1.78	0.65
1:A:31:ASP:H	1:A:41:GLU:HG2	1.63	0.64
1:B:87:ILE:O	1:B:91:SER:HB2	2.01	0.61
1:B:76:VAL:O	1:B:80:ILE:HD12	2.00	0.60
1:A:123:TRP:CB	1:A:127:ALA:HB2	2.31	0.60
1:A:216:GLY:O	1:A:220:LYS:NZ	2.35	0.60
1:B:243:CYS:SG	1:B:264:MET:HE2	2.43	0.59
1:A:152:ASN:OD1	1:A:152:ASN:N	2.37	0.58
1:B:62:ILE:O	1:B:100:ILE:HD12	2.04	0.57
1:A:185:PRO:HG3	1:A:203:LEU:HD21	1.84	0.57
1:A:185:PRO:HB3	1:A:206:LEU:HD11	1.87	0.56
1:B:148:ARG:HD3	1:B:150:ASN:ND2	2.19	0.56
1:B:145:MET:HA	1:B:184:ASP:HB3	1.88	0.56
1:B:189:PHE:O	1:B:191:LYS:HG3	2.07	0.55
1:A:227:LEU:HB3	1:A:229:LEU:HD12	1.89	0.54
1:B:80:ILE:HG12	1:B:110:GLN:HG3	1.90	0.53
1:A:75:SER:OG	1:A:76:VAL:N	2.40	0.52
1:B:19:LYS:HE3	1:B:58:HIS:NE2	2.24	0.52
1:B:51:GLU:O	1:B:55:GLU:HG3	2.10	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:161:ILE:HG23	1:B:209:LEU:HD21	1.93	0.51
1:B:214:LEU:C	1:B:214:LEU:HD23	2.36	0.51
1:A:31:ASP:OD2	1:A:41:GLU:HG3	2.11	0.50
1:B:99:SER:HB3	1:B:118:ILE:HB	1.93	0.50
1:A:155:ASN:HD22	1:A:158:ALA:H	1.60	0.50
1:B:26:LEU:HD21	1:B:45:ALA:HB1	1.92	0.50
1:A:194:GLU:HG2	4:A:319:HOH:O	2.11	0.49
1:A:157:MET:SD	1:A:160:MET:CE	2.96	0.49
1:B:53:ARG:NH1	1:B:54:ASP:OD1	2.33	0.49
1:B:155:ASN:ND2	1:B:158:ALA:H	2.10	0.49
1:B:218:SER:HB2	1:B:255:VAL:HA	1.95	0.48
1:A:138:TYR:O	1:A:139:ASP:HB3	2.11	0.48
1:A:218:SER:HB2	1:A:255:VAL:HA	1.95	0.48
1:B:28:VAL:O	1:B:28:VAL:HG13	2.13	0.48
1:B:99:SER:CB	1:B:118:ILE:HB	2.44	0.48
1:B:86:MET:O	1:B:90:VAL:HG23	2.15	0.47
1:A:156:LEU:CG	1:A:160:MET:HE2	2.44	0.47
1:A:8:ARG:HA	1:A:8:ARG:HD2	1.71	0.47
1:B:117:HIS:O	1:B:140:VAL:HB	2.14	0.47
1:A:156:LEU:HG	1:A:160:MET:CE	2.45	0.46
1:A:145:MET:HA	1:A:184:ASP:HB3	1.98	0.46
1:B:244:LEU:HD11	1:B:248:LYS:HE2	1.98	0.46
1:B:203:LEU:HD11	1:B:215:LEU:HD13	1.97	0.45
1:A:145:MET:HG3	1:A:184:ASP:CG	2.41	0.45
1:B:236:GLU:H	1:B:236:GLU:CD	2.25	0.45
1:B:185:PRO:HB3	1:B:206:LEU:HD11	1.97	0.45
1:A:155:ASN:O	1:A:156:LEU:C	2.60	0.45
1:A:160:MET:HE1	1:A:199:ALA:HB2	1.99	0.45
1:B:165:TYR:CE1	1:B:209:LEU:HD22	2.52	0.45
1:A:106:GLU:HA	1:A:106:GLU:OE1	2.16	0.44
1:A:112:ILE:HD12	1:A:112:ILE:H	1.82	0.44
1:A:177:ARG:O	1:A:178:ASP:C	2.58	0.44
1:B:155:ASN:HD22	1:B:156:LEU:N	2.12	0.44
1:B:215:LEU:HG	1:B:217:THR:HG23	2.00	0.43
1:A:244:LEU:HD11	1:A:248:LYS:HE2	2.00	0.43
1:A:79:GLU:OE1	1:A:104:LYS:HD2	2.19	0.42
1:A:43:ASP:O	1:A:47:ARG:HG3	2.19	0.42
1:B:201:ARG:NE	3:B:283:SO4:O4	2.30	0.42
1:A:79:GLU:O	1:A:83:VAL:HB	2.19	0.42
1:B:154:ARG:HE	1:B:154:ARG:HB3	1.66	0.42
1:A:100:ILE:O	1:A:100:ILE:HG23	2.18	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:8:ARG:HH11	1:B:8:ARG:HA	1.85	0.41
1:B:11:GLU:OE1	1:B:95:LYS:NZ	2.54	0.41
1:B:95:LYS:HB3	1:B:95:LYS:HE3	1.67	0.41
1:A:100:ILE:HG22	1:A:118:ILE:O	2.20	0.41
1:A:180:ASN:HD22	1:A:180:ASN:HA	1.68	0.40
1:B:227:LEU:HB3	1:B:229:LEU:HD12	2.03	0.40
1:B:20:THR:HG21	1:B:246:ILE:HD11	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:HIS:CD2	1:A:137:HIS:NE2[7_465]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	263/297 (89%)	246 (94%)	16 (6%)	1 (0%)	30	49
1	B	254/297 (86%)	244 (96%)	10 (4%)	0	100	100
All	All	517/594 (87%)	490 (95%)	26 (5%)	1 (0%)	43	63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	66	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	219/247 (89%)	201 (92%)	18 (8%)	10	23
1	B	214/247 (87%)	198 (92%)	16 (8%)	12	26
All	All	433/494 (88%)	399 (92%)	34 (8%)	11	24

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	LEU
1	A	26	LEU
1	A	28	VAL
1	A	31	ASP
1	A	33	PHE
1	A	41	GLU
1	A	75	SER
1	A	83	VAL
1	A	91	SER
1	A	95	LYS
1	A	100	ILE
1	A	110	GLN
1	A	131	ILE
1	A	137	HIS
1	A	152	ASN
1	A	154	ARG
1	A	180	ASN
1	A	235	LEU
1	B	2	LYS
1	B	7	LEU
1	B	13	THR
1	B	27	ASN
1	B	41	GLU
1	B	74	VAL
1	B	78	GLU
1	B	80	ILE
1	B	82	ARG
1	B	84	VAL
1	B	152	ASN
1	B	154	ARG
1	B	155	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	180	ASN
1	B	194	GLU
1	B	209	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	150	ASN
1	A	155	ASN
1	A	180	ASN
1	B	88	GLN
1	B	155	ASN
1	B	180	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	B	284	-	4,4,4	0.39	0	6,6,6	0.77	0
3	SO4	B	279	-	4,4,4	0.59	0	6,6,6	0.60	0
3	SO4	B	281	-	4,4,4	0.27	0	6,6,6	0.81	0
3	SO4	A	279	-	4,4,4	0.46	0	6,6,6	0.46	0
3	SO4	B	282	-	4,4,4	0.54	0	6,6,6	0.61	0
2	B55	B	902	-	12,13,13	2.28	3 (25%)	13,19,19	2.05	3 (23%)
3	SO4	B	278	-	4,4,4	0.68	0	6,6,6	0.97	0
3	SO4	B	280	-	4,4,4	0.43	0	6,6,6	0.59	0
3	SO4	A	280	-	4,4,4	0.45	0	6,6,6	0.52	0
2	B55	A	901	-	12,13,13	2.30	2 (16%)	13,19,19	1.86	3 (23%)
3	SO4	A	282	-	4,4,4	0.59	0	6,6,6	0.73	0
3	SO4	B	283	-	4,4,4	0.68	0	6,6,6	0.50	0
3	SO4	A	281	-	4,4,4	0.48	0	6,6,6	0.39	0
3	SO4	A	278	-	4,4,4	0.76	0	6,6,6	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	B55	A	901	-	-	-	0/2/2/2
2	B55	B	902	-	-	-	0/2/2/2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	901	B55	O1-C5	6.65	1.36	1.23
2	B	902	B55	O1-C5	6.45	1.35	1.23
2	A	901	B55	C1-N1	2.79	1.35	1.31
2	B	902	B55	C2-C5	-2.47	1.35	1.44
2	B	902	B55	C2-N1	-2.42	1.33	1.38

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	902	B55	C1-N1-C2	-4.43	99.70	109.59
2	A	901	B55	C1-N1-C2	-4.05	100.53	109.59
2	B	902	B55	O1-C5-C2	-3.53	117.21	126.53
2	B	902	B55	O1-C5-N4	2.75	125.28	120.11
2	A	901	B55	O1-C5-C2	-2.35	120.32	126.53
2	A	901	B55	C5-C2-N1	2.15	132.33	129.81

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	283	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	267/297 (89%)	0.78	19 (7%) 22 19	56, 65, 86, 95	0
1	B	260/297 (87%)	0.79	17 (6%) 25 22	59, 69, 83, 93	0
All	All	527/594 (88%)	0.79	36 (6%) 23 20	56, 67, 84, 95	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	28	VAL	5.2
1	A	33	PHE	4.7
1	A	67	THR	4.5
1	A	28	VAL	4.5
1	B	66	SER	4.5
1	A	27	ASN	4.0
1	A	151	MET	3.5
1	A	66	SER	3.2
1	A	76	VAL	3.0
1	B	65	GLU	3.0
1	A	7	LEU	3.0
1	B	9	CYS	2.9
1	B	74	VAL	2.8
1	B	272	ILE	2.6
1	A	30	PRO	2.5
1	A	274	LYS	2.5
1	A	153	TYR	2.4
1	B	27	ASN	2.4
1	B	76	VAL	2.4
1	A	190	ALA	2.4
1	B	240	ALA	2.4
1	A	272	ILE	2.3
1	A	29	THR	2.3
1	A	35	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	39	TYR	2.3
1	A	2	LYS	2.2
1	B	262	SER	2.2
1	B	39	TYR	2.2
1	B	227	LEU	2.2
1	B	82	ARG	2.1
1	B	267	MET	2.1
1	A	9	CYS	2.1
1	A	225	HIS	2.1
1	B	112	ILE	2.1
1	B	106	GLU	2.0
1	B	235	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	B	282	5/5	0.51	0.15	100,105,106,106	0
3	SO4	B	283	5/5	0.53	0.17	97,101,102,103	0
3	SO4	A	282	5/5	0.59	0.15	91,92,94,97	0
3	SO4	B	278	5/5	0.59	0.19	96,96,98,100	0
3	SO4	A	281	5/5	0.62	0.16	92,94,95,96	0
3	SO4	B	284	5/5	0.63	0.15	90,90,91,95	0
3	SO4	B	280	5/5	0.64	0.16	100,101,101,101	0
3	SO4	A	280	5/5	0.65	0.15	89,89,90,93	0
3	SO4	B	281	5/5	0.71	0.15	89,89,91,92	0
2	B55	A	901	12/12	0.77	0.17	82,83,85,87	0
3	SO4	A	279	5/5	0.78	0.15	89,92,93,95	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	A	278	5/5	0.80	0.26	80,82,85,85	0
2	B55	B	902	12/12	0.84	0.13	84,85,85,88	0
3	SO4	B	279	5/5	0.87	0.14	82,83,87,87	0

6.5 Other polymers [i](#)

There are no such residues in this entry.